

Measuring Neutrino Flux at Low Energies: The Low ν Method (for $E_\nu > 0.5$ GeV)


Arie Bodek

University of Rochester

Plenary talk Nufact 2012

Thursday July 26. 2012

35min +5 min



$$Q^2 = -q^2 = -(k - k')^2 > 0,$$

$$s = (P + k)^2,$$

$$W^2 = (P + q)^2,$$

$$x = \frac{Q^2}{2P \cdot q},$$

$$y = \frac{P \cdot q}{P \cdot k},$$

$$\nu = \frac{P \cdot q}{M}.$$

ν = the energy of the final state hadrons in the lab (Ehad)

ν = the energy of the particles that recoil against the final state lepton (Erecoil)

References on low ν method.

For High Neutrino Energies ($E > 30$ GeV)

S.R. Mishra (CCFR), in Proceedings of the Workshop on Hadron Structure Functions and Parton Distributions, ed. by D. Geesaman et al.

W. Seligman, Ph.D. thesis (CCFR) Columbia, University, 1997, Nevis

For Intermediate Neutrino Energies ($E > 3.5$ GeV)

D. Battacharya, PhD. Thesis (MINOS), U. Pittsburgh (2009)

P. Adamson et al. (MINOS), Phys. Rev. **D 81**, 072002 (2010)

For Low Neutrino Neutrino Energies ($E > 0.4$ GeV)

A. Bodek, U. Sarica, D. Naples, L. Ren, “Methods to Determine Neutrino Flux at Low Energies: Investigation of the Low ν Method”. Eur.Phys.J. **C72**, 1973 (2012) arXiv:1201.3025 [hep-ex] (for $E_\nu > 0.7$ GeV)

A. Bodek, U. Sarica, K.S. Kuzmin, V.A. Naumov. “Extraction of Neutrino Flux with the Low ν Method at MiniBooNE Energies”. to be published in proceedings of CIPANP 2012, St. Petersburg (2012). arXiv:1207.1247 [hep-ex] (for $E_\nu > 0.4$ GeV)

The following two methods were used in the determination of the neutrino flux in high energy (E_ν) neutrino beams.

1. *All experiments use the Standard Technique which is to model the distribution of pions and kaons produced by incident proton beam in the target. Then, tracking the pions and kaons through the Horn focusing magnetic fields, and modeling the decays of pions and kaons in the decay pipe. THIS METHOD TYPICALLY HAS errors of 15%.*
2. *CCFR, NuTeV, MINOS also use the “low- v ” method for the determination of the energy dependence of the relative neutrino and antineutrino flux and normalize to a well known cross section at high energy. This method reduces the systematic errors.*

Here v is the energy transfer to the target (sometimes called E_{had})

- *CCFR and NuTeV use this method for $E_\nu > 30$ GeV.*
- *MINOS uses this method for $E_\nu > 3$ GeV for neutrinos and $E_\nu > 6$ GeV for antineutrinos. The method only yields the relative flux vs energy. Both experiments normalize to the world average total cross section at high energy.*

In this talk we investigate the use of the method down to $E_\nu = 0.4$ GeV which is in the MiniBooNE, MINERvA and T2K range of interest.

Are there other Techniques to determine flux?

The following is a list of all the techniques

1. Modeling the distribution of pions and kaons produced by incident proton beam in the target. Then, tracking the pions and kaons through the Horn focussing magnetic fields, and modeling the decays of pions and kaons in the decay pipe. Standard method
2. Measuring the muon flux that exits the decay pipe and relating it to the neutrino flux.
3. Monitoring Inverse muon decay events ($\nu_{\mu} + e \rightarrow \mu^{-} + \nu_e$) in the detector.
4. Monitoring neutrino-electron scattering events ($\nu_{\mu} + e \rightarrow \nu_{\mu} + e$) in the detector.
5. The “low- ν ” method for the determination of the energy dependence of the relative neutrino and antineutrino flux. Low nu method

1. In method 1, the differential cross sections for the production of pions and kaons by protons incident on a thick nuclear target must be known very well. In addition, the magnetic field of the horn focussing magnets must be modeled reliably.
2. In method 2, the response of the muon detectors at the end of the decay pipe must be very well understood (for absolute calibration of the neutrino flux). The response of the muon detectors is sensitive to δ rays. In addition, since the energy of the muons is not measured, it is difficult to determine the energy dependence of the neutrino flux.
3. In method 3, the threshold for the reaction $\nu_\mu + e \rightarrow \mu^- + \nu_e$ is about 12 GeV. Therefore, this method can only be used at higher energies. Unfortunately, this method cannot be used for the determination of the flux for antineutrinos. Inverse muon decay was used by NOMAD to constrain their neutrino flux at high energies.
4. In method 4, only the sum of the fluxes for neutrinos and antineutrinos can be measured. This is because calorimetric detectors such as MINERvA cannot determine the charge of final state electron in $\nu_\mu + e \rightarrow \nu_\mu + e$ events.

1. Modeling the distribution of pions and kaons

Standard method, errors of order 15% .

2. Measuring the muon flux that exits the decay pipe

Used at CERN, difficult technically

3. Monitoring Inverse muon decay events

Statistically limited, Good only for $E > 12$ GeV, not valid for antineutrinos

4. Monitoring neutrino-electron scattering events

Statistically limited,, yields sum of neutrino and antineutrino flux

The Low ν Method for High Energies

- At high E, cross section for DIS dominates.

$$\frac{d\sigma}{d\nu} = A \left(1 + \frac{B \nu}{A E} - \frac{C \nu^2}{A 2E^2} \right)$$

$$A = \frac{G_F^2 M}{\pi} \int_0^1 \mathcal{F}_2(x) dx,$$

$$B = -\frac{G_F^2 M}{\pi} \int_0^1 (\mathcal{F}_2(x) \mp x \mathcal{F}_3(x)) dx,$$

$$C = B - \frac{G_F^2 M}{\pi} \int_0^1 \mathcal{F}_2(x) \bar{R} dx,$$

Multiply both sides by flux \rightarrow

$$\frac{dN}{d\nu} = \phi A \left(1 + \frac{B \nu}{A E} - \frac{C \nu^2}{A 2E^2} \right) \text{ and}$$


$$\bar{R} = \left(\frac{1 + \frac{2Mx}{\nu}}{1 + R(x, Q^2)} - \frac{Mx}{\nu} - 1 \right).$$

As $\nu \rightarrow 0$,


$$\frac{dN}{d\nu} \approx \phi A$$

“A term” only depends on \mathcal{F}_2 , other terms are small at low ν . Correction terms proportional to ν/E depend on q_{bar} ($x\mathcal{F}_3$) and $R = \sigma_L/\sigma_T$ ($2x\mathcal{F}_1$)

Cross Sections per Nucleon for Neutrino on Carbon with ν Cut



Cross Sections per Nucleon for Antineutrino on Carbon with ν Cut






Used
In
MINOS

Our
study

The low ν cross section is approximately constant, with some model corrections. The method has been used by MINOS for neutrino energies greater than 3 GeV and antineutrino energies of 6 GeV


- The MINOS Collaboration suggested the number of low ν events should not exceed 60% of the total cross section
- However, this number should also be statistically significant.
- Hence, MINOS used $\nu < 1$ GeV for $E > 3$ GeV for neutrinos and $E > 5$ GeV for antineutrinos; $\nu < 2$ GeV and $\nu < 5$ GeV for $E > 9$ GeV and $E > 18$ GeV, respectively.





Total neutrino cross sections extracted by MINOS for neutrino energies $E_\nu > 3$ GeV using the low ν method

We investigate the uncertainties in the method for neutrino energies as low as $E_\nu = 0.4$ GeV.



Total antineutrino cross sections extracted by MINOS for antineutrino energies $E_\nu > 6$ GeV using the low ν method


We investigate the uncertainties in the method for antineutrino energies as low as $E_\nu = 0.7$ GeV.

Fraction of low ν events used to determine the flux must be less than 60% of the total number of events. The low ν cut must be less than 25% of the beam energy

$\nu < 0.5$ GeV means $Q^2 < 1$ GeV², 2/3 QE events, 1/3 Delta

$\nu < 0.2$ GeV means $Q^2 < 0.4$ GeV², all QE events

$\nu < 0.1$ GeV means $Q^2 < 0.2$ GeV², all QE events



Need to investigate uncertainties in modeling QE (and Delta production) processes on nuclear targets.

In Eur. Phys. J. C 72, 1973 (2012) we investigated:

$\nu < 0.25$ and $\nu < 0.5$ GeV for $E_\nu > 0.7$ GeV.

Here, we investigate $\nu < 0.2$ and $\nu < 0.1$ for $E_\nu > 0.4$ GeV (see also arXiv:1207.1247)

The Low ν Method for Low Energies

- The low ν cross section is approximately constant, so it can be used to determine the flux.
- The question is, what are the model uncertainties of the energy dependent corrections when we use the method at very low energies (between 0.4 GeV and 5 GeV).
- We need to use events with very low ν (i.e. less than 0.10 , 0.20, 0.25, and 0.5 GeV).
- We studied all of the above low ν regions.

The low ν for low neutrino energies.
At low ν the cross section is dominated by W_2

$$\begin{aligned} \frac{d\sigma^{ud}}{dQ^2 d\nu} &= S_{\cos} \frac{1}{2E^2} \mathcal{W}_1 [Q^2 + m_\mu^2] \\ &+ S_{\cos} \mathcal{W}_2 \left[\left(1 - \frac{\nu}{E}\right) - \frac{(Q^2 + m_\mu^2)}{4E^2} \right] \\ &+ S_{\cos} \mathcal{W}_3 \left[\frac{Q^2}{2ME} - \frac{\nu}{4E} \frac{Q^2 + m_\mu^2}{ME} \right] \\ &+ S_{\cos} \mathcal{W}_4 \left[m_\mu^2 \frac{(Q^2 + m_\mu^2)}{4M^2 E^2} \right] \\ &- S_{\cos} \mathcal{W}_5 \left[\frac{m_\mu^2}{ME} \right] \end{aligned}$$

$$\begin{aligned} \sigma_{\nu cut}(E) &= \int_{\nu_{min}(E)}^{\nu_{cut}} \frac{d^2\sigma}{dQ^2 d\nu} dQ^2 d\nu \\ &= \sigma_{W_2} + \sigma_2 + \sigma_1 + \sigma_3 + \sigma_4 + \sigma_5, \end{aligned}$$

**= constant plus correction terms
that depend on ν/E .**

$$\sigma_2 = S_{\cos} \int_{\nu_{min}(E)}^{\nu_{cut}} \left[-\frac{\nu}{E} - \frac{Q^2 + m_\mu^2}{4E^2} \right] \mathcal{W}_2 d\nu. \quad \text{known}$$

$$\sigma_1 = S_{\cos} \int_{\nu_{min}(E)}^{\nu_{cut}} - \left[\frac{(Q^2 + m_\mu^2)}{2E^2} \right] \mathcal{W}_1 d\nu.$$

$$\sigma_3 = S_{\cos} \int_{\nu_{min}(E)}^{\nu_{cut}} \left[\frac{Q^2}{2ME} - \frac{\nu}{4E} \frac{Q^2 + m_\mu^2}{ME} \right] \mathcal{W}_3 d\nu.$$

$$\sigma_4 = S_{\cos} \int_{\nu_{min}(E)}^{\nu_{cut}} \left[m_\mu^2 \frac{(Q^2 + m_\mu^2)}{4M^2 E^2} \right] \mathcal{W}_4 d\nu. \quad \text{Very small}$$

$$\sigma_5 = S_{\cos} \int_{\nu_{min}(E)}^{\nu_{cut}} \left[\frac{-m_\mu^2}{ME} \right] \mathcal{W}_5 d\nu. \quad \text{Very small}$$

$$\sigma_{tot}(E) = \sigma_{W_2}(\infty) [f_C]$$

$$f_C = [f_{W_2} + f_2 + f_1 + f_3 + f_4 + f_5]$$

$$f_{W_2} = \frac{\sigma_{W_2}}{\sigma_{W_2}(\infty)} (\approx 1)$$

$$f_2 = \frac{\sigma_2}{\sigma_{W_2}(\infty)} (= \textit{kinematic correction})$$

$$f_1 = \frac{\sigma_1}{\sigma_{W_2}(\infty)} (= \textit{important})$$

$$f_3 = \frac{\sigma_3}{\sigma_{W_2}(\infty)} (= \textit{important})$$

$$f_4 = \frac{\sigma_4}{\sigma_{W_2}(\infty)} (= \textit{very small})$$

$$f_5 = \frac{\sigma_4}{\sigma_{W_2}(\infty)} (= \textit{very small})$$

Correction terms that make the cross section deviate from a constant are proportional to v/E

$$\sigma_{\nu cut}(E) = \sigma_{W_2}(\infty) [f_C],$$

$$f_C = [f_{W_2} + f_2 + f_1 + f_3 + f_4 + f_5],$$

$$f_{W_2} = \frac{\sigma_{W_2}}{\sigma_{W_2}(\infty)} \approx 1,$$


$$f_i = \frac{\sigma_i}{\sigma_{W_2}(\infty)}.$$

1. $f_{W_2} = \frac{\sigma_{W_2}}{\sigma_{W_2}(\infty)} \approx 1$ is well known and does not contribute to the uncertainty in f_C .
2. The energy dependent correction f_2 is explicit and therefore does not contribute to the uncertainty in f_C .
3. The contributions of f_4 and f_5 are small since they are proportional to the square of the muon mass, and therefore have a negligible contribution to the uncertainty in f_C . (Note that the vector parts of f_4 and f_5 are known very well since they can be expressed in terms of the vector parts of W_1 and W_2).
4. The only non-negligible uncertainty originates from the modeling of the contributions of f_1 and f_3 (primarily from f_3).

Lowest ν cut in MINOS was 1 GeV (poor calorimeter). For MINERvA we can go to lower hadron energies. Start with a ν cut of 0.5 GeV

$$\dots \nu_{max} = 0.5: \bar{f}_C \dots$$


$\bar{f}_C(15.1)$ for Neutrino on Carbon $\nu < 0.5$



Relative low ν Neutrino cross section normalized to the cross section at 15.1 GeV

... $\nu_{max}=0.5: \bar{f}_C...$

$\bar{f}_C(15.1)$ for Antineutrino on Carbon $\nu < 0.5$




Relative low ν Anti-neutrino cross section normalized to the cross section at 15.1 GeV

Uncertainty in the relative low ν cross section for neutrinos (from uncertainties in QE and Delta production cross sections and form factors)

$$\dots \nu_{max} = 0.5 \dots$$


$\bar{F}_C(15.1)$ Error for Neutrino on Carbon $\nu < 0.5$, $W < 1.4$



Uncertainty in the relative low ν cross section for antineutrinos neutrinos from uncertainties in QE and Delta production cross sections and form factors

$$\dots \nu_{max} = 0.5 \dots$$

$\bar{F}_C(15.1)$ Error for Antineutrino on Carbon $\nu < 0.5$, $W < 1.4$



... $\nu_{max}=0.5$: Conclusion

The error in f_c is about than 2.6 % for neutrinos (above 1.2 GeV)

The error in f_c is about than 1.4 % for antineutrinos (above 2 GeV)

For these energies, the fraction of $\nu < 0.5$ GeV events is less than 60%).

For lower energies need to use ν cuts less than 0.5 GeV.

Note the number of event is proportional to the ν cut, so the number of events with $\nu < 0.5$ GeV is twice the number of $\nu < 0.25$ GeV).

Next: investigate $\nu < 0.2$ and $\nu < 0.1$ GeV cuts.

$$W_{1-Qelastic}^{\nu-vector} = \delta(\nu - \frac{Q^2}{2M})\tau |G_M^V(Q^2)|^2$$

$$W_{1-Qelastic}^{\nu-axial} = \delta(\nu - \frac{Q^2}{2M})(1 + \tau) |F_A(Q^2)|^2$$

$$W_{2-Qelastic}^{\nu-vector} = \delta(\nu - \frac{Q^2}{2M}) |F_V(Q^2)|^2$$

$$W_{2-Qelastic}^{\nu-axial} = \delta(\nu - \frac{Q^2}{2M}) |F_A(Q^2)|^2$$

$$W_{3-Qelastic}^{\nu} = \delta(\nu - \frac{Q^2}{2M}) |2G_M^V(Q^2)F_A(Q^2)|^2$$

where

$$G_E^V(Q^2) = G_E^p(Q^2) - G_E^n(Q^2),$$

$$G_M^V(Q^2) = G_M^p(Q^2) - G_M^n(Q^2).$$

and

$$|F_V(Q^2)|^2 = \frac{[G_E^V(Q^2)]^2 + \tau [G_M^V(Q^2)]^2}{1 + \tau}.$$

For $\nu < 0.25$ GeV , the uncertainties in the relative low ν cross section depends on uncertainties in the modeling of the QE cross section.





We investigate several models that fit either the MiniBooNE cross section or the free nucleon cross sections:

1. Free nucleon form factors with M_A varying between 1.014 and 1.3 GeV and a **dipole fit form** for F_A .
2. **Transverse enhancement of GMp and GMn** as observed in electron scattering on bound nucleons, and a **dipole form for F_A**
3. Transverse enhancement of GMp and GMn as observed in electron scattering on bound nucleons, and a **modified dipole form for F_A** fit to all free nucleon data (our best model).

We use BBBA form factors. Duality based parametrization of the ratio of free EM nucleon form factors to dipole (G_{ep} , G_{mp} , G_{en} , G_{mn}).

A. Bodek, S. Avvakumov, R. Bradford, and H. Budd: Vector and Axial Nucleon Form Factors:

Phys. J. C53, 349 (2008).



1. Ratios of $G_{E\nu}$ (a), $G_{M\nu}/\mu_\nu$ (b), G_{En} (c) and G_{Mn}/μ_n (d) to G_D . The short-dashed line in each plot is the old Kelly

BBBA form factors are also available for the axial form factor. Duality base parametrization of the ratio of free nucleon F_A to dipole with $M_A = 1.015$.

A. Bodek, S. Avvakumov, R. Bradford, and H. Budd: Vector and Axial Nucleon Form Factors: *Phys. J. C53*, 349 (2008).




Fig. 3. (a) $F_A(Q^2)$ re-extracted from neutrino-deuteron data divided by $G_D^A(Q^2)$ [22]. (b) $F_A(Q^2)$ from pion electroproduction divided by $G_D^A(Q^2)$ [22], corrected for hadronic effects[12]. Solid line - duality based fit; Short-dashed line - $F_A(Q^2)_{A2=V2}$

Transverse Enhancement Carbon 12




Fig. 7. The transverse enhancement ratio[14] (\mathcal{R}_T) as a function of Q^2 . Here, \mathcal{R}_T is ratio of the integrated transverse response function for QE electron scattering on nucleons bound in carbon divided by the integrated response function for independent nucleons. The black points are extracted from Carlson *et al*[16], and the blue bands are extracted from a fit[19] to QE data from the JUPITER[18] experiment (Jlab experiment E04-001). The curve is a fit to the data of the form $\mathcal{R}_T = 1 + AQ^2e^{-Q^2/B}$. The dashed lines are the upper and lower error bands (color online).


Nuclear Corrections from electron scattering data (BBC Parametrization of transverse enhancement of form factors in Carbon)

A. Bodek, H. Budd and E. Christy, *Eur.Phys. J. C*71, 172 (2011)


Use RT factor to enhance GMp and GMn in bound nucleons extracted from in electron scattering experiments on Carbon.

We will refer to TE cross sections with this correction as BBC form factors for a nucleus. (i.e. BBBA times RT).

Fraction of $\nu < 0.20$ Cross Section to Total QE Cross Section for Neutrino




Fraction of $\nu < 0.20$ Cross Section to Total QE Cross Section for Antineutrino



Want fraction of events in the low ν flux sample to be less than 60% of the QE cross section²⁵


Fraction of $\nu < 0.10$ Cross Section to Total QE Cross Section for Neutrino



Neutrinos:
Can use $\nu < 0.1$ GeV for
 $E > 0.4$ GeV

Fraction of $\nu < 0.10$ Cross Section to Total QE Cross Section for Antineutrino


$\nu < 0.1$ GEV



AntiNeutrinos:
Can use $\nu < 0.1$ GeV for
 $E > 0.6$ GeV


GENIE cross section has no TE

Want fraction of events in the low ν flux sample to be less than 60% of the QE cross section²⁶




$\nu < 0.2$ GeV
NEUTRINOS

BBC (TE) form factors
with modified dipole
 F_A ($M_A = 1.014$)



Normalize to the
low ν cross
section at $E = 1.1$
GeV and look at
the relative low ν
cross section to
get the correction
factor.

$\nu < 0.2$ can only
be used down to
 $E_\nu = 0.7$ GeV.




To go to lower energy need a lower ν cut.



$\nu < 0.1$ GeV Neutrinos



BBC (TE) form
Factors with
modified dipole F_A
($M_A = 1.014$)

Normalize to the
low ν cross section
at $E_\nu = 1.1$ GeV and
look at the relative
low ν cross section
to get the
correction factor.




$\nu < 0.1$ GeV can be
used down to
 $E_\nu = 0.4$ GeV.






Error in f_C for Antineutrino ($\nu < 0.10$)



Uncertainty in correction factor is independent of M_A . Also independent of the uncertainties in the TE parameters. Difference between with and without TE is a maximum of 0.035.

Error in f_C for Antineutrino ($\nu < 0.20$)



Conclusion 1

- The energy dependence of the flux in low energy neutrino beams can be determined by using the low ν events with

$\nu < 0.2$ GeV for $E_\nu > 0.7$ GeV (for neutrino beams)

$\nu < 0.1$ GeV for $E_\nu > 0.4$ GeV (for neutrino beams)

$\nu < 0.2$ GeV for $E_\nu > 2$ GeV (for antineutrino beams)

$\nu < 0.1$ GeV for $E_\nu > 0.7$ GeV (for antineutrino beams)

The uncertainty in the model dependent correction factors is small so any model which fits the data can be used.

(note the absolute flux cannot be determined with the low ν method, so it must be normalized at some energy).

- The primary error will be experimental, i.e. statistical and systematic experimental error in the extracted number of low ν events as a function of neutrino energy, resolution smearing and unfolding.

Example: Application to *MiniBooNE* *see* arXiv:1207.1247

MiniBooNE reconstructs E_ν^{QE} by Assuming QE kinematics for the muon.

We extract $\nu = E_\nu^{\text{QE}} - E_\mu$



$$E_\nu^{\text{QE}} = \frac{2(M'_n)E_\mu - ((M'_n)^2 + m_\mu^2 - M_p^2)}{2 \cdot [(M'_n) - E_\mu + \sqrt{E_\mu^2 - m_\mu^2} \cos \theta_\mu]}, \quad (1)$$

$$Q_{QE}^2 = -m_\mu^2 + 2E_\nu^{\text{QE}}(E_\mu - \sqrt{E_\mu^2 - m_\mu^2} \cos \theta_\mu), \quad (2)$$


where $E_\mu = T_\mu + m_\mu$ is the total muon energy and M_n , M_p , m_μ are the neutron, proton, and muon masses. The adjusted neutron mass, $M'_n = M_n - E_B$, depends on the binding energy (or more carefully stated, the separation energy) in carbon, E_B , which for this analysis is set to 34 ± 9 MeV.

MiniBooNE removes pion production events and subtracts non QE background (e.g. Delta)

MiniBooNE flux and estimated fractional error




MiniBooNE published flux weighted cross section in $\text{Cos}\theta_\mu$ and Muon Kinetic, T_μ bins. We need to the data binned in $E_{\nu^{\text{QE}}}$ and ν .



$M_A = 1.30$


Ratio of data to model with $M_A = 1.3$ (Plot from MiniBooNE paper)



We use average of the ratio of data to model for bins that fall below $\nu < 0.1$ and $\nu < 0.2$ in bins of E_{ν}^{QE} to extract the ratio of the flux from low ν events to published MiniBooNE flux.


The MiniBooNE binning of 0.1 in $\text{Cos}\theta_{\mu}$ is not ideal for this study. Bins of 0.05 in $\text{Cos}\theta_{\mu}$ would be better, or better yet bins in E_{ν}^{QE} and ν . We use the published data as is as an example. ³⁶

Simple test: Statistical errors only. No unfolding for resolution in E_{ν}^{QE}



Ratio of low ν flux to flux used by MiniBooNE. Ratio at 1.0 GeV is defined as 1.0

MiniBooNE unfolded QE cross section



Conclusion 2, a low ν analysis with E_{ν}^{QE} and ν bins (and accounting for systematic errors and unfolding) can provide an independent check and reduce the uncertainty on the energy dependence of the measured QE cross section.


The extracted low ν flux is within the quoted systematic error in the flux.

Next we test the effect on M_A

Statistical errors only. No unfolding for experimental resolution in neutrino energy.

Testing the effect on M_A

Ratio of low ν flux to flux used by MiniBoNE.
Ratio at 1.0 GeV is defined as 1.0



Possible Flux correction fit A

Possible Flux correction B

Error on the published MiniBoone flux

Ratio of flux extracted from low ν events in *MiniBoone* to the published flux is within the published *MiniBoone* systematic errors on the flux. Next we test the effect of possible flux modifications (A or B) which are consistent with the low ν flux on the extraction of M_A

Fit to the double differential ($d^2\sigma/d\cos\theta_\mu dE_\mu$) cross sections (137 points) (2D)

M_A (in GeV)	N	χ^2/NDF
BBBA(07), dipole F_A		
1.348 \pm 0.018	0.994 \pm 0.009	39.798/135=0.295 (published flux)
1.347 \pm 0.018	0.994 \pm 0.009	42.709/135=0.316 (Flux correction A)
1.340 \pm 0.018	0.996 \pm 0.009	39.360/135=0.292 (Flux correction B)
BBBA(07), modified F_A (our best estimate of free nucleon form factors)		
1.299 \pm 0.016	0.994 \pm 0.009	30.434/135=0.225 (published flux)
1.288 \pm 0.016	0.991 \pm 0.009	42.939/135=0.318 (Flux correction A)
1.283 \pm 0.016	0.993 \pm 0.009	37.666/135=0.279 (Flux correction B)
BBC(11)(TE), dipole F_A		
1.219 \pm 0.018	1.013 \pm 0.009	43.683/135=0.324 (published flux)
1.217 \pm 0.018	1.011 \pm 0.009	40.700/135=0.302 (Flux correction A)
1.206 \pm 0.018	1.012 \pm 0.009	44.562/135=0.330 (Flux correction B)
BBC(11) (TE), modified F_A (our best nuclear model based on electron scattering data)		
1.174 \pm 0.017	1.010 \pm 0.009	37.190/135=0.276 (published flux)
1.171 \pm 0.016	1.008 \pm 0.009	35.453/135=0.263 (Flux correction A)
1.162 \pm 0.016	1.009 \pm 0.009	38.307/135=0.284 (Flux correction B)

We let the absolute normalization float in the fit

Conclusion 3: Possible low v flux modification do not change the extracted value of M_A . Expected since flux changes the energy dependence of cross section but not the Q^2 distribution

Conclusion 4: The extracted values of M_A with BBC (Transverse Enhancement and modified dipole) from the 2D fits is 1.17 with similar χ^2/DF as all other fits

With our best model, the MiniBooNE 2D data favors TE with modified dipole F_A , and $M_A=1.17$

Fit to the differential ($d\sigma/dQ^2$) cross sections (17 data points) (1D).

M_A (in GeV)	N	χ^2/NDF
BBBA(07), dipole F_A		
1.410 \pm 0.040	0.993 \pm 0.020	11.764/15=0.784 (published flux)
1.403 \pm 0.040	0.991 \pm 0.020	12.045/15=0.803 (Flux correction A)
1.388 \pm 0.039	0.991 \pm 0.020	12.222/15=0.815 (Flux correction B)
BBBA(07), modified F_A (our best estimate of free nucleon form factors)		
1.330 \pm 0.036	0.986 \pm 0.021	5.611/15=0.374 (published flux)
1.324 \pm 0.036	0.984 \pm 0.021	5.674/15=0.378 (Flux correction A)
1.311 \pm 0.035	0.984 \pm 0.021	5.896/15=0.393 (Flux correction B)
BBC(11), dipole F_A		
1.170 \pm 0.034	1.050 \pm 0.022	19.714/15=1.314 (published flux)
1.165 \pm 0.034	1.048 \pm 0.022	18.537/15=1.236 (Flux correction A)
1.152 \pm 0.033	1.047 \pm 0.022	18.516/15=1.234 (Flux correction B)
BBC(11) (TE) , modified F_A (our best nuclear model based on electron data)		
1.110 \pm 0.031	1.039 \pm 0.023	19.042/15=1.270 (published flux)
1.106 \pm 0.031	1.037 \pm 0.023	17.766/15=1.184 (Flux correction A)
1.095 \pm 0.030	1.036 \pm 0.023	17.711/15=1.181 (Flux correction B)



Here also, possible low ν flux modification do not change the extracted value of M_A . Expected since flux changes the energy dependence of cross section but not the Q^2 distribution

Conclusion 5: The extracted values of M_A with BBC (Transverse Enhancement and modified dipole) for the 1D fit is 1.10 with similar χ^2/DF as all other fits. With our best model, the MiniBooNE 1D Q^2 distribution favors TE with modified dipole F_A and $M_A = 1.10$

We let the absolute normalization float 1D fits to MiniBooNE Q^2 distribution

Next: compare $M_A = 1.17$ and $M_A = 1.10$ with modified dipole with free nucleon data.

Ratio of F_A data on deuterium to dipole with $M_A = 1.015$ GeV (solid line=modified dipole)



Orange=modified dipole with $M_A = 1.17$, Green = modified dipole with $M_A = 1.10$

Conclusion 6:

Modified dipole with $M_A = 1.10$ is consistent with pion electroproduction data.

It is also consistent with neutrino data on deuterium for $Q^2 < 0.5$ GeV².

It is not consistent with the average of the neutrino deuterium data for $Q^2 > 1$ GeV.

Final Conclusion

- We can greatly reduce the systematic errors in the neutrino cross section for energies greater than 0.4 GeV by using the low ν method.
- The low ν method must be normalized to a cross section at some energy. MINERvA can extend the cross section measurements to overlap both MINOS and MiniBooNE, thus normalizing to known cross sections at high energies.