

The EMC Effect of Tritium and Helium-3 from the JLab MARATHON Experiment

Michael Nycz

On behalf of the MARATHON Collaboration



JLUO Annual Meeting, Jefferson Lab
June 25, 2026


Outline

- Deep Inelastic Scattering and the EMC effect
- MARATHON Experiment and $A=3$ mirror nuclei
- MARATHON EMC Results
- Summary and Outlook

More details found in

[D. Abrams et al., Phys. Rev. Lett. 135 \(2025\)](#)


Deep Inelastic Scattering



$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'}{Q^4} \left[\frac{2F_1(x)}{M} \sin^2\left(\frac{\theta}{2}\right) + \frac{F_2(x)}{\nu} \cos^2\left(\frac{\theta}{2}\right) \right]$$

$$F_1 = \frac{1}{2} \sum_i e_i^2 q_i(x) \quad F_2 = x \sum_i e_i^2 q_i(x)$$

Deep Inelastic Scattering



$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'}{Q^4} \left[\frac{2F_1(x)}{M} \sin^2\left(\frac{\theta}{2}\right) + \frac{F_2(x)}{\nu} \cos^2\left(\frac{\theta}{2}\right) \right]$$


$$F_1 = \frac{1}{2} \sum_i e_i^2 q_i(x) \qquad F_2 = x \sum_i e_i^2 q_i(x)$$

Absorption cross section of **longitudinal** or **transverse** photons

$$R = \frac{\sigma_l}{\sigma_t} \qquad F_1 = \frac{MF_2(\nu^2 + Q^2)}{Q^2\nu(1 + R)}$$

$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'}{Q^4} \cos^2\left(\frac{\theta}{2}\right) F_2(x) \left[\frac{1}{\nu} + \frac{1 + \frac{Q^2}{\nu^2}}{xM(1 + R) \tan^2\left(\frac{\theta}{2}\right)} \right]$$

Deep Inelastic Scattering




$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'}{Q^4} \cos^2\left(\frac{\theta}{2}\right) F_2(x) \left[\frac{1}{\nu} + \frac{1 + \frac{Q^2}{\nu^2}}{xM(1+R) \tan^2\left(\frac{\theta}{2}\right)} \right]$$

Ratio of structure functions

$$\frac{\sigma_{A1}}{\sigma_{A2}} = \frac{F_2^{A1}}{F_2^{A2}}$$


Expectation from Scaling and Nucleon Fermi Motion

$$AF_2^A = ZF_2^p + (A - Z)F_n^n$$




EMC Effect

$$AF_2^A = ZF_2^p + (A - Z)F_n^n$$




quark momentum distributions in nuclei are not simply the sum of free proton and neutron distributions




[J. J. Aubert et al., Phys. Lett. B 123, 275 \(1983\)](#)

EMC Effect: Confirmation




[A. Bodek et al., Phys. Rev. Lett. 50, 1431 \(1983\)](#)


EMC Effect



[A. Bodek et al., Phys. Rev. Lett. 50, 1431 \(1983\)](#)




[J. Gomez et al., Phys. Rev. D49 \(1994\)](#)




[J. Seely et al., Phys. Rev. Lett. 103, \(2009\)](#)


EMC Effect



[A. Bodek et al., Phys. Rev. Lett. 50, 1431 \(1983\)](#)




[J. Gomez et al., Phys. Rev. D49 \(1994\)](#)




[J. Seely et al., Phys. Rev. Lett. 103, \(2009\)](#)

EMC Effect: Models and Predictions


- 1. Multiquark clusters**
- 2. Dynamical Rescaling**
 - a. $F_2^A(x, Q^2) = F_2^N(x, \xi_A(Q^2) \cdot Q^2)$
- 3. Off-shell correction**
 - a. $\delta f(x)$
- 4. And many, many more.....**



A. Bodek et al., Phys. Rev. Lett. 50, 1431 (1983)





J. Gomez et al., Phys. Rev. D49 (1994)




J. Seely et al., Phys. Rev. Lett. 103, (2009)

EMC-SRC Correlation




[O. Hen et al., Phys. Rev. C 85 \(2012\) 047301](#)



[N Fomin, et al. Phys. Rev. Lett \(2012\)](#)

40+ Years: No Consensus




MARATHON


MeAsurement of the F_2^n/F_2^p , d/u RAtioS and A=3 EMC Effect in Deep Inelastic Electron Scattering Off the Tritium and Helium MirrOr Nuclei.

Jefferson Lab PAC37 Proposal, December 2010

The JLab MARATHON Collaboration




1. Ratio of mirror nuclei provides a novel method of the F_2^n/F_2^p ratio at large x
2. Measurement of EMC of light nuclei (A=3)



[A. Accardi et al. Phys. Rev. D 84 \(2011\)](#)

The MARATHON Experiment

Hall A



Data collection: Jan. - April 2018


- LHRS and RHRS independently to detect e^-
- Standard detector configurations

- Sealed target cells
 - Cooled to 40 K
- ^3H filled at Savannah River National Lab


Kin	E (GeV)	E' (GeV)	Q^2 [[GeV/c] ²]	θ (deg)	x
0	10.59	3.1	2.79	16.90	0.19
1	10.59	3.1	3.06	17.58	0.22
2	10.59	3.1	3.61	19.12	0.26
3	10.59	3.1	4.19	20.58	0.29
4	10.59	3.1	4.76	21.93	0.34
5	10.59	3.1	5.31	23.21	0.38
7	10.59	3.1	6.47	25.59	0.46
9	10.59	3.1	7.56	27.77	0.54
11	10.59	3.1	8.71	29.92	0.62
13	10.59	3.1	9.84	31.73	0.70
15	10.59	3.1	10.96	33.56	0.78
16	10.59	2.9	11.83	36.12	0.82



MARATHON Analysis




${}^3\text{H}$ Decay: $\tau = 4500 \pm 8$ days
 ${}^3\text{H} \rightarrow {}^3\text{He} + e^- + \bar{\nu}_e$




Charge Normalized Yield


$$Y = \frac{N_e C}{LTQ\rho L}$$




- Corrections
- ${}^3\text{H}$ Decay
 - Density
 - Radiative
 - Charge symmetric
 - Endcap
 - Bin centering



MARATHON Analysis




^3H Decay: $\tau = 4500 \pm 8 \text{ days}$
 $^3\text{H} \rightarrow ^3\text{He} + e^- + \bar{\nu}_e$




Charge Normalized Yield

$$Y = \frac{N_e C}{LTQ\rho L}$$



- Corrections
- ^3H Decay
 - Density
 - Radiative
 - Charge symmetric
 - Endcap
 - Bin centering



EMC Effect of ^3H and ^3He

Normalization

- F_2^n / F_2^p can be extracted from the ratios: σ_h / σ_d , σ_t / σ_d and σ_d / σ_p
- Around $x \approx 0.31$ nuclear corrections expected to be minimal
 - F_2^n / F_2^p should be mutually consistent

$$F_2^n / F_2^p = \frac{2R_{hd}(\sigma_d / \sigma_h) - 1}{1 - R_{hd}(\sigma_d / \sigma_h)} = \frac{2R_{td}(\sigma_d / \sigma_t) - 1}{1 - R_{td}(\sigma_d / \sigma_t)}$$


1. Data
2. Model Input

- K-P Model
- Data-driven
 - Describes very well all available DIS nucleon and nuclear world data


A comparison among these extractions found:
 ^3H & ^3He EMC ratios needed to be normalized to be consistent with σ_d / σ_p extraction

σ_t / σ_d Normalization factor: 0.996 ± 0.005	σ_h / σ_d Normalization factor: 1.021 ± 0.005
--	--


Normalized A=3 EMC Ratios




Isoscalar Correction: Neutron & Proton Excess




${}^3\text{H}$



${}^2\text{H}$






${}^3\text{He}$



${}^2\text{H}$

Isoscalar Correction: Neutron & Proton Excess



$$F_{iso} = \frac{A(1 + \sigma_n/\sigma_p)}{2[Z + (A - Z)\sigma_n/\sigma_p]}$$



[D. Abrams et al., Phys. Rev. Lett. 128 \(2022\)](#)


Normalized A=3 EMC Ratios

First measurement




- Systematic Uncertainties for EMC Ratios**
- Target density due beam-heating: (0.1-0.5)%
 - Radiative corrections: (0.25-0.45)%
 - Acceptance: 0.2%

Normalized A=3 EMC Ratios



World data after applying Marathon's Isoscalar correction to existing ${}^3\text{He}$ data



- Systematic Uncertainties for EMC Ratios**
- Target density due beam-heating: (0.1-0.5)%
 - Radiative corrections: (0.25-0.45)%
 - Acceptance: 0.2%

Summary and Outlook

PHYSICAL REVIEW LETTERS **135**, 062502 (2025)

Editors' Suggestion

EMC Effect of Tritium and Helium-3 from the JLab MARATHON Experiment

- The MARATHON reported the measurement of the EMC effect of the $A=3$ mirror nuclei
 - [D. Abrams et al., Phys. Rev. Lett. 135 \(2025\)](#)
 - First measurement of the EMC effect of ${}^3\text{H}$
- $A=3$ data will provide
 - Unique input for QCD studies of the nucleons
 - Partonic structure of the few-body nuclear systems in the valence quark region
- The Hall A tritium program has proven to be a great success
 - More than 10 PhDs
 - PRL and Nature publications
 - A lot of interesting physics!
- Renewed interest in tritium
 - E12-20-05: Approved tritium experiment in Hall B

