

# Measurement of Deeply Virtual Compton Scattering on the Proton at Multiple Beam Energies with the CLAS12 Detector

**Yijie Wang, Volker Burkert, Latifa Elouadrhiri,  
Richard G. Milner, Bhawani Singh**

Run Group K



Massachusetts  
Institute of  
Technology



Mar, 2026

# Way to Understand Proton's Structure

## Generalized Parton Distribution

**GPDs** encode the correlated momentum and spatial structure of quarks and gluons inside hadrons. Each term is a complex function.

Measurable variables are Compton Form Factor:

$$\mathcal{F} = \int_{-1}^1 dx \left[ \frac{1}{x - \xi - i\epsilon} - \frac{1}{x + \xi - i\epsilon} \right] F(x, \xi, t)$$

CFF
GPD

With polarized electron beam and proton, we can have access to CFFs'  $\text{Im}(\mathcal{H})$  and  $\text{Re}(\mathcal{H})$ .

		Quark Polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$H$		$2\tilde{H}_T + E_T$
	L		$\tilde{H}$	$\tilde{E}_T$
	T	$E$	$\tilde{E}$	$H_T, \tilde{H}_T$

# Way to Understand Proton's Structure

## Gravitational Form Factor

- Mass/energy distribution
- Angular momentum distribution
- Shear forces and pressure distributions


$$\langle p' | T^{\mu\nu} | p \rangle = \bar{u}(p') \left[ A(t) \gamma^{(\mu} P^{\nu)} + B(t) \frac{i\sigma^{(\mu\alpha} \Delta_\alpha P^{\nu)}}{2M} + D(t) \frac{\Delta^\mu \Delta^\nu - g^{\mu\nu} \Delta^2}{4M} \right] u(p)$$

$$\int_{-1}^1 dx x H(x, \xi, t) = A(t) + \xi^2 D(t)$$

$$\int_{-1}^1 dx x E(x, \xi, t) = B(t) - \xi^2 D(t)$$

$$p(r) = \frac{1}{6M} \nabla^2 \widetilde{D}(r)$$


$$s(r) = -\frac{1}{4M} r \frac{d}{dr} \left( \frac{1}{r} \frac{d}{dr} \widetilde{D}(r) \right)$$



Burkert, V. D., Elouadrhiri, L., & Girod, F. X. (2018). The pressure distribution inside the proton. *Nature*, 557(7705), 396-399.


# Physics Processes

## Deeply Virtual Compton Scattering $ep \rightarrow ep\gamma$



perturbative QED

QCD



### Cross Section of BH-DVCS

$$\frac{d^5\sigma}{dx_B dQ^2 d|t| d\phi d\phi_S} = \frac{\alpha^3 x_B}{16\pi^2 Q^4 \sqrt{1+\epsilon^2}} |\mathcal{T}|^2$$

$$|\mathcal{T}|^2 = |\mathcal{T}_{BH} + \mathcal{T}_{DVCS}|^2 = |\mathcal{T}_{BH}|^2 + |\mathcal{T}_{DVCS}|^2 + \mathcal{I}$$

$\mathcal{I}$  and  $\mathcal{T}$  are functions of CFF

# DVCS at Multiple Beam Energies

## Deeply Virtual Compton Scattering $ep \rightarrow ep\gamma$

$$\frac{d^5\sigma}{dx_B dQ^2 d|t| d\phi d\phi_S} = \frac{\alpha^3 x_B}{16\pi^2 Q^4 \sqrt{1+\epsilon^2}} |\mathcal{T}|^2 \quad |\mathcal{T}|^2 = |\mathcal{T}_{\text{BH}} + \mathcal{T}_{\text{DVCS}}|^2 = |\mathcal{T}_{\text{BH}}|^2 + |\mathcal{T}_{\text{DVCS}}|^2 + \mathcal{I}$$




Different beam energies give different  $y$  at same  $(x_B, Q^2, t)$  point

$\mathcal{I} \sim 1/y^3$  Separate DVCS and interface term!

$$|\mathcal{T}^{\text{DVCS}}|^2 \sim 1/y^2$$

### Previous Hall A Result

Georges, F., Rashad, M. N. H., Stefanko, A., Dlamini, M., Karki, B., Ali, S. F., ... & Urciuoli, G. M. (2022). Deeply virtual compton scattering cross section at high Bjorken  $x_B$ . *Physical review letters*, 128(25), 252002.



# DVCS at Multiple Beam Energies


## Deeply Virtual Compton Scattering $ep \rightarrow ep\gamma$

Different beam energies give different  $y$  at same  $(x_B, Q^2, t)$  point


$$\mathcal{F} \sim 1/y^3$$

Separate DVCS and interface term!


$$|\mathcal{F}^{\text{DVCS}}|^2 \sim 1/y^2$$



Christiaens, G., et al. "First CLAS12 measurement of deeply virtual Compton scattering beam-spin asymmetries in the extended valence region." *Physical Review Letters* 130.21 (2023): 211902.



Sangbaek Lee, et al, Submitted to collaboration review




Joshua Artem Tan, et al, Submitted to collaboration review


# Run Group K

## LH2 Target

RGK Fall2018




RGK Spring2024



Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80

# Analysis Method









# Fiducial Cut

## Electron

### DC, R1

*edge* > 3 cm









# Fiducial Cut

## Electron

### ECAL, deposited energy


$v, w > 13.5$  cm



# Fiducial Cut






## Electron ECAL, EC-in as Example

*remove bad strips*



# Fiducial Cut


**Proton  
CVT**  
*edge* > 0 cm



# Fiducial Cut

## Proton CVT

*remove support bands*




# Fiducial Cut

## Photon


### FT

*remove bad circles*

photon FTCal (no fiducial cuts)




photon FTCal




# PID Cut

## Electron Triangle Cut




# PID Cut

## Electron Sampling Fraction Cut







# Energy Loss Correction

Proton  
FD








$$\Delta p = A(\theta) + B(\theta)/p + C(\theta)/p^2$$




# Energy Loss Correction

## Proton CD



# Energy Loss Correction

## Proton After Correction











# Exclusivity Cut

## Exclusivity Cut

### (FD,FD) as an Example

Variable	Definition
$MM_{ep}^2$	Missing mass squared of the $ep$ system
$MM_{ep\gamma}^2$	Missing mass squared of the $ep\gamma$ system
$MM_{e\gamma}^2$	Missing mass squared of the $e\gamma$ system
$E_{\text{miss}}$	Missing energy of the $ep\gamma$ system
$P_T^{\text{miss}}$	Missing transverse momentum of the $ep\gamma$ system
$\theta_{\gamma\gamma'}$	Angle between the measured photon and the photon reconstructed from the $ep$ kinematics
$\theta_{e\gamma}$	Opening angle between the scattered electron and photon
$\Delta E$	Energy imbalance of the $ep\gamma$ system
$\Delta\phi$	Coplanarity angle between the photon plane and the recoil-proton plane



# Exclusivity Cut

## First Selection Cut (FD,FD) as an Example

$$Q^2 > 1 \text{ GeV}^2$$

$$-t < 1 \text{ GeV}^2$$

$$W > 2 \text{ GeV}$$









$$|MM^2(ep)| < 1.5 \text{ GeV}^2$$

$$E_{miss} < 1 \text{ GeV}$$

$$P_{miss}^T < 0.25 \text{ GeV}$$

$$\theta(e, \gamma) > 5^\circ$$









$$\theta(\gamma, \gamma') < 3^\circ$$



# Exclusivity Cut

## Exclusivity Cut 1


$MM^2(ep)$  cut  
 $MM^2(ep\gamma)$  cut



# Exclusivity Cut

## Exclusivity Cut 2

$MM^2(e\gamma)$  cut  
 $E_{miss}$  cut











# Exclusivity Cut

## Exclusivity Cut 3

$P_{miss}^T$  cut










$\theta(\gamma, \gamma')$  cut

$\Delta\phi$  cut



# Exclusivity Cut


## After Exclusivity Cut



# DV $\pi^0$ P Background Exclusivity Cut


## Exclusivity Cut

Variable	Definition
$M_{\pi^0}$	Invariant mass of the two-photon system
$MM_{ep\pi^0}^2$	Missing mass squared of the $ep\pi^0$ system
$MM_{ep}^2$	Missing mass squared of the $ep$ system
$MM_{e\pi^0}^2$	Missing mass squared of the $e\pi^0$ system
$E_{miss}^0$	Missing energy of the $ep\pi^0$ system
$P_{T,miss}^0$	Missing transverse momentum of the $ep\pi^0$ system
$\theta_{\pi^0\pi^0'}$	Angle between the $\pi^0$ reconstructed from $\gamma\gamma$ kinematics and the $\pi^0$ reconstructed from $ep$ kinematics
$\Delta\phi_{\pi^0}$	Coplanarity angle between the $\pi^0$ plane and the recoil-proton plane
$\theta_{e\gamma_1}, \theta_{e\gamma_2}$	Opening angles between the scattered electron and each decay photon
$\theta_{\gamma_1\gamma_2}$	Opening angle between the two decay photons




# DV $\pi^0$ P Background Exclusivity Cut

## After Exclusivity Cut



# Data Binning

Boundary from the minimum electron energy cut









Boundary from the minimum W cut

Boundary from the minimum electron angle cut

# DV $\pi^0$ P Background Subtraction

$$B_{\pi^0} = \frac{N_{DVCS,rec}^{DV\pi^0 P,sim}(x_B, Q^2, t, \phi)}{N_{DV\pi^0 P,rec}^{DV\pi^0 P,sim}(x_B, Q^2, t, \phi)} \cdot \frac{N_{exp}^{DV\pi^0 P,rec}(x_B, Q^2, t, \phi)}{N_{exp}^{DVCS,rec}(x_B, Q^2, t, \phi)}$$





# BSA Extraction

## RGK Fall18 Pass2 7.5GeV BSA

$$A_{LU,DVCS}^{\text{corr}} = \frac{A_{LU,DVCS} - A_{\pi^0} \cdot A_{LU,DV\pi^0P}}{1 - A_{\pi^0}}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80



# RGK Fall18 Pass2 7.5GeV Integrated Charge

## Integrated Charge


Trigger Config:

- 1) electron in ECAL  
or electron in FT & 1 forward hadron
- 2) electron in ECAL, pre-scale by **2**  
or electron in FT & 1 forward hadron

Thus, the effective charge is:


$$Q = Q_{v2} + Q_{v4} \cdot \frac{1}{2} = 9.14996 \text{ mC.}$$

Not 13.69730 mC !









# Acceptance Correction

$$A_{acc} = \frac{N_{rec}(x_B, Q^2, t, \phi)}{N_{gen}(x_B, Q^2, t, \phi)}$$









# Background Merging Efficiency Correction

$$A_{eff} = 1 + \frac{I}{45nA} \left( -1 + \frac{N_{bkg}(x_B, Q^2, t, \phi)}{N_{no\ bkg}(x_B, Q^2, t, \phi)} \right)$$



# Radiation Correction

$$A_{rad} = \frac{N_{rad}(x_B, Q^2, t, \phi)}{N_{no\ rad}(x_B, Q^2, t, \phi)}$$




# Bin Centering Correction

$$A_{bcc} = \frac{d\sigma}{\langle d\sigma \rangle}$$

Evenly distributed grid in  
 $x_B, Q^2, t, \phi$  4D space

Size :  $n \times n \times n \times n$

Final Choice:  $n = 20$



# Lepton Propagator Cut

$$d\sigma_{\text{BH}^2} = \Gamma \frac{1}{x_B^2 y^2 (1 + \epsilon^2)^2 t \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)} (c_0^{\text{BH}} + c_1^{\text{BH}} \cos(\phi) + c_2^{\text{BH}} \cos(2\phi))$$

$$d\sigma_{\text{DVCS}^2} = \Gamma \frac{1}{y^2 Q^2} (c_0^{\text{DVCS}} + c_1^{\text{DVCS}} \cos(\phi) + c_2^{\text{DVCS}} \cos(2\phi))$$

$$d\sigma_{\mathcal{I}} = \Gamma \frac{1}{x_B y^3 t \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)} (c_0^{\mathcal{I}} + c_1^{\mathcal{I}} \cos(\phi) + c_2^{\mathcal{I}} \cos(2\phi) + c_3^{\mathcal{I}} \cos(3\phi))$$

$$P_1 = - \frac{1}{y(1 + \epsilon^2)} (J + 2K \cos \phi_{\text{BMK}}) \quad \text{Can reach 0 at collinear limit} \rightarrow \text{Diverge}$$

dvcsген require  $|P_1| > P_1^c$       default  $P_1^c = 0.005$       set another  $P_1^c = 0.00005$


When  $P_1$  is far away from 0,  $d\sigma(|P_1| > 0.005)/d\sigma(|P_1| > 0.00005) = 1$

When  $P_1 \sim 0$ ,  $d\sigma(|P_1| > 0.005)/d\sigma(|P_1| > 0.00005) < 1$

# Lepton Propagator Cut

Thus, define  $C_{P_1} = d\sigma(|P_1| > 0.005)/d\sigma(|P_1| > 0.00005)$

When  $C_{P_1} < \text{Certain Value}$ , we reject that data point.





# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Unpolarized Cross Section

$$d\sigma = \frac{N}{\mathcal{L} \Delta V} \cdot \frac{1 - B_{\pi_0}}{A_{acc} A_{eff} A_{rad}} \cdot A_{bcc}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80







# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Unpolarized Cross Section $0.4 \text{ GeV}^2 < |t| < 0.5 \text{ GeV}^2$

$$d\sigma = \frac{N}{\mathcal{L} \Delta V} \cdot \frac{1 - B_{\pi_0}}{A_{acc} A_{eff} A_{rad}} \cdot A_{bcc}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80





# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Polarized Cross Section

$$d\sigma = \frac{N}{\mathcal{L} \Delta V} \cdot \frac{1 - B_{\pi_0}}{A_{acc} A_{eff} A_{rad}} \cdot A_{bcc}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80




# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Polarized Cross Section $0.4 \text{ GeV}^2 < |t| < 0.5 \text{ GeV}^2$

$$d\sigma = \frac{N}{\mathcal{L} \Delta V} \cdot \frac{1 - B_{\pi_0}}{A_{acc} A_{eff} A_{rad}} \cdot A_{bcc}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80





# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Cross Section Difference

$$\Delta\sigma = d\sigma_{\rightarrow} - d\sigma_{\leftarrow}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80




# Cross Section Extraction

## RGK Fall18 Pass2 7.5GeV Cross Section Difference $0.4 \text{ GeV}^2 < |t| < 0.5 \text{ GeV}^2$

$$\Delta\sigma = d\sigma_{\rightarrow} - d\sigma_{\leftarrow}$$

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80



# Rosenbluth Separation

$$\frac{d^5\sigma}{dx_B dQ^2 d|t| d\phi d\phi_S} = \frac{\alpha^3 x_B}{16\pi^2 Q^4 \sqrt{1+\epsilon^2}} |\mathcal{T}|^2 \quad |\mathcal{T}|^2 = |\mathcal{T}_{\text{BH}} + \mathcal{T}_{\text{DVCS}}|^2 = |\mathcal{T}_{\text{BH}}|^2 + |\mathcal{T}_{\text{DVCS}}|^2 + \mathcal{I}$$

Different beam energies give different  $y$  at same  $(x_B, Q^2, t)$  point

$\mathcal{I} \sim 1/y^3$   
 $|\mathcal{T}^{\text{DVCS}}|^2 \sim 1/y^2$

Separate DVCS and interface term!

Fitting Formular:

$$\begin{pmatrix} b_{UU} \\ b_{LU} \end{pmatrix} = \begin{pmatrix} A_0(y) & A_1(y, \phi) & A_2(y, \phi) & A_3(y, \phi) & 0 \\ 0 & 0 & 0 & 0 & y(2-y)K\Gamma_I(\phi)\sin(\phi) \end{pmatrix} \begin{pmatrix} C^{\text{DVCS}} \\ \text{Re}C^{\text{INT}} \\ \text{Re}C^{\text{INT,V}} \\ \text{Re}C^{\text{INT,A}} \\ F_\Delta \end{pmatrix}$$

$\sigma$  Degree of Freedom : 4

$A_{LU}$  Degree of Freedom : 1

Beam Energy/GeV	Accumulated Charge/mC
6.4	~90
6.5	~10
7.5	~10
8.5	~80



+  
RGA: 10.6 GeV

Too close, consider as one beam energy.

All beam energies from RGK are crucial!

# Rosenbluth Separation

## Pseudo Data Test ( ~ 20% uncertainty)




Twist-2 DVCS Amplitude separated!

Agree with true(generated) value

# Rosenbluth Separation

## Estimated Experiment Test – CFF Extraction

$\text{Im } \mathcal{H} \text{ vs } \xi \quad (Q^2 = 1.10 \text{ GeV}^2)$



Conservative Systematic Uncertainty


$$\sigma_{\text{sys}} = (\sigma_{\text{fit}, 0\mathcal{F}} - \sigma_{\text{fit}, \mathcal{F}})$$

$$\sigma = \sqrt{\sigma_{\text{fit}}^2 + \sigma_{\text{sys}}^2}$$

# Rosenbluth Separation

## Estimated Experiment Test – CFF Extraction

Re  $\mathcal{H}$  vs  $\xi$  ( $Q^2 = 1.10 \text{ GeV}^2$ )




Conservative Systematic Uncertainty

$$\sigma_{\text{sys}} = (\sigma_{\text{fit}, 0\mathcal{F}} - \sigma_{\text{fit}, \mathcal{F}})$$

$$\sigma = \sqrt{\sigma_{\text{fit}}^2 + \sigma_{\text{sys}}^2}$$

# Status & Summary



**We have successfully extracted the beam spin asymmetry (BSA) as well as the unpolarized and polarized cross sections from the RGK 7.5GeV data. The Rosenbluth separation procedure is currently being refined using simulations.**

## Next Steps:

- Evaluate full systematic uncertainties.
- Analyze the full RGK data set as soon as it becomes available.
- Perform Rosenbluth separation to disentangle the various contributions, combining results across all RGK beam energies (6.4, 6.5, 7.5, and 8.4 GeV) and RGA data sets (10.2 and 10.6 GeV) to improve coverage, statistical precision, and control of systematics of extracted physics quantities.

Building on the previous studies of the proton's gravitational structure with CLAS data, this work using CLAS12 data with **multiple beam energies** will enable high-resolution imaging to map the proton's internal pressure and force distributions.

Thank you for your listening!

**Yijie Wang, Volker Burkert, Latifa Elouadrhiri,  
Richard G. Milner, Bhawani Singh**

Run Group K



Massachusetts  
Institute of  
Technology




Mar, 2026

# Backup: Way to Understand Proton's Structure

## Generalized Parton Distribution

### 3D Imaging of proton




Lin, H. W. (2021). Nucleon tomography and generalized parton distribution at physical pion mass from lattice QCD. *Physical Review Letters*, 127(18), 182001.

### Spin & Angular Momentum







$$\frac{1}{2} = \frac{1}{2} \Delta \Sigma + L_q + J_g$$

$$J_{q/g} = \lim_{t, \xi \rightarrow 0} \frac{1}{2} \int x dx [H^{q/g}(x, \xi, t) + E^{q/g}(x, \xi, t)]$$



# Backup: Fiducial Cut

**Electron  
DC, R2**  
*edge > 3 cm*









# Backup: Fiducial Cut

## Electron

### DC, R3







*edge* > 10 cm



# Backup: Fiducial Cut







## Proton DC, R1

*edge* > 3 cm



# Fiducial Cut







**Proton  
DC, R2**  
*edge > 3 cm*



# Fiducial Cut






Proton  
DC, R3

*edge* > 5 cm



# Backup: Fiducial Cut

## Proton CVT *edge* > 5 cm









# Backup: Fiducial Cut

Photon


ECAL,  
deposited  
energy

$v, w > 13.5$  cm



# Backup: Fiducial Cut

## Photon ECAL, EC-in as an Example *remove bad strips*



# Backup: Data Skimming

## Event Selection Criteria (DVCS analysis)

- $\geq 1$  electron detected in FD
- $\geq 1$  photon detected in FT or FD
- $\geq 1$  proton detected in FD or CD

## Kinematic cuts:


- $E_e > 0.1 E_{beam}$
- $E_p > M_p + 5 \text{ MeV}$
- $P_\gamma > 0.15 E_{beam}$
- $Q^2 > 0.8 \text{ GeV}^2$
- $W > 1.8 \text{ GeV}$

## Exclusivity cuts:

- Missing energy:  $-1 < E_{miss} < 2 \text{ GeV}$
- Proton missing mass<sup>2</sup>:  $0.25 < MM_p^2 < 2.0 \text{ GeV}^2$
- System missing mass<sup>2</sup>:  $-0.1 < MM_{ep\gamma}^2 < 0.1 \text{ GeV}^2$
- Transverse missing momentum:  $P_T^{miss} < 0.75 \text{ GeV}$
- Photon-missing photon angle:  $\theta_{\gamma\gamma'} < 7.5^\circ$

# Backup: Data Kinematics Coverage Constrained by Proton

ALL proton  $p$  vs theta in  $\theta > 0.0^\circ$



## Boundary From Proton:


1. Strong boundary in FD, due to bending of low momentum protons
2. Boundary depends on  $p$  and  $\theta$

Proton momentum is equivalent to  $t$

For Example:

$$t = 0.25 \text{ GeV}^2 \Leftrightarrow p = 0.51 \text{ GeV}$$

# Backup: Data Kinematics Coverage Constrained by Proton




For Example:

$$t = 0.25 \text{ GeV}^2 \Leftrightarrow p = 0.51 \text{ GeV}$$

Read from the left plot: only  $\theta > 40^\circ$  proton is within boundary



This boundary depends on  $\phi_{BMK}$




# Backup: Data Kinematics Coverage Constrained by Proton

$t = 0.25 \text{ GeV}^2$

$t = 0.35 \text{ GeV}^2$



# Backup: RGA — DVCS @ 12GeV



Sangbaek Lee