

Investigation of the Hadronic Decay $\eta \rightarrow \pi^+ \pi^- \pi^0$ with CLAS

01.02.2017 | 7th Workshop of the APS Topical Group Hadronic Physics 2017


Institute for Nuclear Physics - Jülich Research Center

One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/c ²]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP - eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes





One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/c ²]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP - eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes





One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/c ²]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP - eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes





One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/ c^2]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP -eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes





One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/ c^2]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP - eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes




One Meson, many Opportunities

Properties of the η -meson

m_η [GeV/ c^2]	0.5478
Γ_η [keV]	(1.31 ± 0.05)
$\bar{\tau}$ [s]	$5 \cdot 10^{-19}$
J^{PC}	0^{-+}

- The η -meson is a C -, P -, G - and CP - eigenstate
- All strong and electromagnetic decays are forbidden to first order
- Access to rare decay processes



Focus of this Talk

Decay Dynamics of $\eta \rightarrow \pi^+ \pi^- \pi^0$

System	Isospin State $ I, I_z\rangle$	C-Eigenvalue	G-Eigenvalue
η	$ 0, 0\rangle$	+1	+1
$(\pi^+ \pi^- \pi^0)$	$ 0, 0\rangle$	-1	-1
$(\pi^+ \pi^- \pi^0)$	$ 1, 0\rangle$	+1	-1


- Decay $\eta \rightarrow \pi^+ \pi^- \pi^0$ is G-violating \Rightarrow Forbidden to first order
- Decay is driven by isospin breaking part of strong interaction \Rightarrow C is conserved
- Decay width: $\Gamma \propto Q^{-4}$

$$\text{with: } Q^2 = \left(\frac{m_s}{m_d}\right)^2 \times \left[1 - \left(\frac{m_u}{m_d}\right)^2\right]^{-1}$$


\Rightarrow **Determine decay width $\Gamma \Rightarrow$ Access to quark mass ratio**



- Measure $\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)$, e.g. via $\frac{\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)}{\Gamma(\eta \rightarrow \gamma \gamma)}$
- Dalitz Plot Analysis**



Dalitz Plot Analysis of $\eta \rightarrow \pi^+ \pi^- \pi^0$



(a) KLOE coll., *JHEP*, 05, (2008)

Dimensionless Dalitz Plot Variables:



$$X = \sqrt{3} \frac{T_{\pi^+} - T_{\pi^-}}{T_{\pi^+} + T_{\pi^-} + T_{\pi^0}}$$

$$Y = 3 \frac{T_{\pi^0}}{T_{\pi^+} + T_{\pi^-} + T_{\pi^0}} - 1$$

- Describe three body decay by two variables (here: X and Y)
- Complete information about decay dynamics
- Parameterise decay width Γ :

$$\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3 + gX^2Y + \dots)$$
- $c \neq 0$ and $e \neq 0$:
 - i) Imply C-violation
 - ii) Cause asymmetries within the Dalitz Plot
- Compare Dalitz Plot parameters a,b,d,f from experiment and theory

Recent Measurements I





Most recent result from the KLOE-Collaboration:^(f)

- η -Mesons produced via: $e^+e^- \rightarrow \Phi \rightarrow \eta\gamma$
- $\approx 4.7 \cdot 10^6$ $\eta \rightarrow \pi^+\pi^-\pi^0$ events in the final data sample
- Fit function: $\text{Norm} \times (1 + aY + bY + cX + dX^2 + eXY + fY^3)$
- Determined asymmetries of the Dalitz Plot \Rightarrow Consistent with zero
 \Rightarrow No C-violation

(f) KLOE coll., *JHEP*, 019, (2016)

Recent Measurements II



Result from the WASA-at-COSY Collaboration:^(d)

- η -Mesons produced via: $pd \rightarrow {}^3\text{He}\eta$
- $\approx 120 \text{ k } \eta \rightarrow \pi^+\pi^-\pi^0$ events in the final data sample
- Translate each pair (X,Y) into a global bin $i(X, Y)$
→ Obtain one dimensional Dalitz Plot
- Fit function: $\text{Norm} \times (1 + aY + bY + cX + dX^2 + eXY + fY^3)$

^(d) WASA-at-COSY coll., *Phys. Rev.*, C90(045207), 2014

Recent Measurements and Theoretical Predictions

Parameter:		- a	b	d	f
Exp.	KLOE (08) ^(a)	1.090(5)($^{+8}_{-19}$)	0.124(6)(10)	0.057(6)($^{+7}_{-16}$)	0.14(1)(2)
	WASA ^(d)	1.144(18)	0.219(19)(47)	0.086(18)(15)	0.115(37)
	KLOE (16) ^(f)	1.104(3)(2)	0.142(3)($^5_{-4}$)	0.073(3)($^{+4}_{-3}$)	0.154(6)($^{+4}_{-5}$)
Theor.	ChPT (NNLO) ^(b)	1.271(75)	0.394(102)	0.055(57)	0.025(160)
	NREFT ^(c)	1.213(14)	0.308(23)	0.050(3)	0.083(19)
	PWA ^(e)	1.116(32)	0.188(12)	0.063(4)	0.091(3)

(a) KLOE coll., *JHEP*, 05, (2008)

(b) J. Bijnens and K. Ghorbani., *JHEP*, 11, (2007)



(c) S- P. Schneider et al., *JHEP*, 028, (2011)

(d) WASA-at-COSY coll., *Phys. Rev.*, C90(045207), 2014

(e) Peng Guo et al., *Phys. Rev.*, D92(05016), (2015)

(f) KLOE coll., *JHEP*, 019, (2016)


- WASA-at-COSY results used for Partial Wave Analysis (PWA) from JPAC* group
 \Rightarrow Direct calculation of: $Q = 21.4 \pm 0.4_{\text{stat}}$ ^(e)
- Dalitz Plot Analysis and determination of Q for $\gamma p \rightarrow p\eta[\eta \rightarrow \pi^+\pi^-\pi^0]$ with CLAS



* See plenary talk "Hadron Spectroscopy at JPAC" by Alessandro Pilloni at 4.40 p.m.

Small Outlook: Dalitz Plot Analysis for $\omega \rightarrow \pi^+ \pi^- \pi^0$ with WASA-at-COSY

- Theoretical description of this decay: VMD Model, Lagrangian Approach^(a), Dispersive Analysis^{(b),(c)}
→ **Input from experiment needed**
- Look at: $\frac{d^2\Gamma}{dZd\Phi} \propto (1 + 2\alpha Z + 2\beta Z^{3/2} \sin^3 \Phi + \mathcal{O}(Z^2) + \dots)$
- Analysis ongoing^{(d),(e)} for the reaction:
 $pd \rightarrow {}^3\text{He} \omega [\omega \rightarrow \pi^+ \pi^- \pi^0]$ and $pp \rightarrow pp \omega [\omega \rightarrow \pi^+ \pi^- \pi^0]$



(a) S. Leupold et al., Eur. Phys. J. A **39**, 205-212, (2009)

(b) N. Nieceknig et al., Eur. Phys. J. C **39**, 2014, (2012)

(c) I.V.Danilkin et al., Phys. Rev. **D91**, 094029, (2015)


(d) Ph.D.-Project of Lena Heijkenskjöld

(e) Ph.D.-Project of Siddhesh Sawant

Small Outlook: Dalitz Plot Analysis for $\eta' \rightarrow \pi^+ \pi^- \eta$ with CLAS


- The decay $\eta' \rightarrow \pi^+ \pi^- \eta$ allows to probe the low energy regime of QCD
 \Rightarrow Test ChPT
- Compare theory and experiment by checking decay kinematics
 \Rightarrow Dalitz Plot
- Analysis* performed using the CLAS g12 data set
- ≈ 87 k events reconstructed

* PhD-Project of S. Ghosh




CEBAF Large Acceptance Spectrometer - CLAS

CEBAF
Large
Acceptance
Spectrometer





The CLAS g12 $\gamma p \rightarrow pX$ Data Set



- Photon beam: $E_{\gamma, \text{beam}} \in [1.1 \text{ GeV}, 5.45 \text{ GeV}]$
- (Main) Contributions from:
 - Direct pion production (e.g. $\gamma p \rightarrow p\pi^+\pi^-\pi^0$)
 - π^0 , η , ω and ρ decays


Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$ Events



Decay Specific Analysis Steps


- i) Kinematic fit with reaction hypothesis: $\gamma p \rightarrow p \pi^+ \pi^- (\pi^0)$
- ii) Kinematic Limit: $M(\pi^+, \pi^-) \leq m_\eta + 3\sigma_{\eta, res} - m_{\pi^0}$
 ≈ 700 k $\eta \rightarrow \pi^+ \pi^- \pi^0$ events reconstructed so far

Towards the Dalitz Plot



- i) Look at $M_x(p)$ -spectrum as a function of the global bin $i(X, Y)$
- ii) Subtract non-resonant background
- iii) Correct for contributions from $\eta \rightarrow \pi^+ \pi^- \gamma$

Current Status



- **Left:** Dalitz Plot after background subtraction and correction for $\eta \rightarrow \pi^+ \pi^- \gamma$ events (see previous slide)
- **Right:** One Dimensional Dalitz Plot after applying efficiency correction
- Fit function: $\text{Norm} \times (1 + aY + bY^2 + dX^2 + fY^3)$
- Systematic checks and validation of current results ongoing

Summary and Outlook

1. Features of the hadronic decay: $\eta \rightarrow \pi^+ \pi^- \pi^0$
 - C-conserving
 - Isospin-violating
 - Decay amplitude driven by quark mass ratio
2. Determine decay amplitude via Dalitz Plot Analysis
 - Kinematics of a three body decay described by two variables
 - Dalitz Plot Parameters: a,b,c,d,e and f
 - Asymmetries in Dalitz Plot \leftrightarrow c and e are non-zero \leftrightarrow C-Violation
3. Several experimental and theoretical efforts to determine Dalitz Plot
 - $c = e = 0$ confirmed by experiments
 - Latest measurements done by WASA-at-COSY and KLOE:
 - i) KLOE confirmed Dalitz Plot asymmetries to be consistent with zero
 - ii) WASA-at-COSY results used in PWA (JPAC-group) to determine the quark mass ratio
 - JPAC: Partial Wave Analysis as a tool to fit experimental Dalitz Plot distributions
4. Dalitz Plot Analysis of $\eta \rightarrow \pi^+ \pi^- \pi^0$ with CLAS
 - ≈ 700 k events reconstructed so far
 - Final (one dimensional) Dalitz Plot distribution
 - To do:
 - i) Systematics checks (ongoing)
 - ii) Determination of quark mass ratio


Contents

- ▶ (2) One Meson, many Opportunities
- ▶ (3) Decay Dynamics of $\eta \rightarrow \pi^+ \pi^- \pi^0$
- ▶ (4) Dalitz Plot Analysis of $\eta \rightarrow \pi^+ \pi^- \pi^0$
- ▶ (5) Recent Measurements I
- ▶ (5) Recent Measurements II
- ▶ (7) Recent Measurements and Theoretical Predictions
- ▶ (8) Small Outlook: Dalitz Plot Analysis for $\omega \rightarrow \pi^+ \pi^- \pi^0$ with Wasa-at-COSY
- ▶ (9) Small Outlook: Dalitz Plot Analysis for $\eta' \rightarrow \pi^+ \pi^- \eta$ with CLAS
- ▶ (10) CEBAF Large Acceptance Spectrometer - CLAS
- ▶ (11) The CLAS g12 $\gamma p \rightarrow pX$ Data Set
- ▶ (12) Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$ Events
- ▶ (13) Towards the Dalitz Plot
- ▶ (14) Current Status
- ▶ (15) Summary and Outlook

Stuff that is not shown

- ▶ (18) Three Body Decays and the Dalitz Plot
- ▶ (19) Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$
- ▶ (21) The Dalitz Plot in one Dimension
- ▶ (22) Features of the 1D Dalitz Plot

Backup: Three Body Decays and the Dalitz Plot



Kinematic Constraints	NDF
3 Lorentz-Vectors	12
Momentum Conservation	-3
Energy Conservation	-1
3 Masses	-3
3 Euler Angles	-3
Total*	2



* Valid for all particles being scalars

Dalitz Plot

- Two* variables sufficient to describe three body decay
- Display decay kinematics
- Show possible resonances

Backup: Reconstruction of $\eta \rightarrow \pi^+\pi^-\pi^0$



I) The Kinematic Fit



- $\sim 2/3$ of the available CLAS g12 data set analysed so far
- Use least squares kinematic fit
 - Reaction hypothesis: $\gamma p \rightarrow p\pi^+\pi^-(\pi^0)$
 - Reject events with a probability $< 10\%$

Backup: Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$



I) The Kinematic Fit



- $\sim 2/3$ of the available CLAS g12 data set analysed so far
- Use least squares kinematic fit
 - Reaction hypothesis: $\gamma p \rightarrow p \pi^+ \pi^- (\pi^0)$
 - Reject events with a probability $< 10\%$

Backup: Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$



II) Reject Contributions from $\omega \rightarrow \pi^+ \pi^- \pi^0$



- $\sim 2/3$ of the available CLAS g12 data set analysed so far
- Use least squares kinematic fit
- Use kinematic limit: $M(\pi^+, \pi^-) \leq m_\eta + 3\sigma_{\eta,rec} - m_{\pi^0}$



Backup: Reconstruction of $\eta \rightarrow \pi^+ \pi^- \pi^0$

II) Reject Contributions from $\omega \rightarrow \pi^+ \pi^- \pi^0$



- $\sim 2/3$ of the available CLAS g12 data set analysed so far
- Use least squares kinematic fit
- Use kinematic limit: $M(\pi^+, \pi^-) \leq m_\eta + 3\sigma_{\eta, rec} - m_{\pi^0}$

Backup: The Dalitz Plot in one Dimension





Follow WASA-at-COSY analysis^(a):

- Divide Dalitz Plot in $N \times N$ bins
- Translate each pair (X, Y) into a global bin number $i \in [0, N^2 - 1]$ (e.g. $X = Y = 0 \equiv i = 60$)
- Take kinematic boundaries (see black line in left plot) into account



(a) WASA-at-COSY coll., *Phys. Rev.*, C90(045207), 2014

Backup: Features of the 1D Dalitz Plot





- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot





- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot





- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot





- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot





- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot



- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$

Backup: Features of the 1D Dalitz Plot



- Look at 1D Dalitz Plot with WASA-at-COSY values for a,b,d and f
- Center of Dalitz Plot is at $i(0, 0) = 60$
- $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + cX + dX^2 + eXY + fY^3)$