

High-precision determination of the pion–nucleon σ -term from Roy–Steiner equations

J. Ruiz de Elvira

Institute for Theoretical Physics, University of Bern

In collaboration with:

M. Hoferichter, B. Kubis, U.-G. Meißner.

PRL 115 (2015) 092301, PRL 115 (2015) 192301 , Phys.Rept. 625 (2016), Phys.Lett. B760 (2016) 74-78,
Eur. Phys. J. A52 (2016), 331, arXiv:1610.08978

The proton mass Workshop, Trento, April 6th, 2017

u^b

b
UNIVERSITÄT
BERN

AEC
ALBERT EINSTEIN CENTER
FOR FUNDAMENTAL PHYSICS



Decomposition of the nucleon mass

$$m_N = \langle N | \theta_\mu^\mu | N \rangle = \langle N | \underbrace{\frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G_a^{\mu\nu}}_{\text{Trace anomaly}} + \underbrace{m_u \bar{u}u + m_d \bar{d}d + m_s \bar{s}s + \dots}_{\text{Higgs}} | N \rangle$$

- Mass largely generated by gluon field energy via the trace anomaly of the QCD energy-momentum tensor

- $\langle N | m_u \bar{u}u + m_d \bar{d}d | N \rangle \doteq \sigma_{\pi N}$

→ measures light-quark contribution to the nucleon mass, how large precisely?

- $\langle N | m_s \bar{s}s | N \rangle \doteq \sigma_s$

→ strangeness content of the nucleon

The pion-nucleon σ -term

Scalar form factor of the nucleon:

$$\sigma(t) = \langle N(p') | \hat{m} (\bar{u}u + \bar{d}d) | N(p) \rangle \quad t = (p' - p)^2 \quad \sigma_{\pi N} = \sigma(0)$$

- $\sigma_{\pi N}$ measures the **light-quark contribution** to the nucleon mass
- Unfortunately, no direct experimental access to it
- Only very recent precise lattice results
- Linked to πN via the **Cheng-Dashen theorem**

[Cheng, Dashen 1971]


$$F_\pi^2 \bar{D}^+(\nu = 0, t = 2M_\pi^2) = \underbrace{\sigma(2M_\pi^2)}_{\sigma_{\pi N} + \Delta_\sigma} + \Delta_R$$
$$\underbrace{F_\pi^2(d_{00}^+ + 2M_\pi^2 d_{01}^+)}_{F_D} + \Delta_D$$

$$|\Delta_R| \lesssim 2 \text{ MeV}$$

[Bernard, Kaiser, Meißner 1996]

$$\Delta_D - \Delta_\sigma = (-1.8 \pm 0.2) \text{ MeV}$$

[Hoferichter et al. 2012]



Phenomenological status


- **Karlsruhe/Helsinki** partial-wave analysis KH80 [Höhler et al. 1980s]
 - ↪ comprehensive analyticity constraints, old data
- Formalism for the extraction of $\sigma_{\pi N}$ via the **Cheng–Dashen low-energy theorem**
 - ↪ “canonical value” $\sigma_{\pi N} \sim 45$ MeV, based on KH80 input [Gasser, Leutwyler, Locher, Sainio 1988,1991]
- **GWU/SAID** partial-wave analysis [Pavan, Strakovsky, Workman, Arndt 2002]
 - ↪ much larger value $\sigma_{\pi N} = (64 \pm 8)$ MeV
- More recently: ChPT in different regularizations (w/ and w/o Δ) [Alarcón et al. 2012]
 - ↪ fit to PWAs, $\sigma_{\pi N} = 59 \pm 7$ MeV

Phenomenological status


- **Karlsruhe/Helsinki** partial-wave analysis KH80 [Höhler et al. 1980s]
 - ↪ comprehensive analyticity constraints, old data
- Formalism for the extraction of $\sigma_{\pi N}$ via the **Cheng–Dashen low-energy theorem**
 - ↪ “canonical value” $\sigma_{\pi N} \sim 45$ MeV, based on KH80 input [Gasser, Leutwyler, Locher, Sainio 1988,1991]
- **GWU/SAID** partial-wave analysis [Pavan, Strakovsky, Workman, Arndt 2002]
 - ↪ much larger value $\sigma_{\pi N} = (64 \pm 8)$ MeV
- More recently: ChPT in different regularizations (w/ and w/o Δ) [Alarcón et al. 2012]
 - ↪ fit to PWAs, $\sigma_{\pi N} = 59 \pm 7$ MeV
- This talk: two new sources of information on low-energy πN scattering
 - Precision extraction of πN **scattering lengths** from **hadronic atoms** [Baru et al. 2011]
 - **Roy-equation** constraints: analyticity, unitarity, crossing symmetry

Hadronic atoms: constraints for πN

- $\pi H/\pi D$: bound state of π^- and p/d spectrum sensitive to threshold πN amplitude
- Combined analysis of πH and πD :
 - $a_0^+ \equiv a^+ = (7.5 \pm 3.1) \cdot 10^{-3} M_\pi^{-1}$
 - $a_0^- \equiv a^- = (86.0 \pm 0.9) \cdot 10^{-3} M_\pi^{-1}$
 - ↪ Large a^+ suggests a large $\sigma_{\pi N}$,
- But: a^+ very sensitive to isospin breaking, PWA based on $\pi^\pm p$ channels
 - ↪ use instead
- Isospin breaking in $\sigma_{\pi N}$ could be important
- We revisit the Cheng-Dashen low-energy theorem



[Baru et al. 2011]



$$\bar{a}^+ = a^+ + \frac{1}{1+M_\pi/m_p} \left\{ \frac{M_\pi^2 - m_0^2}{\pi F_\pi^2} c_1 - 2\alpha f_1 \right\}$$

Motivation: Why Roy-Steiner equations?

Roy(-Steiner) eqs. = Partial-Wave (Hyperbolic) Dispersion Relations coupled by unitarity and crossing symmetry

- **Respect all symmetries:** analyticity, unitarity, crossing
- **Model independent** \Rightarrow the actual parametrization of the data is irrelevant once it is used in the integral.
- Framework allows for **systematic improvements** (subtractions, higher partial waves, ...)
- **PW(H)DRs** help to study processes with **high precision**:
 - $\pi\pi$ -scattering: [Ananthanarayan et al. (2001), García-Martín et al. (2011)]
 - πK -scattering: [Büttiker et al. (2004)]
 - $\gamma\gamma \rightarrow \pi\pi$ scattering: [Hoferichter et al. (2011)]

Warm up: Roy-equations for $\pi\pi$

- $\pi\pi \rightarrow \pi\pi \Rightarrow$ fully crossing symmetric in Mandelstam variables s , t , and $u = 4M_\pi - s - t$
- Start from twice-subtracted fixed-t DRs of the generic form

$$T^I(s, t) = c(t) + \frac{1}{\pi} \int_{4m_\pi^2}^{\infty} \frac{ds'}{s'^2} \left[\frac{s^2}{(s' - s)} - \frac{u^2}{(s' - u)} \right] \text{Im } T^I(s', t)$$

- Subtraction functions $c(t)$ are determined via crossing symmetry

↪ functions of the $l=0,2$ scattering lengths: a_0^0 and a_0^2

- PW-expansion of these DRs yields the **Roy-equations**

[Roy (1971)]

$$t_J^l(s) = ST_J^l(s) + \sum_{J'=0}^{\infty} (2J' + 1) \sum_{l'=0,1,2} \int_{4m_\pi^2}^{\infty} ds' K_{JJ'}^{ll'}(s', s) \text{Im } t_{J'}^{l'}(s')$$

- $K_{JJ'}^{ll'}(s', s) \equiv$ kernels \Rightarrow analytically known

Roy–Steiner equations for πN : differences to $\pi\pi$ Roy equations

Key differences compared to $\pi\pi$ Roy equations


- **Crossing:** coupling between $\pi N \rightarrow \pi N$ (s-channel) and $\pi\pi \rightarrow \bar{N}N$ (t -channel)

⇒ need a different kind of dispersion relations

[Hite, Steiner 1973, Büttiker et al. 2004]

- **Unitarity** in t-channel, e.g. in single-channel approximation

$$\text{Im}f_{\pm}^J(t) = \sigma_t^\pi f_{\pm}^J(t) t_J^I(t)^*$$



⇒ **Watson's theorem:** phase of $f_{\pm}^J(t)$ equals δ_{IJ}

[Watson 1954]

↪ solution in terms of Omnès function

[Muskhelishvili 1953, Omnès 1958]

- **Large pseudo-physical region** in t -channel

↪ $\bar{K}K$ intermediate states for s-wave in the region of the $f_0(980)$

Limited range of validity

$$\sqrt{s} \leq \sqrt{s_m} = 1.38 \text{ GeV}$$

$$\sqrt{t} \leq \sqrt{t_m} = 2.00 \text{ GeV}$$

Input/Constraints


- S- and P-waves **above** matching point
 $s > s_m$ ($t > t_m$)
- Inelasticities
- Higher waves (D-, F-, \dots)
- Scattering lengths from hadronic atoms

[Baru et al. 2011]


Output

- S- and P-wave **phase-shifts** at low energies
 $s < s_m$ ($t < t_m$)
- Subthreshold parameters
 - ▷ Pion-nucleon σ -term
 - ▷ Nucleon **form factor** spectral functions
 - ▷ ChPT **LECs**

Uncertainties: s-channel pw



Uncertainties: Imaginary part t-channel pw



Results for the sigma-term

$$\sigma_{\pi N} = F_\pi^2 \left(d_{00}^+ + 2M_\pi^2 d_{01}^+ \right) + \Delta_D - \Delta_\sigma - \Delta_R$$

- subthreshold parameters output of the Roy–Steiner equations

$$d_{00}^+ = -1.36(3) M_\pi^{-1} \quad [\text{KH}: -1.46(10) M_\pi^{-1}],$$

$$d_{01}^+ = 1.16(2) M_\pi^{-3} \quad [\text{KH}: 1.14(2) M_\pi^{-3}]$$

- $|\Delta_R| \lesssim 2 \text{ MeV}$

[Bernard, Kaiser, Meißner 1996]

- $\Delta_D - \Delta_\sigma = -(1.8 \pm 0.2) \text{ MeV}$

[Hoferichter et al. 2012]

- Isospin breaking in the CD theorem shifts $\sigma_{\pi N}$ by $+3.0 \text{ MeV}$

- Final results: $\sigma_{\pi N} = (59.1 \pm 1.9_{\text{RS}} \pm 3.0_{\text{LET}}) \text{ MeV} = (59.1 \pm 3.5) \text{ MeV}$

[MH, JRE, Kubis, Meißner]

- $\sigma_{\pi N}$ depends linearly on the scattering lengths:

$$\sigma_{\pi N} = 59.1 + \sum_{l_s} c_{l_s} \Delta a_{0+}^{l_s}$$

$\sigma_{\pi N}$: comparison with KH80 and SAID

Comparison with KH80

- RS eqs. with KH80 input $\hookrightarrow \sigma_{\pi N} = 46$ MeV

\hookrightarrow to be compared with $\sigma_{\pi N} = 45$ MeV

Gasser, Leutwyler, Sainio 1988, Gasser, Leutwyler, Sainio 1991

\hookrightarrow KH80 is internally **consistent** but at odd with the modern **SL** determinations

How are d_{00}^+ and d_{01}^+ extracted in KH80 and SAID?

- Standard approach:

Gasser, Leutwyler, Locher, Sainio 1988

replace d_{00}^+ and d_{01}^+ in favor of threshold parameters: a_{0+}^+ and a_{1+}^+


\hookrightarrow corrections from PWA via DRs (D^+ and E^+)

	Born	a_{0+}^+	a_{1+}^+	D^+	E^+	$\Sigma_d = F_\pi^2 (d_{00}^+ + 2M_\pi^2 d_{01}^+)$
KH80	-133	-7	+352	-91	-72	50
SAID	-127	0	+351	-88	-69	67
diff	+6	7	-1	+3	+3	17

- difference with KH80 due the a_{0+}^+
- large weight of a_{1+}^+ \Rightarrow **It has to be known extremely accurately!**
 \hookrightarrow the difference 132.7 (SAID)/131.2 (RS) translates in 5 MeV in the $\sigma_{\pi N}$

Comparison with lattice $\sigma_{\pi N}$ results

- Recent lattice determination of $\sigma_{\pi N}$ at (almost) the physical point
 - BMW $\sigma_{\pi N} = 38(3)(3)$ MeV [Durr et al. 2015]
 - χ QCD $\sigma_{\pi N} = 44.4(3.2)(4.5)$ MeV [Yang et al. 2015]
 - ETMC $\sigma_{\pi N} = 37.22(2.57)(1)$ MeV [Abdel-Rehim et al. 2015]
 - RQCD $\sigma_{\pi N} = 35(6)$ MeV [Bali et al. 2016]
- The linear dependence of $\sigma_{\pi N}$ on the scattering lengths introduces an additional constraint



- Inconsistent with the hadronic atom phenomenology

→ determine the πN scattering lengths in the lattice

Comparison with experimental cross-section data

Unravel the tension around the σ -term comparing with the experimental πN data base

- Generate RS differential cross sections

- ▷ RS S and P waves

- ▷ higher partial waves from SAID and KH80

[Workman et al. 2006,2012, Höhler et al. 1980s]

- ▷ EM interactions implemented using Tromborg procedure

[Tromborg et al. 1977]

- Uncertainties from statistical effects, SL, input variation

- ▷ below $T_\pi = 50$ MeV uncertainties dominated by scattering length errors

- ↪ disentangle RS SL solutions by looking at the data base


- Define:

$$\chi_{a'_{0+}}^2 = \sum_{i,j} \frac{(\mathcal{O}_{i,j}^{\text{exp}} - \mathcal{O}_{i,j}^{\text{RS}}(a'_{0+}))^2}{\Delta \mathcal{O}_{i,j}^{\text{exp}}}$$

- Discrepancy concentrated in the $\pi^+ p \rightarrow \pi^+ p$ channel

	RS	KH80
$a'_{0+}^{1/2} [10^{-3} M_\pi^{-1}]$	169.8 ± 2.0	173 ± 3
$a'_{0+}^{3/2} [10^{-3} M_\pi^{-1}]$	-86.3 ± 1.8	-101 ± 4

Cross-section data: $\pi^+ p \rightarrow \pi^+ p$ channel



Comparison with experimental cross-section data

- Linearized version of RS $d\sigma/d\Omega$ around the HA scattering lengths
 - ▷ good description in the KH80 region
- minimize the χ^2 -like as a function of a_{0+}^l


channel	SL combination	result	HA SL	KH80 SL
$\pi^+ p \rightarrow \pi^+ p$	$a_{0+}^{3/2}$	-86.6	-86.3 ± 1.8	-101 ± 4
$\pi^- p \rightarrow \pi^- p$	$2a_{0+}^{1/2} - a_{0+}^{3/2}$	256.2	253.3 ± 4.4	245 ± 7

- Current challenges
 - ▷ CXS channel
 - ↪ inconsistencies in the data base
 - ▷ GWU/SAID data base normalization constants
 - ↪ D'Agostini bias ⇒ unbiased low energy fit



In progress

Nucleon form factor spectral functions

- $\pi\pi \rightarrow \bar{N}N$ partial waves + F_π^V pion form factor
↪ $\pi\pi$ contribution to the **isovector spectral functions**
- consistent $\pi\pi$ phase shifts in f_\pm^\pm and F_π^V
↪ Watson theorem is satisfied
- modern pion form factor data
- **isospin breaking:** $m_p - m_n$ in pole terms, subthreshold parameters, consistent $\rho - \omega$ mixing



[BaBar 2009, KLOE 2012, BESIII 2015]



[Hoferichter, Kubis, JRE, Hammer, Meißen 2016]

$\pi\pi$ continuum and proton radius puzzle

- sum rules for the isovector radii:

$$\langle r_{E/M}^2 \rangle^v = \frac{6}{\pi} \int_0^\Lambda dt' \frac{\text{Im } G_{E/M}^v(t')}{t'^2}$$

	$\Lambda = 1 \text{ GeV}$	$\Lambda = 2m_N$
$\langle r_E^2 \rangle^v [\text{fm}^2]$	0.418(32)	0.405(36)
$\langle r_M^2 \rangle^v [\text{fm}^2]$	1.83(10)	1.81(11)

- correcting normalization by single heavier resonance: ρ' , ρ'' :

reduces the radii only to: $\Delta \langle r_E^2 \rangle^v = -(0.006 \dots 0.008) \text{ fm}^2$

$$\Delta \langle r_M^2 \rangle^v = -(0.05 \dots 0.07) \text{ fm}^2$$

- with $\langle r_E^2 \rangle^n = -0.1161(22) \text{ fm}^2$ (n scattering on heavy atoms):

↪ proton radius puzzle \iff isovector radius puzzle

$$\langle r_E^2 \rangle^v = 0.412 \text{ fm}^2 (\mu\text{H}) \quad \text{vs.} \quad \langle r_E^2 \rangle^v = 0.442 \text{ fm}^2 (\text{CODATA})$$

▷ mild preference for small proton charge radius

[Hoferichter, Kubis, JRE, Hammer, Meißner 2016]

Matching to Chiral Perturbation Theory

Matching to ChPT at the subthreshold point:

- Chiral expansion expected to work best at **subthreshold point**
 - ▷ Maximal distance from threshold **singularities**
 - ▷ πN amplitude can be expanded as **polynomial**
- Preferred choice for NN scattering due to proximity of relevant kinematic regions

Express the subthreshold parameters in terms of the LECs to $\mathcal{O}(p^4)$

$$d_{00}^+ = -\frac{2M_\pi^2(2\tilde{c}_1 - \tilde{c}_3)}{F_\pi^2} + \frac{g_a^2(3 + 8g_a^2)M_\pi^3}{64\pi F_\pi^4} + M_\pi^4 \left\{ \frac{16\bar{e}_{14}}{F_\pi^2} - \frac{2c_1 - c_3}{16\pi^2 F_\pi^4} \right\}$$

- Chiral πN amplitude to $\mathcal{O}(p^4)$ depends on **13** low-energy constants
- Roy–Steiner system contains **10 subtraction constants**
 - ▷ Calculate remaining **3** from **sum rules**
 - ▷ **Invert the system** to solve for LECs

Chiral low-energy constants

	NLO	N ² LO	N ³ LO
c_1 [GeV ⁻¹]	-0.74 ± 0.02	-1.07 ± 0.02	-1.11 ± 0.03
c_2 [GeV ⁻¹]	1.81 ± 0.03	3.20 ± 0.03	3.13 ± 0.03
c_3 [GeV ⁻¹]	-3.61 ± 0.05	-5.32 ± 0.05	-5.61 ± 0.06
c_4 [GeV ⁻¹]	2.17 ± 0.03	3.56 ± 0.03	4.26 ± 0.04
$\bar{d}_1 + \bar{d}_2$ [GeV ⁻²]	—	1.04 ± 0.06	7.42 ± 0.08
\bar{d}_3 [GeV ⁻²]	—	-0.48 ± 0.02	-10.46 ± 0.10
\bar{d}_5 [GeV ⁻²]	—	0.14 ± 0.05	0.59 ± 0.05
$\bar{d}_{14} - \bar{d}_{15}$ [GeV ⁻²]	—	-1.90 ± 0.06	-12.18 ± 0.12
\bar{e}_{14} [GeV ⁻³]	—	—	0.89 ± 0.04
\bar{e}_{15} [GeV ⁻³]	—	—	-0.97 ± 0.06
\bar{e}_{16} [GeV ⁻³]	—	—	-2.61 ± 0.03
\bar{e}_{17} [GeV ⁻³]	—	—	0.01 ± 0.06
\bar{e}_{18} [GeV ⁻³]	—	—	-4.20 ± 0.05

- Subthreshold errors tiny, chiral expansion dominates uncertainty
- \bar{d}_i at N³LO increase by an order of magnitude
 - ↪ due to terms proportional to $g_A^2(c_3 - c_4) = -16$ GeV⁻¹
 - ↪ mimic loop diagrams with Δ degrees of freedom
- What's going on with chiral convergence?
 - ↪ look at convergence of threshold parameters with LECs fixed at subthreshold point

Convergence of the chiral series

	NLO	N ² LO	N ³ LO	RS
$a_{0+}^+ [10^{-3} M_\pi^{-1}]$	-23.8	0.2	-7.9	-0.9 ± 1.4
$a_{0+}^- [10^{-3} M_\pi^{-1}]$	79.4	92.9	59.4	85.4 ± 0.9
$a_{1+}^+ [10^{-3} M_\pi^{-3}]$	102.6	121.2	131.8	131.2 ± 1.7
$a_{1+}^- [10^{-3} M_\pi^{-3}]$	-65.2	-75.3	-89.0	-80.3 ± 1.1
$a_{1-}^+ [10^{-3} M_\pi^{-3}]$	-45.0	-47.0	-72.7	-50.9 ± 1.9
$a_{1-}^- [10^{-3} M_\pi^{-3}]$	-11.2	-2.8	-22.6	-9.9 ± 1.2
$b_{0+}^+ [10^{-3} M_\pi^{-3}]$	-70.4	-23.3	-44.9	-45.0 ± 1.0
$b_{0+}^- [10^{-3} M_\pi^{-3}]$	20.6	23.3	-64.7	4.9 ± 0.8

- N³LO results bad due to large Delta loops
- matching to ChPT with the explicit Δ 's
 - improvement of the chiral convergence
- Conclusion: lessons for few-nucleon applications
 - either include the Δ to reduce the size of the loop corrections or use LECs from subthreshold kinematics
 - error estimates: consider chiral convergence of a given observable, difficult to assign a global chiral error to LECs


[Siemens, JRE, Epelbaum, Hoferichter, Krebs, Kubis, Meißner 2016]

The “ruler plot” vs. ChPT

Lattice QCD simulations can be performed at different quark/pion masses

Pion mass dependence of m_N up to NNNLO in ChPT, using

- Input from Roy–Steiner solution



thanks to A. Walker-Loud for providing the lattice data

- range of convergence of the chiral expansion is very limited
- huge cancellation amongst terms to produce a linear behavior


- Derived a closed system of Roy–Steiner equations for πN
- Numerical solution and error analysis of the full system of RS eqs.
- Precise determination of the $\sigma_{\pi N}$
 - ▷ Roy–Steiner formalism reproduces KH80 result with KH80 input
 - ▷ With modern input for scattering lengths and coupling constant $\sigma_{\pi N}$ increases
 - ▷ results from hadronic atom results compatible with low-energy πN scattering data
- t- channel → nucleon form factor spectral functions
 - sum rules for isovector radii → proton radius puzzle
- Extraction of the ChPT LECs
- Study of the chiral convergence

Thank you

Spare slides

Motivation: Why πN scattering?


- **Low energies:** test chiral dynamics in the baryon sector
⇒ low-energy theorems e.g. for the scattering lengths
- **Higher energies:** resonances, baryon spectrum
- **Input for NN scattering:** LECs c_i , πNN coupling




- **Crossed channel $\pi\pi \rightarrow \bar{N}N$:** nucleon form factors

⇒ probe the structure of the nucleon

- ▷ scalar form factors (S-wave)
- ▷ electromagnetic form factors (P-waves)
- ▷ generalized PDF (D-waves)



Roy-Steiner equations for πN : flow of information



- **Roy-equations** rigorously valid for a finite energy range

⇒ introduce a matching point s_m

- only partial waves with $J \leq J_{\max}$ are solved
- assume isospin limit

- **Input**

- High-energy region: $\text{Im}t_{IJ}(s)$ for $s \geq s_m$ and for all J
- Higher partial waves: $\text{Im}t_{IJ}(s)$ for $J > J_{\max}$ and for all s

- **Output**

- Self-consistent solution for $\delta_{IJ}(s)$ for $J \leq J_{\max}$ and $s_{\text{th}} \leq s \leq s_m$
- Constraints on subtraction constants

πN -scattering basics

$$\pi^a(q) + N(p) \rightarrow \pi^b(q') + N(p')$$

- **Isospin Structure:**

$$T^{ba} = \delta^{ba} T^+ + \epsilon^{ab} T^-$$

- **Lorentz Structure:** $I \in \{+, -\}$

$$T^I = \bar{u}(p') \left(A^I + \frac{\not{q} + \not{q}'}{2} B^I \right) u(p)$$

$$D^I = A^I + \nu B^I, \quad \nu = \frac{s-u}{4m}$$

- **Isospin basis:** $I_s \in \{1/2, 3/2\}$


$$\{T^+, T^-\} \Leftrightarrow T^{1/2}, T^{3/2}$$

- **PW projection:**

$$s\text{-channel pw: } f_{I\pm}^I$$

$$t\text{-channel pw: } f_{\pm}^J$$

Bose symmetry \Rightarrow even/odd $J \Leftrightarrow I = +/-$



πN -scattering basics: partial waves

- **s-channel** projection:

$$f_{l\pm}^I(W) = \frac{1}{16\pi W_1} \left\{ (E+m)[A_l^I(s) + (W-m)B_l^I(s)] + (E-m)[-A_{l\pm 1}^I(s) + (W+m)B_{l\pm 1}^I(s)] \right\}$$

$$X_l^I(s) = \int_{-1}^1 dz_s P_l(z_s) X^I(s, t) \Big|_{t=t(s, z_s)=-2q^2(1-z_s)} \quad \text{for } X \in \{A, B\} \text{ and } W = \sqrt{s}$$

- **McDowell symmetry:** $f_{l+}^I(W) = -f_{(l+1)-}^I(-W) \quad \forall I \geq 0$

- **t-channel** projection:

$$f_+^J(t) = -\frac{1}{4\pi} \int_0^1 dz_t P_J(z_t) \left\{ \frac{p_t^2}{(p_t q_t)^J} A^I(s, t) \Big|_{s=s(t, z_t)} - \frac{m}{(p_t q_t)^{J-1}} z_t B^I(s, t) \Big|_{s=s(t, z_t)} \right\} \quad \forall J \geq 0$$


$$f_-^J(t) = \frac{1}{4\pi} \frac{\sqrt{J(J+1)}}{2J+1} \frac{1}{(p_t q_t)^{J-1}} \int_0^1 dz_t [P_{J-1}(z_t) - P_{J+1}(z_t)] B^I(s, t) \Big|_{s=s(t, z_t)} \quad \forall J \geq 1$$

- **Bose symmetry** \Rightarrow even/odd $J \Leftrightarrow I = +/-$

πN -scattering basics: Unitarity relations


- **s-channel** unitarity relations ($I_s \in \{1/2, 3/2\}$):

$$\text{Im } f_{I\pm}^{Is}(W) = q |f_{I\pm}^{Is}(W)|^2 \theta(W - W_+) + \frac{1 - (\eta_{I\pm}^{Is}(W))^2}{4q} \theta(W - W_{\text{inel}})$$



- **t-channel** unitarity relations: 2-body intermediate states: $\pi\pi + \bar{K}K + \dots$

$$\text{Im } f_{\pm}^J(t) = \sigma_t^\pi (t_{\mathbf{J}}^J(t))^* f_{\pm}^J(t) \theta(t - t_\pi) + 2c_J \sqrt{2} k_t^{2J} \sigma_t^K (g_{\mathbf{J}}^J(t))^* h_{\pm}^J(t) \theta(t - t_K)$$



- Only linear in $f_{\pm}^J(t) \Rightarrow$ less restrictive

Roy-Steiner equations for πN : HDR's

- **Hyperbolic DRs:** $(s - a)(u - a) = b = (s' - a)(u' - a)$ with $a, b \in \mathbb{R}$

$$A^+(s, t; a) = \frac{1}{\pi} \int_{s_+}^{\infty} \textcolor{red}{ds'} \left[\frac{1}{s' - s} + \frac{1}{s' - u} - \frac{1}{s' - a} \right] \text{Im } A^+(s', t') + \frac{1}{\pi} \int_{t_\pi}^{\infty} \textcolor{blue}{dt'} \frac{\text{Im } A^+(s', t')}{t' - t}$$
$$B^+(s, t; a) = N^+(s, t) + \frac{1}{\pi} \int_{s_+}^{\infty} \textcolor{red}{ds'} \left[\frac{1}{s' - s} - \frac{1}{s' - u} \right] \text{Im } B^+(s', t') + \frac{1}{\pi} \int_{t_\pi}^{\infty} \textcolor{blue}{dt'} \frac{\nu}{\nu'} \frac{\text{Im } B^+(s', t')}{t' - t}$$
$$N^+(s, t) = g^2 \left(\frac{1}{m^2 - s} - \frac{1}{m^2 - u} \right) \quad \text{similar for } A^-, B^- \text{ and } N^- \quad [\text{Hite/Steiner (1973)}]$$

- Why **HDR?**

- Combine all physical regions \Rightarrow crucial for t-channel projection
- Evade double-spectral regions \Rightarrow the PW decompositions converge
- Range of convergence can be maximized by tuning the free hyperbola parameter a
- No kinematical cuts, manageable kernel functions

Roy-Steiner equations for πN : derivation

- Recipe to derive **Roy-Steiner** equations:
 - **Expand** imaginary parts in terms of s- and t-channel partial waves
 - **Project** onto s- and t-channel partial waves
 - **Combine** the resulting equations using s- and t-channel **PW unitarity relations**

- Similar structure to $\pi\pi$ **Roy equations**

- **Validity:** assuming Mandelstam analyticity

- s-channel \Rightarrow optimal for $a = -23.2M_\pi^2$

$$s \in [s_+ = (m + M_\pi)^2, 97.30 M_\pi^2] \quad \Leftrightarrow \quad W \in [W_+ = 1.08 \text{ GeV}, 1.38 \text{ GeV}]$$

- t-channel \Rightarrow optimal for $a = -2.71M_\pi^2$

$$t \in [t_\pi = 4M_\pi^2, 205.45 M_\pi^2] \quad \Leftrightarrow \quad \sqrt{t} \in [\sqrt{t_\pi} = 0.28 \text{ GeV}, 2.00 \text{ GeV}] .$$

Roy-Steiner equations for πN : subtractions

- **Subtractions** are necessary to ensure the convergence of DR integrals
⇒ asymptotic behavior
- Can be introduced to lessen the dependence of the low-energy solution on the high-energy behavior
- Parametrize high-energy information in (a priori unknown) subtraction constants
⇒ matching to ChPT
- Subthreshold expansion around $\nu = t = 0$

$$\begin{aligned}\bar{A}^+(\nu, t) &= \sum_{m,n=0}^{\infty} a_{mn}^+ \nu^{2m} t^n & \bar{B}^+(\nu, t) &= \sum_{m,n=0}^{\infty} b_{mn}^+ \nu^{2m+1} t^n, \\ \bar{A}^-(\nu, t) &= \sum_{m,n=0}^{\infty} a_{mn}^- \nu^{2m+1} t^n & \bar{B}^-(\nu, t) &= \sum_{m,n=0}^{\infty} b_{mn}^- \nu^{2m} t^n,\end{aligned}$$

where

$$\begin{aligned}\bar{A}^+(s, t) &= A^+(s, t) - \frac{g^2}{m} & \bar{B}^+(s, t) &= B^+(s, t) - g^2 \left[\frac{1}{m^2 - s} - \frac{1}{m^2 - u} \right], \\ \bar{A}^-(s, t) &= A^-(s, t), & \bar{B}^-(s, t) &= B^-(s, t) - g^2 \left[\frac{1}{m^2 - s} + \frac{1}{m^2 - u} \right] + \frac{g^2}{2m^2},\end{aligned}$$

RS-eqs for πN : subthreshold expansion

- Subthreshold expansion around $\nu = t = 0$

$$A^+(\nu, t) = \frac{g^2}{m} + d_{00}^+ + d_{01}^+ t + d_{10}^+ \nu^2 + \mathcal{O}(\nu^2 t, t^2)$$

$$A^-(\nu, t) = \nu a_{00}^- + a_{01}^- \nu t + a_{10}^- \nu^3 + \mathcal{O}(\nu^5, \nu t^2, \nu^3 t)$$

$$B^+(\nu, t) = g^2 \frac{4m\nu}{(m^2 - s_0)^2} + \nu b_{00}^+ + \mathcal{O}(\nu^3, \nu t) ,$$

$$B^-(\nu, t) = g^2 \left[\frac{2}{m^2 - s_0} - \frac{t}{(m^2 - s_0)^2} \right] - \frac{g^2}{2m^2} + b_{00}^- + b_{01}^- t + b_{10}^- \nu^2 + \mathcal{O}(\nu^2, \nu^2 t, t^2)$$

- pseudovector Born terms: $D^I = A^I + \nu B^I$

$$\bar{D}^+ = d_{00}^+ + d_{01}^+ t + d_{10}^+ \nu^2$$

$$d_{mn}^+ = a_{mn}^+ + b_{m-1,n}^+, \quad d_{mn}^- = a_{mn}^- + b_{mn}^- .$$

- Sum rules for subthreshold parameters:


$$d_{00}^+ = -\frac{g^2}{m} + \frac{1}{\pi} \int_{s_+}^{\infty} ds' h_0(s') [\text{Im } A^+(s', z_s')]_{(0,0)} + \frac{1}{\pi} \int_{t_\pi}^{\infty} \frac{dt'}{t'} [\text{Im } A^+(t', z_t')]_{(0,0)}$$

$$h_0(s') = \frac{2}{s' - s_0} - \frac{1}{s' - a}$$

Solving Roy-Steiner equations for πN : Recoupling schemes


- **s-channel** subproblem:

- Kernels are diagonal for $I \in \{+, -\}$, but unitarity relations are diagonal for $I_s \in \{1/2, 3/2\} \Rightarrow$ all partial-waves are interrelated
- Once the t-channel PWs are known
 \Rightarrow Structure similar to $\pi\pi$ Roy-equations



- **t-channel** subproblem:

- Only higher PWs couple to lower ones
- Only PWs with even or odd J are coupled
- No contribution from f_+^J to f_-^{J+1}
 \Rightarrow Leads to Muskhelishvili-Omnès problem



s-channel RS equations

$$\begin{aligned}
 f_{l+}^l(W) &= N_{l+}^l(W) + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{l'=0}^{\infty} \left\{ K_{ll'}^l(W, W') \operatorname{Im} f_{l'+}^{l'}(W') + K_{ll'}^l(W, -W') \operatorname{Im} f_{(l'+1)-}^{l'}(-W') \right\} \\
 &\quad + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \sum_J \left\{ G_{lJ}(W, t') \operatorname{Im} f_+^J(t') + H_{lJ}(W, t') \operatorname{Im} f_-^J(t') \right\} \\
 &= -f_{(l+1)-}^l(-W) \quad \forall l \geq 0 , \quad [\text{Hite/Steiner (1973)}]
 \end{aligned}$$

- e $K_{ll'}^l(W, W')$, $G_{lJ}(W, t')$ and $H_{lJ}(W, t')$ -Kernels: analytically known,
e.g. $K_{ll'}^l(W, W') = \frac{\delta_{ll'}}{W' - W} + \dots \quad \forall l, l' \geq 0$,
- **Validity**: assuming Mandelstam analyticity
 \Rightarrow optimal for $a = -23.2M_\pi^2$

$$s \in [s_+ = (m + M_\pi)^2, 97.30 M_\pi^2] \quad \Leftrightarrow \quad W \in [W_+ = 1.08 \text{ GeV}, 1.38 \text{ GeV}]$$

Roy-Steiner equations for πN : t-channel

t-channel RS equations

$$\begin{aligned} f_+^J(t) &= \tilde{N}_+^J(t) + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{l=0}^{\infty} \left\{ \tilde{G}_{Jl}(t, W') \operatorname{Im} f_{l+}^l(W') + \tilde{G}_{Jl}(t, -W') \operatorname{Im} f_{(l+1)-}^l(W') \right\} \\ &\quad + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \sum_{J'} \left\{ \tilde{K}_{JJ'}^1(t, t') \operatorname{Im} f_+^{J'}(t') + \tilde{K}_{JJ'}^2(t, t') \operatorname{Im} f_-^{J'}(t') \right\} \quad \forall J \geq 0 , \\ f_-^J(t) &= \tilde{N}_-^J(t) + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{l=0}^{\infty} \left\{ \tilde{H}_{Jl}(t, W') \operatorname{Im} f_{l+}^l(W') + \tilde{H}_{Jl}(t, -W') \operatorname{Im} f_{(l+1)-}^l(W') \right\} \\ &\quad + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \sum_{J'} \tilde{K}_{JJ'}^3(t, t') \operatorname{Im} f_-^{J'}(t') \quad \forall J \geq 1 , \end{aligned}$$

- **Validity:** assuming Mandelstam analyticity
⇒ optimal for $a = -2.71 M_\pi^2$

$$t \in [t_\pi = 4M_\pi^2, 205.45 M_\pi^2] \Leftrightarrow \sqrt{t} \in [\sqrt{t_\pi} = 0.28 \text{ GeV}, 2.00 \text{ GeV}] .$$

- Assumption: Mandelstam analyticity


Mandelstam (1958,1959)

- $\Rightarrow T(s,t)$ can be written in terms **double spectral densities**: ρ_{st} , ρ_{su} , ρ_{yt}


$$T(s, t) = \frac{1}{\pi^2} \iint ds' du' \frac{\rho_{su}(s', u')}{(s' - s)(u' - u)} + \frac{1}{\pi^2} \iint dt' du' \frac{\rho_{tu}(t', u')}{(t' - t)(u' - u)} + \frac{1}{\pi^2} \iint ds' dt' \frac{\rho_{st}(s', t')}{(s' - s)(t' - t)}$$

→ integration ranges defined by the support of the **double spectral** densities ρ

- Boundaries of ρ are given lowest lying intermediate states



- They limit the range of validity of the HDRS:



Solving t-channel: single channel

- Elastic-channel approximation: generic form of the integral equation

$$f(t) = \Delta(t) + (a + bt)(t - 4m^2) + \frac{t^2(t - 4m^2)}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\text{Im } f(t')}{t'(t'^2 - 4m^2)(t' - t)}$$

- $\Delta(t)$: Born terms, s-channel integrals, higher t -channel partial waves
⇒ left-hand cut
- Introduce subtractions at $\nu = t = 0$ ⇒ subthreshold parameters a, b
- Solution in terms of Omnès function:

$$\begin{aligned} f(t) &= \Delta(t) + (t - 4m^2)\Omega(t)(1 - t\dot{\Omega}(0))a + t(t - 4m^2)\Omega(t)b \\ &\quad - \Omega(t) \frac{t^2(t - 4m^2)}{\pi} \left\{ \int_{4M_\pi^2}^{t_m} dt' \frac{\Delta(t') \text{Im } \Omega(t')^{-1}}{t'^2(t' - 4m^2)(t' - t)} + \int_{t_m}^{\infty} dt' \frac{\Omega(t')^{-1} \text{Im } f(t')}{t'(t' - 4m^2)(t' - t)} \right\} \\ \Omega(t) &= \exp \left\{ \frac{t}{\pi} \int_{t_\pi}^{t_m} \frac{dt'}{t'} \frac{\delta(t')}{t' - t} \right\} \end{aligned}$$



Solving t-channel: input and subtractions

- elastic channel approximation: $\sqrt{t}_m = 0.98 - 1.1 \text{ GeV}$, for $t > t_m \operatorname{Im} f_{\pm}^J(t) = 0$
- First step: check **consistency** with KH80 Höhler 1983
- Input needed:
 - $\pi\pi$ phase shifts:
 - πN phase shifts: SAID, KH80
 - πN at high energies: Regge model
 - πN parameters: KH80


Caprini, Colangelo, Leutwyler, (in preparation), Madrid group

Arndt et al. 2008, Höhler 1983

Huang et al. 2010




Solving t-channel: P-wave results



MO solutions in general consistent with KH80 results

Solving t-channel: P, D and F waves up to $\bar{N}N$



Solving t-channel: coupled channels

- Generic coupled-channel integral equation

$$\mathbf{f}(t) = \Delta(t) + \frac{1}{\pi} \int_{t_\pi}^{t_m} dt' \frac{T^*(t') \Sigma(t') \mathbf{f}(t')}{t' - t} + \frac{1}{\pi} \int_{t_m}^{\infty} dt' \frac{\text{Im } \mathbf{f}(t')}{t' - t}$$

- Formal solution as in the single-channel case (now with Omnès matrix $\Omega(t)$)

⇒ Two-channel Muskhelishvili-Omnès problem

$$\mathbf{f}(t) = \begin{pmatrix} f_+^0(t) \\ h_+^0(t) \end{pmatrix} \quad \text{Im } \Omega(t) = (T(t))^* \Sigma(t) \Omega(t)$$

- Two linearly independent solutions Ω_1, Ω_2

Muskhelishvili 1953

- In general no analytical solution for the Omnès matrix
but for its determinant


Moussallam 2000

$$\det \Omega(t) = \exp \left\{ \frac{t}{\pi} \int_{t_\pi}^{t_m} dt' \frac{\psi(t')}{t'(t' - t)} \right\}.$$



Solving t-channel S-wave equations: input

- Input needed:
 - $\pi\pi$ s-wave partial waves: Caprini, Colangelo, Leutwyler, (in preparation)
 - $K\bar{K}$ s-wave partial waves: Büttiker. (2004)
 - πN and KN s-wave pw: SAID, KH80 Arndt et al. 2008, Höhler 1983,
 - πN at high energies: Regge model Huang et al. 2010
 - πN parameters: KH80
 - Hyperon couplings from Jülich model 1989
 - KN subthreshold parameters neglected
- Two-channel approximation breaks down at $\sqrt{t_0} = 1.3$ GeV $\Rightarrow 4\pi$ channel
- From t_0 to $t = 2$ GeV, different approximations considered

Solving t-channel: S-wave results



Solving t-channel: S-wave results



MO solutions in general consistent with KH80 results

Solving s-channel: S-wave results

- General form of the s-channel integral equation

$$f_{I+}^I(W) = \Delta_{I+}^I(W) + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{I'=0}^{\infty} \left\{ K_{II'}^I(W, W') \operatorname{Im} f_{I'+}^I(W') + K_{II'}^I(W, -W') \operatorname{Im} f_{(I'+1)-}^I(W') \right\}$$

⇒ form of $\pi\pi$ Roy-Equations

- $\Delta_{I+}^I(W)$ ≡ t-channel contribution and pole term

- valid up to $W_m = 1.38$ GeV

- Input:**

- RS t-channel solutions for S and P waves
- s-channel partial waves for $J > 1$
- s-channel partial waves for $W_m < W < 2.5$ GeV
- high energy contribution for $W > 2.5$ GeV: Regge model

SAID analysis

SAID analysis

Huang et al. 2010

- Output:**

- Self-consistent solution for S and P waves for $J \leq J_{\max}$ and $s_{\text{th}} \leq s \leq s_m$
- Constraints on subtraction constants ⇒ subthreshold parameters

Solving s-channel: subtractions

- Existence and uniqueness of solutions

Gasser, Wanders 1999

⇒ no-cusp condition for each pw + 2 additional constraints are needed

- Take advantage of the precise data for pionic atoms

Gotta et al. 2005, 2010

⇒ Impose as a **constraint** scattering lengths from a combined analysis of pionic hydrogen and deuterium

Baru et al. 2011

$$a_{0+}^{1/2} = (169.8 \pm 2.0) 10^{-3} M_\pi^{-1} \quad a_{0+}^{3/2} = (-86.3 \pm 1.8) 10^{-3} M_\pi^{-1}$$

$$\text{Re } f_{l\pm}^I(s) = \mathbf{q}^{2l} \left(a_{l\pm}^I + b_{l\pm}^I \mathbf{q}^2 + \dots \right)$$

10 subthreshold parameters are needed to match d.o.f
⇒ three subtractions

Solving s-channel: strategy

- Parameterize S and P waves up to $W < W_m$
 - Using SAID partial waves as starting point
- Impose as **constraints** the hadronic atom **scattering lengths**
- Introduce as many **subtractions** as necessary to **match d.o.f**
- Minimize difference between **LHS** and the **RHS** on a grid of points W_j



[Gasser, Wanders 1999]

$$\chi^2 = \sum_{l,l_s,\pm} \sum_{j=1}^N \frac{\left(\text{Re } f_{l\pm}^{ls}(W_j) - F[f_{l\pm}^{ls}](W_j) \right)^2}{\text{Re } f_{l\pm}^{ls}(W_j)}$$

$F[f_{l\pm}^{ls}](W_j)$ \equiv right hand side of RS-equations



- Parametrization and subthreshold parameters are the fitting parameters

Solving s-channel: results





blue/red
↔
LHS/RHS

after the fit





gray/black
↔
LHS/RHS
before the fit





Notation: $L_2 l_5 2J$



Results: s-channel PWs



blue/red
 \Updownarrow
 LHS/RHS



after the fit





gray/black
 \Updownarrow
 LHS/RHS



before the fit

Notation: $L_2 I_s 2J$



Results: s-channel PWs



sizable
 $f_2(1275)$ effect
 blue/red



\Updownarrow
LHS/RHS
 after the fit




\Updownarrow
LHS/RHS

before the fit

Notation: $L_2 l_5 2J$

Results: t-channel PWs



blue







before the fit

red



after the fit



KH80

Solving the full RS system: strategy


- Full solution: self-consistent, **iterative** solution of the **full RS** system
⇒ consistent set of **s**- and **t**-channel PWs & **low-energy parameters**
- However:
 - t-channel RS eqs. depend only weakly on s-channel PWs
 - resulting s-channel PW change little from **SAID**

A **full solution** can be achieved including in the **s-channel** RS eqs. the **t-channel** dependence on the **subthreshold parameters**


Uncertainties

- Statistical errors (at intermediate energies)
 - ▷ important correlations between subthreshold parameters
 - ▷ shallow fit minima
 - ⇒ Sum rules for subthreshold parameters become essential to reduce the errors
- Input variation (small)
 - ▷ small effect for considering s-channel KH80 input
 - ▷ very small effects from $L > 5$ s-channel PWs
 - ▷ small effect from the different S-wave extrapolation for $t > 1.3$ GeV
 - ▷ negligible effect of ρ' and ρ''
 - ▷ very significant effects of the D-waves ($f_2(1275)$)
 - ▷ F-waves shown to be negligible
- matching conditions (close to W_m)
- scattering length (SL) errors (on S-waves and subthreshold parameters)
 - ▷ very important for the $\sigma_{\pi N}$

Uncertainties: Real part s-channel pw




Uncertainties: Real part t-channel pw



Decomposition of s-channel RS pw



$$f_{I\pm}^{Is}(W) = N_{I\pm}^{Is}(W) + K_S T(W)_{I\pm}^{Is} + K_t T(W)_{I\pm}^{Is} + D T(W)_{I\pm}^{Is}$$



Comparison with KH80



- Karlsruhe-Helsinki analysis **KH80**
 - comprehensive analyticity constraints based on fixed-t dispersion relations
 - old experimental data
- Here, an update of **KH80** results with modern input
 - HDR increase the range of validity of the equations
 - πN scattering length extracted from hadronic atoms \Rightarrow essential for the $\sigma_{\pi N}$
 - Goldberger-Miyazawa-Oehme sum rule:
$$g_{\pi N}^2 / 4\pi = 13.7 \pm 0.2$$
 Baru et al. 2011
compare:
$$g_{\pi N}^2 / 4\pi = 14.28$$
 Höhler et al. 1983
- s-channel PWs from **SAID**
 - $f_2(1275)$ included \Rightarrow sizable effect
- **KH80** is internally consistent \Rightarrow RS reproduces **KH80** results with **KH80** input

Results: s-channel PWs with KH80 input





blue/red
 \Updownarrow
 LHS/RHS

after the fit




gray/black
 \Updownarrow
 LHS/RHS
 before the fit



Notation: $L_2 l_5 2J$

Results: t-channel PWs with KH80 input



Threshold parameters

- Threshold parameters defined as: $\text{Re } f_{I\pm}^I(s) = q^{2I} \{ a_{I\pm}^I + b_{I\pm}^I q^2 + \dots \}$
- Extracted from hyperbolic sum rules

	RS	KH80
a_{0+}^+ [$10^{-3} M_\pi^{-1}$]	-0.9 ± 1.4	-9.7 ± 1.7
a_{0+}^- [$10^{-3} M_\pi^{-1}$]	85.4 ± 0.9	91.3 ± 1.7
a_{1+}^+ [$10^{-3} M_\pi^{-3}$]	131.2 ± 1.7	132.7 ± 1.3
a_{1+}^- [$10^{-3} M_\pi^{-3}$]	-80.3 ± 1.1	-81.3 ± 1.0
a_{1-}^+ [$10^{-3} M_\pi^{-3}$]	-50.9 ± 1.9	-56.7 ± 1.3
a_{1-}^- [$10^{-3} M_\pi^{-3}$]	-9.9 ± 1.2	-11.7 ± 1.0
b_{0+}^+ [$10^{-3} M_\pi^{-3}$]	-45.0 ± 1.0	-44.3 ± 6.7
b_{0+}^- [$10^{-3} M_\pi^{-3}$]	4.9 ± 0.8	13.3 ± 6.0

- Reasonable agreement with KH80 but for the scattering lengths
- Disagreement in the scattering lengths in $\sim 4\sigma$

Threshold parameters


- Threshold parameters defined as: $\text{Re } f_{I\pm}^I(s) = q^{2I} \{ a_{I\pm}^I + b_{I\pm}^I q^2 + \dots \}$
- Extracted from hyperbolic sum rules

	RS	KH80
$a_{0+}^{1/2} [10^{-3} M_\pi^{-1}]$	169.8 ± 2.0	173 ± 3
$a_{0+}^{3/2} [10^{-3} M_\pi^{-1}]$	-86.3 ± 1.8	-101 ± 4
$a_{1+}^{1/2} [10^{-3} M_\pi^{-3}]$	-29.4 ± 1.0	-30 ± 2
$a_{1+}^{3/2} [10^{-3} M_\pi^{-3}]$	211.5 ± 2.8	214 ± 2
$a_{1-}^{1/2} [10^{-3} M_\pi^{-3}]$	-70.7 ± 4.1	-81 ± 2
$a_{1-}^{3/2} [10^{-3} M_\pi^{-3}]$	-41.0 ± 1.1	-45 ± 2
$b_{0+}^{1/2} [10^{-3} M_\pi^{-3}]$	-35.2 ± 2.2	-18 ± 12
$b_{0+}^{3/2} [10^{-3} M_\pi^{-3}]$	-49.8 ± 1.1	-58 ± 9

- Reasonable agreement with KH80 but for the scattering lengths
- Disagreement in the $a_0^{3/2}$ scattering length in $\sim 4\sigma$

Dispersion relation for the scalar form factor of the nucleon

- Unitarity relation:



$$\text{Im } \sigma(t) = \frac{2}{4m^2 - t} \left\{ \frac{3}{4} \sigma_t^\pi (\mathcal{F}_\pi^S(t))^* f_0^0(t) + \sigma_t^K (\mathcal{F}_K^S(t))^* h_0^0(t) \right\}$$



- Dispersion relation:

$$\sigma(t) = \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\text{Im}\sigma(t')}{(t' - t)} = \sigma_{\pi N} + \frac{t}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\text{Im}\sigma(t')}{t'(t' - t)}$$

Unsubtracted: $\sigma_{\pi N} = \sigma(0)$

Once-subtracted: $\Delta\sigma = \sigma(2M_\pi^2) - \sigma_{\pi N}$

Convergence of the dispersive integral



- **Unsubtracted:** slow convergence
- **Once-subtracted:** stable result for $\Lambda \gtrsim 1$ GeV

Dispersion relation for the πN amplitude

- t-channel expansion of the subtracted pseudo-Born amplitude

$$\bar{D}(\nu = 0, t) = 4\pi \left\{ \frac{1}{p_t^2} \bar{f}_0^+(t) + \frac{5}{2} q_t^2 \bar{f}_2^+(t) + \frac{27}{8} p_t^2 q_t^4 \bar{f}_4^+(t) + \frac{56}{16} p_t^4 q_t^6 \bar{f}_6^+(t) + \dots \right\}$$

- Insert t-channel RS equations for Born-term-subtracted amplitudes $\bar{f}_J^+(t)$

$$\bar{D}(\nu = 0, t) = d_{00}^+ + d_{01}^+ t - 16t^2 \int_{t_\pi}^{\infty} dt' \frac{\text{Im}\bar{f}_0^+(t')}{t'^2(t' - 4m^2)(t' - t)} + \{J \geq 2\} + \{\text{s-channel integral}\}$$

- $\Delta_D = F_\pi^2 (\bar{D}(\nu = 0, t) - d_{00}^+ + d_{01}^+ t)$ from evaluation at $t = 2M_\pi^2$

Summary: σ -term corrections

- Nucleon scalar form factor

$$\Delta_{\sigma} = (13.9 \pm 0.3) \text{ MeV}$$

$$+ Z_1 \left(\frac{g^2}{4\pi} - 14.28 \right) + Z_2 \left(d_{00}^+ M_\pi + 1.46 \right) + Z_3 \left(d_{01}^+ M_\pi^3 - 1.14 \right) + Z_4 \left(b_{00}^+ M_\pi^3 + 3.54 \right)$$

$$Z_1 = 0.36 \text{ MeV}, \quad Z_2 = 0.57 \text{ MeV}, \quad Z_3 = 12.0 \text{ MeV}, \quad Z_4 = -0.81 \text{ MeV}$$

- πN amplitude

$$\Delta_D = (12.1 \pm 0.3) \text{ MeV}$$

$$+ \hat{Z}_1 \left(\frac{g^2}{4\pi} - 14.28 \right) + \hat{Z}_2 \left(d_{00}^+ M_\pi + 1.46 \right) + \hat{Z}_3 \left(d_{01}^+ M_\pi^3 - 1.14 \right) + \hat{Z}_4 \left(b_{00}^+ M_\pi^3 + 3.54 \right)$$

$$\hat{Z}_1 = 0.42 \text{ MeV}, \quad \hat{Z}_2 = 0.67 \text{ MeV}, \quad \hat{Z}_3 = 12.0 \text{ MeV}, \quad \hat{Z}_4 = -0.77 \text{ MeV}$$

→ most of the dependence on the πN parameters cancels in the difference

Full Correction

$$\Delta_D - \Delta_{\sigma} = (-1.8 \pm 0.2) \text{ MeV}$$

Cheng-Dashen theorem in the presence of isospin breaking

- Define as **isoscalar** as

$$X^+ \rightarrow X^P = \frac{1}{2}(X_{\pi^+ p \rightarrow \pi^+ p} + X_{\pi^- p \rightarrow \pi^- p}), \quad X \in \{D, d_{00}, d_{01}, a_0, \dots\}$$

and “**isospin limit**” by proton and charged pion

- Assume virtual photons to be removed

↪ scenario closest to actual πN PWA

- Calculate **IV corrections** in SU(2) ChPT, mainly due to $\Delta_\pi = M_\pi^2 - M_{\pi^0}^2$

- For the σ term no differences at $\mathcal{O}(p^3)$

$$\sigma_{\pi N} = \sigma_p = \sigma_N = -4c_1 M_{\pi^0}^2 - \frac{3g_A^2 M_{\pi^0}^2}{64\pi F_\pi^2} (2M_\pi + M_{\pi^0}) + \mathcal{O}(M_\pi^4)$$

- Slope of the scalar form factor

$$\Delta_\sigma^p = \sigma_p(2M_\pi^2) - \sigma_p = \frac{3g_A^2 M_\pi^3}{64\pi F_\pi^2} + \frac{g_A^2 M_\pi \Delta_\pi}{128\pi F_\pi^2} (-7 + \sqrt{2} \log(3 + 2\sqrt{2})) + \mathcal{O}(M_\pi^4)$$

- Similarly for Δ_D^p

$$\Delta_D^p = F_\pi^2 \left\{ \bar{D}_p(0, 2M_\pi^2) - d_{00}^p - 2M_\pi^2 d_{01}^p \right\} = \frac{23g_A^2 M_\pi^3}{384\pi F_\pi^2} + \frac{g_A^2 M_\pi \Delta_\pi}{256\pi F_\pi^2} (3 + 4\sqrt{2} \log(1 + \sqrt{2})) + \mathcal{O}(M_\pi^4)$$

Cheng-Dashen theorem in the presence of isospin breaking

- Taking everything together

$$\begin{aligned}\sigma_{\pi N} &= F_\pi^2 (d_{00}^p + 2M_\pi^2 d_{01}^p) - \Delta_R + \Delta_D - \Delta_\sigma + (\Delta_D^p - \Delta_D) - (\Delta_\sigma^p - \Delta_\sigma) \\ &\quad + \sigma_p(2M_\pi^2) + F_\pi^2 \bar{D}(0, 2M_\pi^2) \\ &= F_\pi^2 (d_{00}^p + 2M_\pi^2 d_{01}^p) - \underbrace{\Delta_R}_{\lesssim 2 \text{ MeV}} + \underbrace{\Delta_D - \Delta_\sigma}_{(-1.8 \pm 0.2) \text{ MeV}} + \underbrace{\frac{81g_a^2 M_\pi \Delta_\pi}{256\pi F_\pi^2}}_{3.4 \text{ MeV}} + \underbrace{\frac{e^2}{2} F_\pi^2 (4f_1 + f_2)}_{(-0.4 \pm 2.2) \text{ MeV}}\end{aligned}$$

↪ sizable corrections from Δ_π increasing the value of the $\sigma_{\pi N}$


- Electromagnetic nucleon form factor: $\langle N(p') | j_{\text{em}}^\mu | N(p) \rangle = \bar{u}(p') \left[F_1^N(t) \gamma^\mu + \frac{i \sigma^{\mu\nu} q_\nu}{2m_N} F_2^N(t) \right] u(p),$

$$G_E^N(t) = F_1^N(t) + \frac{t}{4m_N^2} F_2^N(t), \quad G_M^N(t) = F_1^N(t) + F_2^N(t).$$

- first inelastic correction from $\pi\pi$ continuum

$$\text{Im } G_E^V(t) = \frac{q_t^3}{m_N \sqrt{t}} (\mathcal{F}_\pi^V(t))^* f_+^1(t) \theta(t - t_\pi)$$

$$\text{Im } G_M^V(t) = \frac{q_t^3}{\sqrt{2t}} (\mathcal{F}_\pi^V(t))^* f_-^1(t) \theta(t - t_\pi)$$



→ rigorous constraint fixed from:

- ▷ RS t-channel partial waves
- ▷ pion form factor


- update of Höhler spectral functions, including also isospin breaking

criticism by Lee et al. 2015

$\pi\pi$ -continuum: $\rho - \omega$ mixing

- Isovector and isoscalar nucleon form factor

$$F_i^s(t) = \frac{1}{2}(F_i^p(t) + F_i^n(t)), \quad F_i^\nu(t) = \frac{1}{2}(F_i^p(t) - F_i^n(t))$$



$$\begin{aligned} \text{Im } G_E^\nu(t) &= \frac{q_t^3}{m_N \sqrt{t}} |\Omega_1^1(t)| |f_+^1(t)| \theta(t - t_\pi) \\ &\times \left(1 + \alpha t + \frac{\varepsilon t}{M_\omega^2 + i M_\omega \Gamma_\omega - t} \right) \\ &+ \varepsilon \text{Im} \left(\frac{t}{M_\omega^2 - i M_\omega \Gamma_\omega - t} \right) \\ &\times \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\frac{q_t'^3}{m_N \sqrt{t'}} |\Omega_1^1(t')| |f_+^1(t')|}{t' - t - i\varepsilon}, \\ \text{Im } G_M^\nu(t) &= \frac{q_t^3}{\sqrt{2}t} |\Omega_1^1(t)| |f_-^1(t)| \theta(t - t_\pi) \\ &\times \left(1 + \alpha t + \frac{\varepsilon t}{M_\omega^2 + i M_\omega \Gamma_\omega - t} \right) \\ &+ \varepsilon \text{Im} \left(\frac{t}{M_\omega^2 - i M_\omega \Gamma_\omega - t} \right) \\ &\times \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\frac{q_t'^3}{\sqrt{2}t'} |\Omega_1^1(t')| |f_-^1(t')|}{t' - t - i\varepsilon}. \end{aligned}$$

Chiral Low Energy Constants with Δ 's

	HB- NN		HB- πN		covariant	
N^2LO	Q^3	ε^3	Q^3	ε^3	Q^3	ε^3
c_1	-1.08(2)	-1.25(3)	-1.08(2)	-1.24(3)	-1.00(2)	-1.19(4)
c_2	3.26(3)	1.71(1.01)	3.26(3)	1.13(1.02)	2.55(3)	1.14(19)
c_3	-5.39(5)	-2.68(84)	-5.39(5)	-2.75(84)	-4.90(5)	-2.56(40)
c_4	3.62(3)	1.57(16)	3.62(3)	1.58(16)	3.08(3)	1.33(20)
d_{1+2}	1.02(6)	0.14(17)	1.02(6)	-0.07(18)	1.78(6)	0.62(16)
d_3	-0.46(2)	-0.84(14)	-0.46(2)	-0.48(15)	-1.12(2)	-1.45(5)
d_5	0.15(5)	0.80(7)	0.15(5)	0.47(6)	-0.05(5)	0.29(6)
d_{14-15}	-1.85(6)	-1.09(30)	-1.85(6)	-0.72(31)	-2.27(6)	-0.98(13)
N^3LO	Q^4	ε^4	Q^4	ε^4	Q^4	ε^4
c_1	-1.11(3)	-1.11(3)	-1.11(3)	-1.11(3)	-1.12(3)	-1.10(3)
c_2	3.61(4)	1.41(38)	3.17(3)	1.28(20)	3.35(3)	1.16(20)
c_3	-5.60(6)	-1.88(45)	-5.67(6)	-2.04(39)	-5.70(6)	-2.10(39)
c_4	4.26(4)	2.03(28)	4.35(4)	2.07(29)	3.97(3)	1.91(27)
d_{1+2}	6.37(9)	1.78(31)	7.66(9)	2.90(30)	4.70(7)	1.78(24)
d_3	-9.18(9)	-3.64(36)	-10.77(10)	-5.91(50)	-5.26(5)	-3.25(14)
d_5	0.87(5)	1.52(7)	0.59(5)	1.03(7)	0.31(5)	0.66(6)
d_{14-15}	-12.56(12)	-4.38(54)	-13.44(12)	-5.17(55)	-8.84(10)	-3.41(41)
e_{14}	1.16(4)	1.64(10)	0.85(4)	1.12(16)	1.17(4)	1.28(11)
e_{15}	-2.26(6)	-4.95(15)	-0.83(6)	-3.30(25)	-2.58(7)	-3.07(13)
e_{16}	-0.29(3)	4.21(16)	-2.75(3)	1.92(43)	-1.77(3)	1.71(17)
e_{17}	-0.17(6)	-0.44(6)	0.03(6)	-0.39(7)	-0.45(6)	-0.51(7)
e_{18}	-3.47(5)	1.34(29)	-4.48(5)	0.67(31)	-1.68(5)	1.30(17)

Threshold kinematics from subthreshold with Δ 's

	HB- NN		HB- πN		covariant		RS
	N ² LO	Q^3	ε^3	Q^3	ε^3	Q^3	ε^3
$a_{0+}^+[10^{-3}M_\pi^{-1}]$	0.5	-9.8(10.9)	0.5	-0.4(9.2)	-14.8	1.0(17.3)	-0.9(1.4)
$a_{0+}^-[10^{-3}M_\pi^{-1}]$	92.2	92.7(1.0)	92.9	90.5(9)	89.9	81.7(1.6)	85.4(9)
$a_{1+}^+[10^{-3}M_\pi^{-3}]$	113.8	125.8(16.7)	121.7	127.2(18.4)	116.4	128.5(9.6)	131.2(1.7)
$a_{1+}^-[10^{-3}M_\pi^{-3}]$	-74.8	-77.4(2.5)	-75.5	-78.4(2.6)	-75.1	-79.7(3.0)	-80.3(1.1)
$a_{1-}^+[10^{-3}M_\pi^{-3}]$	-54.1	-53.4(14.1)	-47.0	-52.5(15.8)	-55.5	-52.5(8.5)	-50.9(1.9)
$a_{1-}^-[10^{-3}M_\pi^{-3}]$	-14.1	-13.1(2.7)	-2.5	-7.8(3.0)	-10.4	-9.7(4.1)	-9.9(1.2)
$b_{0+}^+[10^{-3}M_\pi^{-3}]$	-45.7	-38.1(9.6)	-22.1	-23.7(14.4)	-50.9	-34.7(12.1)	-45.0(1.0)
$b_{0+}^-[10^{-3}M_\pi^{-3}]$	35.9	26.4(1.0)	22.6	17.6(8)	21.6	14.2(2.0)	4.9(8)
	N ³ LO	Q^4	ε^4	Q^4	ε^4	Q^4	ε^4
$a_{0+}^+[10^{-3}M_\pi^{-1}]$	-1.5	-1.5(8.5)	-8.0	1.2(20.4)	-5.7	-0.8(10.3)	-0.9(1.4)
$a_{0+}^-[10^{-3}M_\pi^{-1}]$	68.5	96.3(2.0)	58.6	70.0(3.3)	83.8	83.6(1.9)	85.4(9)
$a_{1+}^+[10^{-3}M_\pi^{-3}]$	134.3	136.0(9.7)	132.1	135.2(8.7)	128.0	132.7(9.0)	131.2(1.7)
$a_{1+}^-[10^{-3}M_\pi^{-3}]$	-80.9	-80.0(3.4)	-90.1	-86.4(2.7)	-78.1	-81.1(3.6)	-80.3(1.1)
$a_{1-}^+[10^{-3}M_\pi^{-3}]$	-55.7	-47.5(10.5)	-73.7	-56.9(7.1)	-53.5	-51.4(7.9)	-50.9(1.9)
$a_{1-}^-[10^{-3}M_\pi^{-3}]$	-10.0	-5.6(4.9)	-23.7	-14.4(6.5)	-11.8	-10.4(5.7)	-9.9(1.2)
$b_{0+}^+[10^{-3}M_\pi^{-3}]$	-42.2	-31.4(8.1)	-44.5	-32.6(21.3)	-54.7	-33.9(8.5)	-45.0(1.0)
$b_{0+}^-[10^{-3}M_\pi^{-3}]$	-31.6	7.1(2.3)	-65.2	-34.1(5.7)	2.3	2.9(2.1)	4.9(8)

Goldberger-Miyazawa-Oehme sum rule

- Fixed- t dispersion relations at threshold \hookrightarrow **GMO sum rule**

$$\frac{g^2}{4\pi} = \left(\left(\frac{m_p + m_n}{M_\pi} \right)^2 - 1 \right) \left\{ \left(1 + \frac{M_\pi}{m_p} \right) \frac{M_\pi}{4} (a_{\pi^- p} - a_{\pi^+ p}) - \frac{M_\pi^2}{2} J^- \right\}$$
$$= 13.69 \pm 0.12 \pm 0.15$$

$$J^- = \frac{1}{4\pi^2} \int_0^\infty dk \frac{\sigma_{\pi^- p}^{\text{tot}}(k) - \sigma_{\pi^+ p}^{\text{tot}}(k)}{\sqrt{M_\pi^2 + k^2}}$$

- J^- known very accurately

Ericson et al. 2002, Abaev et al. 2007

- other determinations

	de Swart et al. 97	Arndt et al. 94	Ericson et al. 02	Bugg et al. 73	KH80
method	NN	πN	GM0	πN	πN
$g^2/4\pi$	13.54 ± 0.05	13.75 ± 0.15	14.11 ± 0.20	14.30 ± 0.18	14.28

- With KH80 scattering lengths $g^2/4\pi = 14.28$ is reproduced exactly

\hookrightarrow discrepancy related to old scattering length values

Trace anomaly

- Classical massless QCD is invariant under scale transformations

$$x \rightarrow \lambda x \quad q(x) \rightarrow \lambda^{3/2} q(\lambda x) \quad A(x) \rightarrow \lambda A(\lambda x) \quad (\text{dilatations})$$

- Quantization/renormalization generates a scale Λ_{QCD} that breaks scale invariance:

→ dimensional transmutation

⇒ Trace anomaly

$$\theta_\mu^\mu = \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G_a^{\mu\nu} + m_u \bar{u}u + m_d \bar{d}d + m_s \bar{s}s + \dots$$

- trace anomaly ≡ signal for the generation of hadron masses
- the mass of any hadron made of light quarks mass is essentially field energy

"Mass without a mass" [Wheeler, 1962]

The role of the pion-nucleon σ -term

- scalar coupling of the nucleon

$$\langle N | m_q \bar{q} q | N \rangle = f_q m_N, \quad f_q = \frac{\sigma_{\pi N}}{2m_N} \quad N \in \{p, n\}$$

↪ Dark Matter detection

↪ $\mu \rightarrow e$ conversion in nuclei



- Condensates in nuclear matter

$$\frac{\langle \bar{q} q \rangle(\rho)}{\langle 0 | \bar{q} q | 0 \rangle} = 1 - \frac{\rho \sigma_{\pi N}}{F_\pi^2 M_\pi^2}$$

- CP violation in πN couplings

↪ hadronic EDMs

$$g_{\eta N} \propto - \left(\frac{\sigma_s}{m_s} - \frac{\sigma_{\pi N}}{2\hat{m}} \right) \bar{\theta}$$



- relate $\sigma_{\pi N}$ to strangeness content of the nucleon:

$$\sigma_{\pi N} = \frac{\hat{m}}{2m_N} \frac{\langle N|\bar{u}u + \bar{d}d - 2\bar{s}s|N\rangle}{1 - y} = \frac{\sigma_0}{1 - y}, \quad y \equiv \frac{2\langle N|\bar{s}s|N\rangle}{\langle N|\bar{u}u + \bar{d}d|N\rangle}$$

$(m_s - \hat{m})(\bar{u}u + \bar{d}d - 2\bar{s}s) \subset$ LQCD produces SU(3) mass splittings:

$$\sigma_{\pi N} = \frac{\sigma_0}{1 - y}, \quad \sigma_0 = \frac{\hat{m}}{m_s - \hat{m}} (m_{\Xi} + m_{\Sigma} - 2m_N) \sim 26 \text{ MeV}$$

higher-order corrections: $\sigma_0 \rightarrow (36 \pm 7) \text{ MeV}$

[Borasoy, Meißner 1997]

- potentially large effects

- from the decuplet
- from relativistic corrections (EOMS vs. heavy-baryon)
- may increase to $\sigma_0 = (58 \pm 8) \text{ MeV}$

[Alarcon et al. 2013, Siemens et al. 2016]

- Conclusion:**

- $\sigma_{\pi N} = (59.1 \pm 3.5) \text{ MeV}$ not incompatible with small y
- chiral convergence of σ_0 (hence $\langle N|\bar{s}s|N\rangle$) very doubtful

Spin-independent WIMP–nucleon scattering

- Effective Lagrangian

$$\mathcal{L} = C_{qq}^{SS} \frac{m_q}{\Lambda^3} \bar{\chi} \chi \bar{q} q + C_{qq}^{VV} \frac{m_q}{\Lambda^2} \bar{\chi} \gamma^\mu \chi \bar{q} \gamma_\mu q + \bar{C}_{gg}^S \frac{\alpha_S}{\Lambda^3} \bar{\chi} \chi G_{\mu\nu} G^{\mu\nu}$$

- WIMP χ **Dirac fermion** and **SM singlet**
- Spin-independent cross section at vanishing momentum transfer



$$\sigma_N^{SI} = \frac{\mu_\chi^2}{\Lambda^4} \left| \left(\frac{m_N}{\Lambda} C_{qq}^{SS} f_q^N - 12\pi C_{gg}^S f_Q^N \right) + C_{qq}^{VV} f_V^N \right|^2$$

$$\mu_\chi = \frac{m_\chi m_N}{m_\chi + m_N} \quad f_q^N = 2 \quad f_q^N = \frac{\sigma_{\pi N}(1-\xi)}{m_N} + \Delta f_q^N$$

- nucleon-matrix elements dominated by $\sigma_{\pi N}$

Roy-equations: $\pi\pi$ results

Solution for the $\pi\pi$ S0-wave



Garcia-Martin, Kaminski, Pelaez, JRE (2011)