




3D Nucleon Tomography

Mar 15th 2017

F.-X. Girod

GPD program and confinement studies in Hall B

Jefferson Lab, Newport News VA




Outline

- 1 Theoretical Context
- 2 Published Data
- 3 Future Measurements
- 4 Projected Impact on GPD Extractions
- 5 Summary and Outlook

Theoretical Context

Deep Exclusive Scattering

Generalized Parton Distributions



$$\gamma^* p \rightarrow \gamma p', \rho p', \omega p', \phi p'$$

Bjorken regime :

$$Q^2 \rightarrow \infty, x_B \text{ fixed}$$

$$t \text{ fixed} \ll Q^2, \xi \rightarrow \frac{x_B}{2-x_B}$$

$$\frac{P^+}{2\pi} \int dy^- e^{ixP^+y^-} \langle p' | \bar{\psi}_{\mathbf{q}}(0) \gamma^+ (1 + \gamma^5) \psi(y) | p \rangle$$

$$= \bar{N}(p') \left[\tilde{H}^q(x, \xi, t) \gamma^+ + \tilde{E}^q(x, \xi, t) i\sigma^{+\nu} \frac{\Delta_\nu}{2M} + \tilde{H}^q(x, \xi, t) \gamma^+ \gamma^5 + \tilde{E}^q(x, \xi, t) \gamma^5 \frac{\Delta^+}{2M} \right] N(p)$$

spin	N no flip	N flip
q no flip	H	E
q flip	$\tilde{H}$	$\tilde{E}$


3-D Imaging conjointly in transverse impact parameter **red** and longitudinal momentum

GPDs and Transverse Imaging


(x_B, t) correlations

$$q_{\mathbf{x}}(x, \vec{b}_{\perp}) = \int \frac{d^2 \vec{\Delta}_{\perp}}{(2\pi)^2} \left[H(x, 0, t) - \frac{E(x, 0, t)}{2M} \frac{\partial}{\partial b_y} \right] e^{-i \vec{\Delta}_{\perp} \cdot \vec{b}_{\perp}}$$

Target polarization



Lattice calculation



GPDs and Energy Momentum Tensor

(x, ξ) correlations

Form Factors accessed via second x-moments :

$$\langle p' | \hat{T}_{\mu\nu}^q | p \rangle = \bar{N}(p') \left[M_2^q(t) \frac{P_\mu P_\nu}{M} + J^q(t) \frac{i(P_\mu \sigma_{\nu\rho} + P_\nu \sigma_{\mu\rho}) \Delta^\rho}{2M} + d_1^q(t) \frac{\Delta_\mu \Delta_\nu - g_{\mu\nu} \Delta^2}{5M} \right] N(p)$$

Angular momentum distribution


$$J^q(t) = \frac{1}{2} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Mass and force/pressure distributions

$$M_2^q(t) + \frac{4}{5} d_1(t) \xi^2 = \frac{1}{2} \int_{-1}^1 dx x H^q(x, \xi, t)$$



$$d_1(t) = 15M \int d^3\vec{r} \frac{j_0(r\sqrt{-t})}{2t} p(r)$$

Distribution of pressure
 $r^2 p(r)$ in GeV fm^{-1}



Deeply Virtual Compton Scattering

The cleanest GPD probe at low and medium energies



$$ep \rightarrow ep\gamma$$

$$\sigma(ep \rightarrow ep\gamma) \propto \left| \text{DVCS} + \text{BH} \right|^2$$

(a) (b) (c)


$$A_{LU} = \frac{d^4\sigma^{\rightarrow} - d^4\sigma^{\leftarrow}}{d^4\sigma^{\rightarrow} + d^4\sigma^{\leftarrow}} \stackrel{\text{twist-2}}{\approx} \frac{\alpha \sin \phi}{1 + \beta \cos \phi}$$

$$\alpha \propto \text{Im} \left(F_1 \mathcal{H} + \xi G_M \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E} \right)$$

$$\mathcal{H}(\xi, t) = i\pi H(\xi, \xi, t) + \mathcal{P} \int_{-1}^1 dx \frac{H(x, \xi, t)}{x - \xi}$$


$$A_{UL} \propto \text{Im} \left(F_1 \tilde{\mathcal{H}} + \xi G_M \mathcal{H} + G_M \frac{\xi}{1 + \xi} \mathcal{E} + \dots \right) \sin \phi$$

Observables sensitivities to GPD



DVCS

	$\mathcal{I}m$	$\mathcal{R}e$
$\mathcal{H}$	A_{LU}	σ A_{LL}, A_{LT}
$\tilde{\mathcal{H}}$	A_{UL}	
$\mathcal{E}$	A_{UT}	




DVMP

	Meson	Flavor
$\mathcal{H}_{T,\mathcal{E}T}$	π^+	$\Delta u - \Delta d$
	π^0	$2\Delta u + \Delta d$
	η	$2\Delta u - \Delta d + 2\Delta s$
$\mathcal{H}, \mathcal{E}$	ρ^+	$u - d$
	ρ^0	$2u + d$
	ω	$2u - d$
	ϕ	g

A global analysis is needed to fully disentangle GPDs

Published Data




$$F_1 \mathcal{H} + \xi G_M \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E}$$

Precision in a large phase-space (x_B , Q^2 , t)

Qualitative model agreement

quantitative constraints on parameters

DVCS Unpolarized Cross-Sections 6 GeV





$$\bullet \quad \frac{d^4\sigma_{ep \rightarrow e\gamma}}{dQ^2 dx_B dt d\Phi} \text{ (nb/GeV}^4\text{)}$$

— BH — VGG (H only)
 KM10 --- KM10a

VGG: Vanderhaeghen, Guichon, Guidal



KM: Kumericki, Mueller





The t -slope becomes flatter with increasing x_B :
valence quarks (higher x_B) at the center of the nucleon and sea quarks (small x_B) at its periphery

$$F_1 \tilde{\mathcal{H}} + \xi G_M \left(\mathcal{H} + \frac{\xi}{1+\xi} \mathcal{E} \right)$$



Model independent extraction

Using only A_{LU} and A_{UL}


6 GeV

GPD dependencies versus x_B mirror their respective ordinary PDFs

$\tilde{H}$ and $H \leftrightarrow \Delta q(x)$ and $q(x)$

Change of $\Delta q(x)$ t-slope vs x_B less pronounced than $q(x)$

Axial charge more concentrated than EM charge




Future Measurements

CLAS12 GPD program

Number	Title	Contact	Days	Energy	Target
E12-06-108	Hard Exclusive Electroproduction of π^0 and η	Stoler	80	11	IH ₂
E12-06-119	Deeply Virtual Compton Scattering	Sabatie	80	11	IH ₂
E12-12-001	Timelike Compton Scat. & J/ ψ prod. in e^+e^-	Nadel-Turonski	120	11	IH ₂
E12-12-007	Exclusive ϕ meson electroproduction	Stoler	60	11	IH ₂
E12-11-003	DVCS on Neutron Target	Niccolai	90	11	ID ₂
E12-06-119	Deeply Virtual Compton Scattering	Sabatie	120	11	NH ₃
C12-12-010	DVCS with a transverse target	Elouadrhiri	110	11	HD-ice
E12-16-010	DVCS with CLAS12 at 6.6 GeV and 8.8 GeV	Elouadrhiri	50+50	6.6 & 8.8	IH ₂

80 days @ $\mathcal{L} = 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ with 85% polarized beam

$$A_{LU} \propto F_1 \mathcal{H} + \xi G_M \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E}$$




Beam Spin Asymmetries
 ϕ dependence

Statistical uncertainties :
from 1 % (low Q^2)
to 10 % (high Q^2)

Unprecedented statistics
over the full ϕ range
up to high $x = 0.6$

80 days @ $\mathcal{L} = 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ with 85% polarized beam

$$A_{LU} \propto F_1 \mathcal{H} + \xi G_M \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E}$$




Beam Spin Asymmetries
 ϕ dependence

Statistical uncertainties :
from 1 % (low Q^2)
to 10 % (high Q^2)

Unprecedented statistics
over the full ϕ range
up to high $x = 0.6$

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3

$$A_{UL} \propto F_1 \tilde{\mathcal{H}} + \xi G_M \left(\mathcal{H} + \frac{\xi}{1+\xi} \mathcal{E} \right) - \dots$$




Target Spin Asymmetries
 ϕ dependence

Statistical uncertainties :
from 2 % (low Q^2)
to 30 % (high Q^2)

Unprecedented statistics
over the full ϕ range
up to high $x = 0.6$

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3

$$A_{UL} \propto F_1 \tilde{\mathcal{H}} + \xi G_M \left(\mathcal{H} + \frac{\xi}{1+\xi} \mathcal{E} \right) - \dots$$




Target Spin Asymmetries
 ϕ dependence

Statistical uncertainties :
from 2 % (low Q^2)
to 30 % (high Q^2)

Unprecedented statistics
over the full ϕ range
up to high $x = 0.6$

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3


$$A_{UL} \propto F_1 \tilde{\mathcal{H}} + \xi G_M \left(\mathcal{H} + \frac{\xi}{1+\xi} \mathcal{E} \right) - \dots$$



TSA t-slopes

Sample kinematics
for target asymmetry

Change of t -slope with x_B
 $\leftrightarrow$
imaging $\Delta q(x_B, b_\perp)$




- Large glue density at $x > 0.1$

PDF from global fits

(F_2 evolution, ν_{DIS} , jets)

Gluons carry more than 30%
of the momentum for $0.1 < x$





- 3D imaging of the nucleon



spatial distribution of valence quarks :
elastic scattering, DVCS, ...

Nucleon gluonic radius ?
exclusive ϕ

Extraction of gluonic profiles



Longitudinal cross-section



Corresponding sensitivity in transverse position space


$$b = 1/\sqrt{-t}$$

Error propagation study
Skewness $\xi \neq 0$ neglected

Projected Impact on GPD Extractions


Projected Impact on GPD Extraction

Using simulated data
based on VGG model.
Input GPD H extracted
with good accuracy




Projected Impact on GPD Extraction

Using simulated data based on VGG model.
Input GPD H extracted with good accuracy



Projected Impact on GPD Extraction

Using simulated data based on VGG model.
Input GPD H extracted with good accuracy




Projection for the Nucleon transverse profile

Model profile


Projected error band

$$Q^2 = 3.75 \text{ GeV}^2$$



Precision tomography in the valence region

Global Fits to extract the D-term




Beam Spin Asymmetries

$$\text{Im}\mathcal{H}(\xi, t) = \frac{r}{1+x} \left(\frac{2\xi}{1+\xi} \right)^{-\alpha(t)} \left(\frac{1-\xi}{1+\xi} \right)^b \left(\frac{1-\xi}{1+\xi} \frac{t}{M^2} \right)^{-1}$$

Unpolarized cross-sections

$$\text{Re}\mathcal{H}(\xi, t) = D + \mathcal{P} \int dx \left(\frac{1}{\xi - x} - \frac{1}{\xi + x} \right) \text{Im}\mathcal{H}(\xi, t)$$




Green : pure Bethe-Heitler

Blue : local fit + uncertainty range

Red : resulting global fit

D-term and Pressure distribution




$$D(\frac{x}{\xi}, t) = \left(1 - \frac{x^2}{\xi^2}\right) \left[d_1(t) C_1^{3/2}(\frac{x}{\xi}) + d_3(t) C_3^{3/2}(\frac{x}{\xi}) + \dots \right]$$

t-dependence of the D-term : unknown

Dipole

Exponential

...



Resulting pressure distribution

$$\text{Stability condition : } \int_0^{\infty} dt \, r^2 p(r) = 0$$




World data fit

CLAS 6 GeV data

Projected CLAS12 data



Summary and Outlook

- Transverse Imaging, Energy Momentum Tensor
- First Generation Experiments Published
- Entering the 12 GeV and High Precision era
-



Supplementary material

Exclusive π^0 production at $W > 2$ 6 GeV



Sensitivity to the chiral-odd GPDs H_T and $2\tilde{H}_T + E_T$

The t -slope parameter is found independent of Q^2 and decreasing with x_B