

A New Approach to Calculating Magnetized Dynamical Friction for JLEIC

David Bruhwiler, Stephen Webb & Dan T. Abell




JLEIC Collaboration Meeting

5 April 2017 – Newport News, VA

This work is supported by the US DOE, Office of Science, Office of Nuclear Physics,
under Award # DE-SC0015212, with partial support from RadiaSoft LLC


Goals

- Simulate magnetized friction force
 - include all relevant real world effects
 - e.g. incoming beam distribution
 - include a wide range of parameters
 - cannot succeed via brute force
 - new theory is required



from Geller & Weisheit, Phys. Plasmas (1977)

- Include key aspects of magnetized e- beam transport
 - imperfect magnetization
 - space charge
 - field errors



from Zhang et al., MEIC design, arXiv (2012)



Hamiltonian for 2-body magnetized collision

$$H(\vec{x}_{ion}, \vec{p}_{ion}, \vec{x}_e, \vec{p}_e) = H_0(\vec{p}_{ion}, y_e, \vec{p}_e) + H_C(\vec{x}_{ion}, \vec{x}_e)$$

$$\vec{B} = B_0 \hat{z} \quad \vec{A} = -B_0 y \hat{x} \quad p_{e,x} = m_e (v_{e,x} - \Omega_L y_e)$$

$$H_0(\vec{p}_{ion}, y_e, \vec{p}_e) = \frac{1}{2m_{ion}} (p_{ion,x}^2 + p_{ion,y}^2 + p_{ion,z}^2) + \frac{1}{2m_e} \left[(p_{e,x} + eB_0 y_e)^2 + p_{e,y}^2 + p_{e,z}^2 \right]$$

$$H_C(\vec{x}_{ion}, \vec{x}_e) = \frac{-Ze^2}{4\pi\epsilon_0} \frac{1}{\sqrt{(x_{ion} - x_e)^2 + (y_{ion} - y_e)^2 + (z_{ion} - z_e)^2}}$$

Resulting equations of motion, in the standard drift-kick symplectic form:

$$M(\Delta t) = M_0(\Delta t/2)M_C(\Delta t)M_0(\Delta t/2)$$

D.L. Bruhwiler and S.D. Webb, “New algorithm for dynamical friction of ions in a magnetized electron beam,” in *AIP Conf. Proc.* **1812**, 050006 (2017); <http://aip.scitation.org/doi/abs/10.1063/1.4975867>



Symplectic drift map for 2-body system

$$M_0(\Delta t): \quad \vec{p}_{ion} = \text{constant} \quad p_{e,x} = \text{constant} \quad p_{e,z} = \text{constant}$$

$$\vec{x}_{ion}(t + \Delta t) = \vec{x}_{ion}(t) + \frac{\vec{p}_{ion}(t)}{m_{ion}} \Delta t \quad z_e(t + \Delta t) = z_e(t) + \frac{p_{e,z}(t)}{m_e} \Delta t$$

$$x_e(t + \Delta t) = x_e(t) + r_L [\cos(\varphi_0 + \Omega_e \Delta t) - \cos(\varphi_0)]$$

$$y_e(t + \Delta t) = y_e(t) - r_L [\sin(\varphi_0 + \Omega_e \Delta t) - \sin(\varphi_0)]$$

$$\varphi_0 = \tan^{-1}(v_{e,x}/v_{e,y}) \quad v_{e,\perp}^2 = v_{e,x}^2 + v_{e,y}^2 \quad \Omega_L = |eB_0|/m_e \quad r_L = V_{e,\perp}/\Omega_L$$



Symplectic kick for 2-body system

$$M_C(\Delta t): \quad \vec{x}_{ion} = \text{constant} \quad \vec{x}_e = \text{constant}$$
$$\Delta \vec{p}_{ion} = \frac{\alpha(\vec{x}_e - \vec{x}_{ion})\Delta t}{b^3(\vec{x}_{ion}, \vec{x}_e)} \quad \Delta \vec{p}_e = \frac{\alpha(\vec{x}_{ion} - \vec{x}_e)\Delta t}{b^3(\vec{x}_{ion}, \vec{x}_e)}$$
$$\alpha = \frac{Ze^2}{4\pi\epsilon_0} \quad b(\vec{x}_{ion}, \vec{x}_e) = \left[(x_{ion} - x_e)^2 + (y_{ion} - y_e)^2 + (z_{ion} - z_e)^2 \right]^{1/2}$$

These 2nd-order equation of motion are simple and robust.

They can be made 4th-order via standard Yoshida algorithm.

However, they require resolution of the gyroperiod and, hence, are slow:

$$\Delta t_{\max} \approx \frac{1}{8} \frac{2\pi}{\Omega_e}$$



Transform to Action-Angle variables

We follow Lichtenberg and Lieberman, *Regular & Chaotic Dynamics* (1992).

We use their canonical generating function of the 2nd kind:

$$F(x_e, y_e, \varphi, y_{gc}) = m_e \Omega_e \left[\frac{1}{2} (y_e - y_{gc})^2 \cot(\varphi) - x_e y_{gc} \right]$$

which yield the following Hamiltonian:

$$H(\vec{x}_{ion}, \vec{p}_{ion}, \varphi, y_{gc}, z_e, p_\varphi, p_{gc}, p_{ez}) = H_0(\vec{p}_{ion}, p_\varphi, p_{ez}) + H_C(\vec{x}_{ion}, \varphi, y_{gc}, z_e, p_\varphi, p_{gc})$$

$$H_0(\vec{p}_{ion}, p_\varphi, p_{e,z}) = \frac{1}{2m_{ion}} \vec{p}_{ion} \cdot \vec{p}_{ion} + \Omega_e p_\varphi + \frac{1}{2m_e} p_{e,z}^2$$

$$H_C(\vec{x}_{ion}, \varphi, y_{gc}, z_e, p_\varphi, p_{gc}, p_{ez}) = \frac{-Ze^2/4\pi\epsilon_0}{\sqrt{\left(x_{ion} - x_{gc}/m_e\Omega_e\right)^2 + \left(y_{ion} - y_{gc}\right)^2 + \left(z_{ion} - z_e\right)^2 + r_L^2 + \dots}} \\ \dots + 2\left(x_{ion} - x_{gc}\right)r_L \cos(\varphi) + 2\left(y_{ion} - y_{gc}\right)r_L \sin(\varphi)$$

$$x_{gc} = p_{gc}/m_e\Omega_e$$

$$r_L = \left(2p_\varphi/m_e\Omega_e\right)^{1/2}$$

Zero'th-order dynamics is now very simple, but H_C is problematic...



Transform to next-order Action-Angle variables

We follow Lichtenberg and Lieberman, *Regular & Chaotic Dynamics* (1992).

We use standard secular perturbation theory, requiring two approximations:

1) H_C is a perturbation. This requires $E_{\text{kinetic}} \gg E_{\text{potential}}$.

2) $r_L \ll \left[(x_{\text{ion}} - x_{\text{gc}})^2 + (y_{\text{ion}} - y_{\text{gc}})^2 + (z_{\text{ion}} - z_e)^2 + r_L^2 \right]^{1/2}$

This is approximately satisfied for relevant trajectories and fails gracefully.

The result is to remove the fast ϕ -dependence from the Hamiltonian:

$$H(\vec{x}_{\text{ion}}, y_{\text{gc}}, z_e, \vec{p}_{\text{ion}}, p_{\text{gc}}, p_{e,z}, J) = H_0(\vec{p}_{\text{ion}}, p_{e,z}, J) + H_C(\vec{x}_{\text{ion}}, y_{\text{gc}}, z_e, p_{\text{gc}}, J)$$

$$J = p_\phi + \frac{Ze^2}{4\pi\epsilon_0} \frac{r_L}{\Omega_e} \frac{(x_{\text{ion}} - x_{\text{gc}})\cos(\phi) + (y_{\text{ion}} - y_{\text{gc}})\sin(\phi)}{\left((x_{\text{ion}} - x_{\text{gc}})^2 + (y_{\text{ion}} - y_{\text{gc}})^2 + (z_{\text{ion}} - z_e)^2 + r_L^2 \right)^{3/2}}$$

$$H_0(\vec{p}_{\text{ion}}, J, p_{e,z}) = \frac{1}{2m_{\text{ion}}} \vec{p}_{\text{ion}} \cdot \vec{p}_{\text{ion}} + \Omega_e J + \frac{1}{2m_e} p_{e,z}^2$$

$$H_C(\vec{x}_{\text{ion}}, y_{\text{gc}}, z_e, J, p_{\text{gc}}) = \frac{-Ze^2}{4\pi\epsilon_0} \left/ \left[\left(x_{\text{ion}} - \frac{p_{\text{gc}}}{m_e \Omega_e} \right)^2 + (y_{\text{ion}} - y_{\text{gc}})^2 + (z_{\text{ion}} - z_e)^2 + \frac{2}{m_e \Omega_e} J \right]^{1/2} \right.$$



Symplectic maps for averaged 2-body system

Equations of motion are still in the standard drift-kick symplectic form:

$$M(\Delta t) = M_0(\Delta t/2)M_C(\Delta t)M_0(\Delta t/2)$$

$$M_0(\Delta t): \quad \vec{p}_{ion} = \text{constant} \quad J = \text{constant} \quad p_{gc} = \text{constant}$$

$$p_{ez} = \text{constant} \quad \theta \text{ is ignored} \quad y_{gc} = \text{constant}$$

$$\vec{x}_{ion}(t + \Delta t) = \vec{x}_{ion}(t) + \frac{\vec{p}_{ion}(t)}{m_{ion}} \Delta t \quad z_e(t + \Delta t) = z_e(t) + \frac{p_{e,z}(t)}{m_e} \Delta t$$

Much larger time steps are now possible:

$$\Delta t_{\max} \approx \frac{1}{8} \frac{|z_{ion} - z_e|}{|v_{ion,z} - v_{e,z}|}$$



Symplectic kick for averaged 2-body system

$$M_C(\Delta t):$$

$$\vec{x}_{ion} = \text{constant}$$

$$\Delta p_{ion,x} = \alpha \left(p_{gc} / m_e \Omega_e - z_{ion} \right) \Delta t / b^3 \left(\vec{x}_{ion}, y_{gc}, z_e, J, p_{gc} \right)$$

$$J = \text{constant}$$

θ is ignored

$$\Delta p_{ion,y} = \alpha \left(y_{gc} - z_{ion} \right) \Delta t / b^3 \left(\vec{x}_{ion}, y_{gc}, z_e, J, p_{gc} \right)$$

$$z_e = \text{constant}$$

$$\Delta p_{ion,z} = \alpha \left(z_e - z_{ion} \right) \Delta t / b^3 \left(\vec{x}_{ion}, y_{gc}, z_e, J, p_{gc} \right)$$

$$\Delta p_{ez} = -\Delta p_{ion,z}$$

$$\alpha = \frac{Ze^2}{4\pi\epsilon_0}$$

$$b(\vec{x}_{ion}, y_{gc}, z_e, J, p_{gc}) = \left[\left(x_{ion} - \frac{p_{gc}}{m_e \Omega_e} \right)^2 + (y_{ion} - y_{gc})^2 + (z_{ion} - z_e)^2 + \frac{2}{m_e \Omega_e} J \right]^{1/2}$$



Time-explicit vs Averaged – approx. agreement





Figure 1: Relative error in the value of ΔP_{ion} computed using four-turn averaging (blue circles) and the Magnus expansion (red squares).



Time-explicit vs Averaged – approx. agreement

$\log_{10}(\text{relative error } \Delta P_{\text{averaging}})$



This graphic shows the relative error made by our averaging computation of the ion momentum kick ΔP . The average is taken over one full gyrotron period. The scale is logarithmic, so the largest errors here are about 1 %.



Magnus expansion yields analytic result

We follow Alex Dragt, *Lie Methods for Nonlinear Dynamics with Applications to Accelerator Physics* (Version of 22 June 2016), p. 861:
<http://www.physics.umd.edu/dsat/dsatliemethods.html>

The result is an analytic calculation of the ion momentum change!

$$M(t) = M_I(t)M_0(t)$$

$$M_I(t) = \exp\left[-: \int_0^t d\sigma H_I(\sigma) : \right] \quad H_I(t) = M_0(t)H_C$$

$$M_I(T) \approx I - : \int_0^T d\sigma H_I(\sigma) :$$

We can evaluate this approximate expression analytically.
It is valid, when the Coulomb interaction is a perturbation.



Analytic calculation of $\Delta \mathbf{p}_{ion}$ (1)

$$C_1 = \left(x_{ion} - \frac{P_{gc}}{m_e \Omega_e} \right)^2 + (y_{ion} - y_{gc})^2 + (z_{ion} - z_e)^2 + \frac{2}{m_e \Omega_e} J \quad (14a)$$

$$C_2 = 2(x_{ion} - x_{gc})v_{ion,x} + 2(y_{ion} - y_{gc})v_{ion,y} + 2(z_{ion} - z_e)(v_{ion,z} - v_{ez}) \quad (14b)$$

$$C_3 = v_{ion,x}^2 + v_{ion,y}^2 + (v_{ion,x} - v_{ez})^2 \quad (14c)$$

$$b = [C_1 + C_2 T + C_3 T^2]^{1/2} \quad \Delta = 4C_1 C_3 - C_2^2 \quad (14d)$$

$$D_1 = \left[\frac{2C_3 T + C_2}{b} - \frac{C_2}{\sqrt{C_1}} \right] \quad (14e)$$

$$D_2 = \left[\frac{2C_1 + C_2 T}{b} - 2\sqrt{C_1} \right] \quad (14f)$$



Analytic calculation of $\Delta \mathbf{p}_{ion}$ (2)

$$\Delta p_{ion,x} = \frac{-2\alpha}{\Delta} \left[(x_{ion} - p_{gc}/m_e \Omega_e) D_1 - (p_{ion,x}/m_{ion}) D_2 \right] \quad (15a)$$

$$\Delta p_{ion,y} = \frac{-2\alpha}{\Delta} \left[(y_{ion} - y_{gc}) D_1 - (p_{ion,y}/m_{ion}) D_2 \right] \quad (15b)$$

$$\Delta p_{ion,z} = \frac{-2\alpha}{\Delta} \left[(z_{ion} - z_e) D_1 - \left(\frac{p_{ion,z}}{m_{ion}} - \frac{p_{ez}}{m_e} \right) D_2 \right] \quad (15c)$$

$$\Delta p_{gc} = -\Delta p_{ion,x} \quad \Delta y_{gc} = -\Delta p_{ion,y}/m_e \Omega_e \quad (15d)$$



Time-explicit vs Magnus – approx. agreement





Figure 1: Relative error in the value of ΔP_{ion} computed using four-turn averaging (blue circles) and the Magnus expansion (red squares).



Time-explicit vs Magnus – approx. agreement


$\log_{10}(\text{relative error } \Delta P_{\text{Magnus}})$



This graphic shows the relative error made by our Magnus computation of ΔP .




Integrate to obtain Friction force



$$F = -\frac{n_e m_e}{T} \int \int \int_{\mathbb{R}^3} d^3 v \int \int \int_V dr dz d\varphi \Delta v(T, r, \varphi, z, v) r p(v)$$



Integrate to obtain Friction force



$$F = -\frac{n_e m_e}{T} \int \int \int_{\mathbb{R}^3} d^3 v \int \int \int_V dr dz d\varphi \Delta v(T, r, \varphi, z, v) r p(v)$$



Include other effects in Magnus Expansion

$$H(\vec{x}_{ion}, \vec{p}_{ion}, \vec{x}_e, \vec{p}_e) = H_0(\vec{p}_{ion}, y_e, \vec{p}_e) + H_C(\vec{x}_{ion}, \vec{x}_e) \\ + H_{space-charge}(\vec{x}_{ion}, \vec{x}_e) + H_{solenoid-field-errors}(??)$$

- Quantitative treatment of space charge & field errors?
 - *space charge should work*
 - *field errors are more challenging*
- Requires generalization of Magnus expansion
 - *we are optimistic this can be done*

