

Evolution of
Pseudo-PDFs
and
Quasi-PDFs

Evolution of Pseudo-PDFs and Quasi-PDFs


A.V. Radyushkin (ODU/Jlab)

- Parton Densities
- Transverse Momentum Cut-off
- Pseudo-distributions
- qPDFs
- qPDF/TMD relation
- Hard tail
- Gauge link
- Renormalization
- Reduced pseudo-ITD

- Evolution in lattice data
- Data
- Building \overline{MS} ITD

Summary

QCD Evolution 2018
May 22, 2018



Parton Densities


Evolution of Pseudo-PDFs and Quasi-PDFs

Parton Densities
 Transverse Momentum Cut-off
 Pseudo-distributions
 qPDFs
 qPDF/TMD relation
 Hard tail
 Gauge link
 Renormalization
 Reduced pseudo-ITD


Evolution in lattice data
 Data
 Building \overline{MS} ITD

Summary

- May 11, 1918 – R.P. Feynman's birthday
- Feynman diagrams, propagator, path integrals, parton model ...
- Original Feynman approach to PDFs $f(x)$: infinite momentum $P_3 \rightarrow \infty$ limit of $k_3 = xP_3$ momentum distributions (\sim quasi-PDFs $Q(x, P_3)$)
- $f(x)$ were treated as k_{\perp} -integrals of more detailed $f(x, k_{\perp})$ distributions
- From the start it was understood that $Q(x, P_3 \rightarrow \infty) \rightarrow f(x)$ limit exists only if $f(x, k_{\perp})$ rapidly decreases with k_{\perp}
- “Transverse momentum cut-off”, $\langle k_{\perp}^2 \rangle \sim 1/R_{\text{hadr}}^2$
- Question 1: why $Q(x, P_3)$ differs from $f(x)$?
- Question 2: how does $Q(x, P_3)$ convert into $f(x)$ when $P_3 \rightarrow \infty$?
- Qualitative answer: yP_3 comes from two sources:
 from the motion of the hadron as a whole (xP_3) and
 from Fermi motion of quarks inside the hadron ($(y - x)P_3 \sim 1/R_{\text{hadr}}$)



- $(y - x)P_3 \sim 1/R_{\text{hadr}}$ part has the same origin as transverse momentum
- \Rightarrow One should be able to relate quasi-PDFs to TMDs



Pseudo-distributions and PDFs

Evolution of Pseudo-PDFs and Quasi-PDFs

Parton Densities

Transverse Momentum Cut-off

Pseudo-distributions

qPDFs

qPDF/TMD relation

Hard tail

Gauge link

Renormalization

Reduced


pseudo-ITD

Evolution in lattice data

Data

Building $\overline{\text{MS}}$ ITD

Summary



- Basic matrix element (ignoring spin)

$$\langle p | \phi(0) \phi(z) | p \rangle = \mathcal{M}(-pz, -z^2)$$

- Lorentz invariance: \mathcal{M} depends on z through $(pz) \equiv -\nu$ and z^2


- loffe time ν : $\mathcal{M}(\nu, -z^2) =$ **loffe-time pseudo-distribution** (pseudo-ITD)
- Pseudo** \equiv off the light cone
- For any Feynman diagram, for **arbitrary** z^2 and **arbitrary** p^2

$$\mathcal{M}(\nu, -z^2) = \int_{-1}^1 dx e^{ix\nu} \mathcal{P}(x, -z^2)$$

- Limits $-1 \leq x \leq 1$, negative x correspond to anti-particles
- On the light cone**: usual ITD and usual PDF $\mathcal{P}(x, 0) = f(x)$
- If $z^2 \rightarrow 0$ limit is singular, regularization (like $\overline{\text{MS}}$) is needed, $f(x) \rightarrow f(x, \mu^2)$ and we have $\overline{\text{MS}}$ ITD

$$\mathcal{M}(\nu, 0)|_{\mu^2} \equiv \mathcal{I}(\nu, \mu^2) = \int_{-1}^1 dx e^{ix\nu} f(x, \mu^2)$$

- Pseudo-PDF** $\mathcal{P}(x, -z^2)$: Fourier transform of pseudo-ITD with respect to ν for fixed z^2



Quasi-PDFs

Evolution of Pseudo-PDFs and Quasi-PDFs


Parton Densities

Transverse Momentum Cut-off
 Pseudo-distributions
 qPDFs
 qPDF/TMD relation
 Hard tail
 Gauge link
 Renormalization
 Reduced pseudo-ITD

Evolution in lattice data

Data
 Building \overline{MS} ITD

Summary



- Basic matrix element (ignoring spin)

$$\langle p | \phi(0) \phi(z) | p \rangle = \mathcal{M}(-(pz), -z^2)$$

- Lorentz invariance: \mathcal{M} depends on z through $(pz) \equiv -\nu$ and z^2


- Take $z = (0, 0, 0, z_3)$, then $-(pz) \equiv \nu = Pz_3$ and $-z^2 = z_3^2$
- Introduce **quasi-PDF** (Ji,2013)

$$Q(y, P) = \frac{P}{2\pi} \int_{-\infty}^{\infty} dz_3 e^{-iyPz_3} \mathcal{M}(Pz_3, z_3^2)$$

- Write $\mathcal{M}(Pz_3, z_3^2)$ through pseudo-PDF

$$Q(y, P) = \frac{P}{2\pi} \int_{-1}^1 dx \int_{-\infty}^{\infty} dz_3 e^{-i(y-x)Pz_3} \mathcal{P}(x, z_3^2)$$

- Quasi-PDFs $Q(y, P)$ are defined for all $-\infty < y < \infty$
- If $\mathcal{P}(x, z_3^2) = f(x)$, then $Q(y, P) = f(y)$



qPDF/TMD relation

Evolution of Pseudo-PDFs and Quasi-PDFs

Parton Densities

Transverse Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation

Hard tail
Gauge link
Renormalization
Reduced pseudo-ITD

Evolution in lattice data

Data
Building \overline{MS} ITD

Summary

- Using $z_3 = \nu/P$ as integration variable

$$Q(y, P) = \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} e^{-iy\nu} \mathcal{M}(\nu, \nu^2/P^2)$$

- Take $z = (z_+ = 0, z_-, z_1, z_2)$. Then $\nu = -p^+ z^-$ and $-z^2 = z_1^2 + z_2^2$
- Introduce **TMD** $\mathcal{F}(x, k_1^2 + k_2^2)$:


$$\mathcal{M}(\nu, z_1^2 + z_2^2) = \int_{-1}^1 dx e^{ix\nu} \int_{-\infty}^{\infty} dk_1 dk_2 e^{i(k_1 z_1 + k_2 z_2)} \mathcal{F}(x, k_1^2 + k_2^2)$$

- Combining with Eq. for $Q(y, P)$ in terms of $\mathcal{M}(\nu, \nu^2/P^2)$

$$Q(y, P) = P \int_{-1}^1 dx \int_{-\infty}^{\infty} dk_1 \mathcal{F}(x, k_1^2 + (y-x)^2 P^2)$$

- Is it possible to study the approach of $Q(y, P)$ to $f(y)$?
- Try factorized model $\mathcal{F}^{\text{fact}}(x, k_{\perp}^2) = f(x)K(k_{\perp}^2)$
- Popular idea: Gaussian dependence $K_G(k_{\perp}^2) = e^{-k_{\perp}^2/\Lambda^2}/\pi\Lambda^2$

$$Q_G^{\text{fact}}(y, P) = \frac{P}{\Lambda\sqrt{\pi}} \int_{-1}^1 dx f(x) e^{-(y-x)P^2/\Lambda^2}$$



Evolution of
Pseudo-PDFs
and
Quasi-PDFs

Parton
Densities


Transverse
Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced
pseudo-ITD

Evolution in
lattice data


Data
Building \overline{MS} ITD

Summary

- Take PDF $f(x) = u_v(x) - d_v(x) = \frac{315}{32} \sqrt{x}(1-x)^3 \theta(0 \leq x \leq 1)$ obtained by pseudo-PDF method (Orginos et al. 2017)



- Curves for $P/\Lambda = 0.75, 1.5, 2.25$ are close to qPDFs obtained by Lin et al (2016), upper momentum $P = 1.3$ GeV, effective $\Lambda \approx 600$ MeV
- Need $P \sim 4.5 \Lambda \approx 2.7$ GeV to get reasonably close to input PDF
- Note a lot of dirt for negative y , even for $P/\Lambda = 4.5$



Renormalizable theories and hard term

7/15

$$Q(y, P) = \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} e^{-iy\nu} \mathcal{M}(\nu, \nu^2/P^2)$$

- In QCD $\mathcal{M}(\nu, z_3^2)$ has logarithmic singularity in z_3^2 . At one loop,

$$\mathcal{M}^{\text{hard}}(\nu, z_3^2) = -\frac{\alpha_s}{2\pi} C_F \ln(z_3^2) \int_0^1 du B(u) \mathcal{M}^{\text{soft}}(u\nu, 0)$$

- Generates perturbative evolution. Altarelli-Parisi (AP) evolution kernel

$$B(u) = \left[\frac{1+u^2}{1-u} \right]_+$$

- The function $\mathcal{M}(\nu, \nu^2/P^2)$ that generates the quasi-PDF gets

$$\mathcal{M}^{\text{hard}}(\nu, \nu^2/P^2) = -\frac{\alpha_s}{2\pi} C_F \ln(\nu^2/P^2) \int_0^1 du B(u) \int_{-1}^1 dx e^{-iux\nu} f^{\text{soft}}(x)$$

- Hard part of the quasi-PDF $Q(y, P)$ has a $\ln P^2$ term

$$Q^{\text{hard}}(y, P) = \ln(P^2) \Delta(y) + \dots$$

- It is nonzero in the $-1 \leq y \leq 1$ region only

$$\Delta(y) = \frac{\alpha_s}{2\pi} C_F \int_0^1 \frac{du}{u} B(u) f^{\text{soft}}(y/u)$$

Evolution of
Pseudo-PDFs
and
Quasi-PDFs

Parton
Densities

Transverse
Momentum Cut-off
Pseudo-distributions

qPDFs
qPDF/TMD relation


Hard tail

Gauge link
Renormalization
Reduced
pseudo-ITD

Evolution in
lattice data

Data
Building $\overline{\text{MS}}$ ITD

Summary



Gauge link complications

Evolution of Pseudo-PDFs and Quasi-PDFs


Parton Densities

Transverse Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced pseudo-ITD

Evolution in lattice data

Data
Building \overline{MS} ITD

Summary




- In QCD, there is one more source of the z^2 -dependence of pseudo-ITD: gauge link $\hat{E}(0, z; A)$
- It has specific ultraviolet divergences
- Use Polyakov regularization $1/\delta z^2 \rightarrow 1/(\delta z^2 - a^2)$ for gluon propagator in coordinate space

- Effect of the UV cut-off a is similar to that of the lattice spacing
- At one loop, link-related UV singular terms have the structure

$$\Gamma_{UV}(z_3, a) \sim -\frac{\alpha_s}{2\pi} C_F \left[2 \frac{|z_3|}{a} \tan^{-1} \left(\frac{|z_3|}{a} \right) - 2 \ln \left(1 + \frac{z_3^2}{a^2} \right) \right]$$

- For fixed a , these terms vanish when $z_3 \rightarrow 0$
- No violation of quark number conservation
- Because of UV singularities, there is large activity to **renormalize out** link-related factor



Renormalize or exterminate?

Evolution of Pseudo-PDFs and Quasi-PDFs

Parton Densities

Transverse Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link




Renormalization

Reduced pseudo-ITD

Evolution in lattice data

Data
Building \overline{MS} ITD

Summary




- Structure of factorization for DIS in Feynman gauge
- Sum of gluon insertions gives $\langle P | \bar{\psi}(0) \gamma S^c(z) \gamma \hat{E}(0, z; A) \psi(z) | P \rangle$ + higher twists
- But: quark self-energy diagram is not factorized as $S^c(z) \times \langle AA \rangle$
- Operator $\bar{\psi}(0) \hat{E}(0, z; A) \psi(z)$ should be accompanied by “no AA contractions”
- Link self-energy diagrams and UV-singular parts of vertex diagrams should be excluded **together with** associated z_3^2 -dependence
- It is **not sufficient** just to subtract UV divergences

- Easy way out: consider reduced pseudo-ITD

$$\mathfrak{M}(\nu, z_3^2) \equiv \frac{\mathcal{M}(\nu, z_3^2)}{\mathcal{M}(0, z_3^2)}$$

- $\mathfrak{M}(\nu, z_3^2)$ has finite $a \rightarrow 0$ limit



Reduced Ioffe-time pseudo-distribution

Evolution of Pseudo-PDFs and Quasi-PDFs

Parton Densities

Transverse Momentum Cut-off Pseudo-distributions

qPDFs qPDF/TMD relation

Hard tail Gauge link Renormalization

Reduced pseudo-ITD

Evolution in lattice data

Data Building $\overline{\text{MS}}$ ITD

Summary


- Reduced pseudo-ITD $\mathfrak{M}(\nu, z_3^2)$ is a physical observable (like, say, DIS structure functions)
- No need to specify renormalization scheme, scale, etc. for link-related terms
- Pseudo-ITD $\mathfrak{M}(\nu, z_3^2)$ is finite for fixed z_3 , but is singular in $z_3 \rightarrow 0$ limit
- $\ln z_3^2$ terms reflect perturbative evolution
- For light-cone PDF, one takes $z^2 = 0$ and uses some scheme for **resulting** UV divergence, say, $\overline{\text{MS}}$
- Ioffe-time distribution $\mathcal{I}(\nu, \mu^2)$ is UV scheme and scale dependent

$$\mathcal{I}(\nu, \mu^2) = \int_{-1}^1 dx e^{ix\nu} f(x, \mu^2)$$

- One-loop relation between $\overline{\text{MS}}$ ITD and reduced pseudo-ITD

$$\begin{aligned} \mathcal{I}(\nu, \mu^2) &= \mathfrak{M}(\nu, z_3^2) + \frac{\alpha_s}{2\pi} C_F \int_0^1 dw \mathfrak{M}(w\nu, z_3^2) \\ &\times \left\{ \frac{1+w^2}{1-w} \left[\ln \left(z_3^2 \mu^2 \frac{e^{2\gamma_E}}{4} \right) + 1 \right] + \left[4 \frac{\ln(1-w)}{1-w} - 2(1-w) \right] \right\}_+ \end{aligned}$$

Evolution in lattice data



Evolution of
Pseudo-PDFs
and
Quasi-PDFs

Parton
Densities

Transverse
Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced
pseudo-ITD

Evolution in
lattice data


Data
Building \overline{MS} ITD

Summary

- Exploratory lattice study of reduced pseudo-ITD $\mathfrak{M}(\nu, z_3^2)$ for the valence $u_v - d_v$ parton distribution in the nucleon [Orginos et al. 2017]
- Real part corresponds to the cosine Fourier transform of $q_v(x) = u_v(x) - d_v(x)$

$$\Re(\nu) \equiv \text{Re } \mathfrak{M}(\nu) = \int_0^1 dx \cos(\nu x) q_v(x)$$

- When plotted as function of ν , data both for real and imaginary parts lie close to respective universal curves




- Overall curve corresponds to the function

$$f(x) = \frac{315}{32} \sqrt{x}(1-x)^3$$

- Obtained by forming cosine Fourier transforms of $x^a(1-x)^b$ -type functions and fitting a, b
- Shape is dominated by points with smaller values of $\text{Re } \mathfrak{M}(\nu, z_3^2)$

- Data for $\mathfrak{M}(\nu, z_3^2) \equiv \mathcal{M}(\nu, z_3^2)/\mathcal{M}(0, z_3^2)$ show no polynomial z_3 -dependence for large z_3 though z_3^2/a^2 changes from 1 to ~ 200
- Apparently no higher-twist terms in the reduced pseudo-ITD
- Meaning: $\mathcal{M}(\nu, z_3^2)$ factorizes as $M(\nu)\mathcal{M}(0, z_3^2)$ for large z_3



Lattice data for small and large z_3

Evolution of Pseudo-PDFs and Quasi-PDFs



Parton Densities

- Transverse Momentum Cut-off
- Pseudo-distributions
- qPDFs
- qPDF/TMD relation
- Hard tail
- Gauge link
- Renormalization
- Reduced pseudo-ITD


Evolution in lattice data

- Data
- Building \overline{MS} ITD

Summary



- Points corresponding to $7a \leq z_3 \leq 13a$ values
- Some scatter for points with $\nu \gtrsim 10$
- Otherwise, practically all the points lie on the universal curve based on $f(x)$.
- No z_3 -evolution visible in large- z_3 data
- Points in $a \leq z_3 \leq 6a$ region
- All points lie higher than the curve based on the $z_3 \geq 7a$ data
- Perturbative evolution increases real part of the pseudo-ITD when z_3 decreases
- Conjecture that the observed higher values of $\text{Re}\mathfrak{M}$ for smaller- z_3 points may be a consequence of evolution



Building \overline{MS} ITD

Evolution of Pseudo-PDFs and Quasi-PDFs


Parton Densities

Transverse Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced pseudo-ITD

Evolution in lattice data

Data
Building \overline{MS} ITD


Summary



- z_3 -dependence of the lattice points for “magic” Ioffe-time value $\nu \equiv z_3 P_3 = 12\pi/16 = 3\pi/4$
- Eye-ball fit line has “Perturbative” $\ln(1/z_3^2)$ behavior for small z_3 , and rapidly tends to a constant for $z_3 \gtrsim 6a$
- $\Re(\nu, z_3^2)$ decreases when z_3 increases
- Starts to visibly deviate from a pure logarithmic $\ln z_3^2$ pattern for $z_3 \gtrsim 5a$
- This sets the boundary $z_3 \leq 4a$ on the “logarithmic region”
- \overline{MS} ITD in terms of reduced pseudo-ITD

$$\mathcal{I}(\nu, \mu^2) = \mathfrak{M}(\nu, z_3^2) + \frac{\alpha_s}{2\pi} C_F \int_0^1 dw \mathfrak{M}(w\nu, z_3^2) \times \left\{ \frac{1+w^2}{1-w} \left[\ln \left(z_3^2 \mu^2 \frac{e^{2\gamma_E}}{4} \right) + 1 \right] + \left[4 \frac{\ln(1-w)}{1-w} - 2(1-w) \right] \right\}_+$$

- $\mathcal{I}(\nu, \mu^2)$ should not depend on z_3
- This happens only if, for some α_s , the $\ln z_3^2$ -dependence of the 1-loop term cancels actual z_3^2 -dependence of the data, visible as scatter in the data



Numerical results for $\overline{\text{MS}}$ ITD

14/15

Evolution of Pseudo-PDFs and Quasi-PDFs



Parton Densities

Transverse Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced pseudo-ITD


Evolution in lattice data

Data
Building $\overline{\text{MS}}$ ITD

Summary



- We choose $\mu = 1/a$ which, at lattice spacing of 0.093 fm is ≈ 2.15 GeV
- Using $\alpha_s/\pi = 0.1$ and $z_3 \leq 4a$ data, we generate the points for $\mathcal{I}_R(\nu, (1/a)^2)$
- Upper curve corresponds to the ITD of the CJ15 global fit PDF for $\mu = 2.15$ GeV
- Evolved points are close to some universal curve with a rather small scatter
- The curve itself corresponds to the cosine transform of a normalized $\sim x^a(1-x)^b$ distribution with $a = 0.35$ and $b = 3$
- $\sim x^{0.35}(1-x)^3$ PDF compared to CJ15 and MMHT global fits for $\mu = 2.15$ GeV
- Unable to reproduce $\sim x^{-0.5}$ Regge behavior
- Possible reasons: large pion mass, quenched approximation



Summary

15/15

Evolution of
Pseudo-PDFs
and
Quasi-PDFs

Parton
Densities

Transverse
Momentum Cut-off
Pseudo-distributions
qPDFs
qPDF/TMD relation
Hard tail
Gauge link
Renormalization
Reduced
pseudo-ITD

Evolution in
lattice data

Data
Building \overline{MS} ITD

Summary

- Analyzed nonperturbative structure of quasi-PDFs $Q(y, P)$ using their relation to pseudo-ITDs and TMDs
- Studied nonperturbative evolution of quasi-PDFs $Q(y, P)$ with P using factorized models for TMDs
- Analyzed perturbative structure of quasi-PDFs using their relation to pseudo-ITDs and TMDs
- Argued that link-related terms should be “exterminated”
- Proposed to use reduced pseudo-ITD
- Studied evolution of exploratory lattice data for reduced pseudo-ITD