




Fragmentation Function measurements in Belle


**FF2019 workshop,
March 14, 2019, Duke University**

Ralf Seidl (RIKEN)


SIDIS



pp collisions



e⁺e⁻ Annihilation



Access to FFs

• SIDIS:

$$\sigma^h(x, z, Q^2, P_{h\perp}) \propto \sum_q e_q^2 q(x, p_t, Q^2) D_{1,q}^h(z, k_t, Q^2)$$

- Relies on unpol PDFs
- Parton momentum known at LO
- Flavor structure directly accessible
- Transverse momenta convoluted between FF and PDF

• pp:

$$\sigma^h(P_T) \propto \int_{x_1, x_2, z} \sum_{a, a' \in q, g} f_a(x_1) \otimes f_{a'}(x_2) \otimes \sigma_{aa'} \otimes D_{1,q}^h(z)$$

- Relies on unpol PDFs
- leading access to gluon FF
- Parton momenta not directly known

• e⁺e⁻:

$$\sigma^h(z, Q^2, k_t) \propto \sum_q e_q^2 (D_{1,q}^h(z, k_t, Q^2) + D_{1,\bar{q}}^h(z, k_t, Q^2))$$

- No PDFs necessary
- Clean initial state, parton momentum known at LO
- Flavor structure not directly accessible

EIC without B factory input?

Very unlikely


- Very limited helicity analysis possible (based on Kretzer or KKP)
- Only model dependent Tensor charge extraction
- Sivers and all TMDs just with naïve Gaussian dependence (no x or z dependence)

Transverse momentum dependence

Aka un-integrated PDFs and FFs

K_T Dependence of FFs in e^+e^-

- Gain also sensitivity into transverse momentum generated in fragmentation
- Two ways to obtain transverse momentum dependence
 - Traditional 2-hadron FF
 - use transverse momentum between two hadrons (in opposite hemispheres)
 - Usual convolution of two transverse momenta
 - Single-hadron FF wrt to **Thrust** or jet axis
 - No convolution
 - Need correction for $q\bar{q}$ axis




Thrust definition

- Event shape variable thrust is defined as:

$$T \equiv \frac{\sum_h |\mathbf{P}_h \cdot \hat{\mathbf{n}}|}{\sum_h |\mathbf{P}_h|}$$

- All final-state particles are included in the sum
- A **two-jet-like** event has a high thrust value
- A completely **spherical** event has a thrust value of 0.5

- Thrust axis \mathbf{n} also defines the hemispheres




Correction chain

Correction	Method	Systematics
PID mis-id	PID matrices (5x5 for $\cos \theta_{\text{lab}}$ and p_{lab})	MC sampling of inverted matrix element uncertainties, variation of PID correction method
Momentum smearing	MC based smearing matrices (2160x2160), SVD unfold	SVD unfolding vs analytically inverted matrix, reorganized binning, MC statistics
Non-qqbar BG removal	eeuu, eess, eecc, tau MC subtraction	Variation of size, MC statistics
Acceptance I (cut efficiency)	In barrel reconstructed vs udsc generated in barrel	MC statistics
Acceptance II	udsc Gen MC barrel to 4π	MC statistics, variation in tunes
Weak decay removal (optional)	udcs check evt record for weak decays	Compare to other Pythia settings
ISR	ISR on vs ISR off in Pythia	Variations in tunes

6 thrust bins [0.5,0.7,0.8,0.85,0.9,0.95,1.0] x 18 z bins x 20 kt bins


PID correction

Using Martin Leitgab's 5x5 PID matrices in fine $17 \times 9 P_{\text{lab}} \times \cos\theta_{\text{lab}}$ binning




Smearing

- Reduced smearing matrices from 2160 x 2160 to filled (ie kinematically reachable bins)
- Using SVDUnfold Method in Root




Non-qqbar removal:

Remove all two-photon and tau events from yields, contributions generally up to several %, slightly higher for kaons rand low thrust




Stacked, relative contributions



Acceptance correction

ACCI: Reconstruction and efficiency correction in Barrel acceptance

ACCII: Barrel to 4π correction



$$\frac{d^2\sigma}{dzdk_{T,h}}$$

3/14/2019


k_{Th}

R.Seidl: e+e- Fragmentation

12

Weak correction(optional)

Traced in gen MC hadrons back to mothers with non ud content \rightarrow if not vetoed (K^* , ssbar, ccbar resonances, some hyperons and excited states) \rightarrow **Weak**




3/14/2019

both cross sections provided in paper

13


ISR correction

All different tunes very similar
except old Belle tune
→ assigned as systematics
-high k_T drop of ratio due to ISR
boost





Overall systematic uncertainties

Systematic uncertainties dominated by acceptance correction (for different tunes), PID uncertainties and ISR correction



Cross sections various hadrons

Belle Preliminary



3/14/2019



k_T


RS. et. al. [arXiv:1902.01552](https://arxiv.org/abs/1902.01552)

R.Seidl: e+e- Fragmentation

16


Fits vs k_T^2

Fit exponential to smaller transverse momenta for
Gaussian k_T dependence and power law at higher k_T



Gaussian widths

first direct (no convolutions) measurement of z dependence of Gaussian widths




[arXiv:1902.01552](https://arxiv.org/abs/1902.01552)

3/14/2019

Current phenomenological models: no or linear z dependence only

Gaussian widths, thrust dependence


Gaussian widths get narrower with higher Thrust



[arXiv:1902.01552](https://arxiv.org/abs/1902.01552)

Gaussian widths comparison to MC

first direct (no convolutions) measurement of z dependence
of Gaussian widths



Single Λ polarization measurements



- Related to open question about Λ polarization in hadron collisions from 40 years ago!
- Fragmentation counterpart to the Sivers Function:

unpolarized parton fragments into transversely polarized baryon with transverse momentum wrt to parton direction

- Reconstruct Λ , its transverse momentum and polarization


YingHui Guan (Indiana/KEK):
[PRL 122 \(2019\), 042001](#)

$$D_{1,q}^{\perp h}(z, k_T, Q^2)$$




Transverse momentum dependence

- Different behavior for low and high- z :
- At low z small
- At intermediate z falling Polarization with k_t
- At high z increasing polarization with k_t




Opposite hemisphere pion correlation

- Interesting z_π and z_Λ dependence :
- At low z_Λ light quark fragmentation dominant, some charm in $\pi^- \rightarrow$ different signs
- At high z_Λ strange + charm fragmentation more relevant \rightarrow same signs



Ongoing work: two-hadron transverse momentum

- Analysis ongoing (Anselm/Charlotte)
 - Differential in z_1, z_2 and q_t for pion and kaon combinations
 - All the correction steps (PID, smearing, non- $q\bar{q}$, acceptance, ISR) similar to recent Belle FF analyses



Ongoing work: Collins multidimensional analysis and Kaon combinations

- Currently revisiting kaon combinations of the Collins asymmetries previously done by Francesca
- While doing so, try to perform a full multi-dimensional analysis:
 - Currently (for testing):
 - $6(z_1) \times 6(z_2) \times 4(k_{t1}) \times 4(k_{t2}) \times 1(\text{costheta}) \times 8(\text{phi})$
 - $6(z_1) \times 6(z_2) \times 10(q_t) \times 1(\text{costheta}) \times 8(\text{phi})$
 - Consider 5 k_t bins and several costheta bins after successful test (ongoing)
- Use most correction steps similar to recent analyses (PID, smearing, non-qqbar removal, acceptance?, ISR?)
- To simplify smearing unfold each z_1 - z_2 bin separately (z smearing almost nonexistent)

Summary

- Kt dependent cross sections and Gaussian widths extracted
 - Very clear z dependence of widths, not as assumed by phenomenologists
 - Pions and kaons similar, protons narrower (diquarks?)
- Lambda polarization paper finally published
- More Belle measurements ongoing (di-hadron kt, Collins)

Differences in Pythia/JetSet settings

Par	0	1	9	10	11	12	13	udscatlas	udschermes
	Pythia def.	belle	Atlas	Aleph	LEP/tev.	Hermes	gen Belle		
PARJ(1)	0.1			0.106	0.073	0.029			0.029
PARJ(2)	0.3			0.285	0.2	0.283			0.283
PARJ(3)	0.4			0.71	0.94	1.2			1.2
PARJ(4)	0.05			0.05	0.032				
PARJ(11)	0.5			0.55	0.31				
PARJ(12)	0.6			0.47	0.4				
PARJ(13)	0.75			0.65	0.54				
PARJ(14)	0.0	0.0	0.0	0.02	0.0	0.0	0.05	0.0	0.0
PARJ(15)	0.0	0.0	0.0	0.04	0.0	0.0	0.05	0.0	0.0
PARJ(16)	0.0		0.0	0.02	0.0	0.0	0.05	0.0	0.0
PARJ(17)	0.0	0.0	0.0	0.2	0.0	0.0	0.05	0.0	0.0
PARJ(19)	1			0.57					
PARJ(21)	0.36			0.37	0.325	0.400	0.28	0.28	0.400
PARJ(25)	1				0.63		0.27	0.27	
PARJ(26)	0.4			0.27	0.12		0	0	
PARJ(33)	0.8		0.8	0.8	0.8	0.3		0.8	0.8
PARJ(41)	0.3			0.4	0.5	1.94	0.32	0.32	1.94
PARJ(42)	0.58			0.796	0.6	0.544	0.62	0.62	0.544
PARJ(45)	0.5					1.05			1.05
PARJ(46)	1.						1.0	1.0	
PARJ(47)	1.				0.67				
PARJ(54)	-0.050	-0.040	-0.050	-0.04	-0.050	-0.050		-0.050	-0.050
PARJ(55)	-0.005	-0.004	-0.005	-0.0035	-0.005	-0.005		-0.005	-0.005
PARJ(81)	0.29			0.292	0.29		0.38	0.38	
PARJ(82)	1.0			1.57	1.65		0.5	0.5	
MSTJ(11)	4			3	5		4	4	
MSTJ(12)	2			3		1			1
MSTJ(26)	2	0	2	2	2	2	0	2	2
MSTJ(45)	5					4			4
MSTJ(107)	0	1	0	0	0	0	1	0	0

VM
suppression

P_x, P_y Gauss
width

Lund params

Λ_{QCD} and E
cutoff

Pythia/Jetset parameters

PARJ(1)	:	Diquark suppression relative to quark antiquark production
PARJ(2)	:	Strangeness suppression relative to u or d pair production
PARJ(3)	:	Extra suppression of strange diquarks relative to strange quark production
PARJ(4)	:	Axial (ud_1) vs scalar (ud_0) diquark suppression
PARJ(11)	:	Light meson with spin 1 probability
PARJ(12)	:	Strange meson with spin 1 probability
PARJ(13)	:	Charm meson with spin 1 probability
PARJ(14)	:	Spin 0 meson with $L = 1$ and $J = 1$ probability
PARJ(15)	:	Spin 1 meson with $L = 1$ and $J = 0$ probability
PARJ(16)	:	Spin 1 meson with $L = 1$ and $J = 1$ probability
PARJ(17)	:	Spin 1 meson with $L = 1$ and $J = 2$ probability
PARJ(19)	:	Extra baryon suppression relative to regular diquark suppression (if MSTJ(12) = 3)
PARJ(21)	:	Gaussian Width of p_x and p_y for primary hadrons
PARJ(25)	:	η production suppression factor
PARJ(26)	:	η' production suppression factor
PARJ(33)	:	Energy cutoff of fragmentation process
PARJ(41)	:	Lund a parameter: $(1 - z)^a$
PARJ(42)	:	Lund b parameter: $\exp(-bm_{\perp}^2/z)$
PARJ(45)	:	addition to a parameter for diquarks
PARJ(46)	:	modification of Lund fragmentation for heavy quarks with Bowler, charm, bottom
PARJ(47)	:	modification of Lund fragmentation for heavy quarks with Bowler, bottom
PARJ(54)	:	charm fragmentation functional form and value if MSTJ(11) = 2 or 3
PARJ(55)	:	bottom fragmentation functional form and value if MSTJ(11) = 2 or 3
PARJ(81)	:	Λ_{QCD} for parton showers
PARJ(82)	:	invariant mass cut-off for parton showers

Fragmentation functions and spin structure of the nucleon


- Unpolarized fragmentation functions:

- Provide flavor information in nucleon
- Most apparent in SIDIS measurements related to $\Delta q(x)$
- But also required for all RHIC hadron asymmetries (especially pion A_{LL} charge ordering)
- Transverse momentum dependence needed for Sivers and other TMDs


- Polarized fragmentation functions:

- For transverse spin almost unique access (require two chiral-odd functions):
 - DY: $\delta q \times \delta q$ or
 - SIDIS/RHIC: $\delta q \times \text{Collins}$ or $\delta q \times \text{IFF}$
- FFs from Belle/Babar


Thrust distributions (lin and log)






Pions vs z (lin)




Pions vs z (log)








Z smearing (integrating over ktbins)




Thrust smearing (integrating over z and k_t bins)



Belle fragmentation

- Initially started by RBRC fellow M. Perdekamp (now UIUC), A. Ogawa (BNL) and S. Lange (now U Giessen)
- Continued by RCS with additional help from UIUC (Matthias) and IU/Duke (A. Vossen) and JSPS




New possibilities:

From Correcting ISR to using it as a tool

- During di-hadron analysis and kt analysis successfully checked actual boost of qqbar pair in ISR events
- General Idea: Use photons reconstructed in Belle(2) to tune the reduced sqrt(s) and scale of the fragmenting qqbar system
- Belle(2) acceptance covers a larger range with EM Calorimetry than for tracking → good at catching not too soft photons
- First test: use all photons that cannot be combined with another photon to be close to the π^0 (0.1-0.17) or η (0.5-0.6) mass range


CMS energy of the qqbar system

reduced CMS energy



- Remove detected photon(s), $E > 0.5 \text{ GeV}$ from system
- Correlation to true CMS energy seen


reduced CMS energy - photon energy



- Possible improvements:
 - Correlation w/ angle to the thrust axis
 - Small angles to the beam directions
 - Mostly from the higher energy beam
 - Total reconstructed energy

New possibilities: jet-substructure+fragmentation

- Could jet-substructure work from HEP be used at EIC (and e+e-/pp/pA)?
 - Possibility to use jet charge and mass to distinguish quark and gluon jets might be very useful
 - even better flavor separation
 - Gluon background removal for chiral-odd measurements?
 - Up-down separation also possible? Sufficient discriminators available in e+e- or ep?



*Eur. Phys. J. C*76(6), 1-23 (2016)


→ Plan to study this in the next years (if JSPS has mercy)

New possibilities: other final state FFs needed?

- Extension of di-hadron analysis to any resonant hadron possible:
 - K_s , K^* , ϕ , ρ , etc
- πK and KK IFF measurements
- Other Collins measurements?

Hyperon Fragmentation

Belle: Niiyama et. al. [PRD 97 \(2018\), 072005](#)




$$x_p = \frac{Ph}{Ph_{max}}$$

- Hyperons similar to light hadron fragmentation \rightarrow peaking at low z (x_p)
- Baryon production not too well described by Pythia 6 default settings


Charmed baryon Fragmentation

Belle: Niiyama et. al. [PRD 97 \(2018\), 072005](#)



- Charmed baryons carry large fraction of parton momentum, similar to charmed mesons
- Charmed fragmentation reasonably described in Pythia for main states

Charmed Fragmentation



PRL.95, 142003 (2005)(Babar)
 PRD73, 032002 (2006) (Belle)
 PRD75, 012003 (2007)(Babar)
 PRL 99, 062001 (2007)(Babar)

- Heavier particles generally plotted vs normalized momentum $x_p = \frac{P^h}{P_{max}^h}$
- Unlike light hadrons charmed hadrons contain large fraction of charm quark momentum