

# Lattice TMD observables at the physical pion mass

Michael Engelhardt

New Mexico State University

In collaboration with:

B. Musch, P. Hägler, J. Negele, A. Schäfer

J. R. Green, N. Hasan, S. Krieg, S. Meinel, A. Pochinsky, S. Syritsyn

T. Bhattacharya, R. Gupta, B. Yoon

S. Liuti, A. Rajan


## Fundamental TMD correlator

$$\tilde{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \dots) \equiv \frac{1}{2} \langle P, S | \bar{q}(0) \Gamma \mathcal{U}[0, \dots, b] q(b) | P, S \rangle$$

$$\Phi^{[\Gamma]}(x, k_T, P, S, \dots) \equiv \int \frac{d^2 b_T}{(2\pi)^2} \int \frac{d(b \cdot P)}{(2\pi) P^+} \exp(i x (b \cdot P) - i b_T \cdot k_T) \frac{\tilde{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \dots)}{\tilde{\mathcal{S}}(b^2, \dots)} \Big|_{b^+=0}$$

- “Soft factor”  $\tilde{\mathcal{S}}$  required to subtract divergences of Wilson line  $\mathcal{U}$
- $\tilde{\mathcal{S}}$  is typically a combination of vacuum expectation values of Wilson line structures
- Here, will consider only ratios in which soft factors cancel

## Gauge link structure motivated by SIDIS



Beyond tree level: Rapidity divergences suggest taking staple direction slightly off the light cone. Approach of Aybat, Collins, Qiu, Rogers makes  $v$  space-like. Parametrize in terms of Collins-Soper parameter

$$\hat{\zeta} \equiv \frac{P \cdot v}{|P||v|}$$

Light-like staple for  $\hat{\zeta} \rightarrow \infty$ . Perturbative evolution equations for large  $\hat{\zeta}$ .

“Modified universality”,  $f^{\text{T-odd, SIDIS}} = -f^{\text{T-odd, DY}}$

## Fundamental TMD correlator

$$\tilde{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \dots) \equiv \frac{1}{2} \langle P, S | \bar{q}(0) \Gamma \mathcal{U}[0, \eta v, \eta v + b, b] q(b) | P, S \rangle$$

$$\Phi^{[\Gamma]}(x, k_T, P, S, \dots) \equiv \int \frac{d^2 b_T}{(2\pi)^2} \int \frac{d(b \cdot P)}{(2\pi) P^+} \exp(i x (b \cdot P) - i b_T \cdot k_T) \frac{\tilde{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \dots)}{\tilde{\mathcal{S}}(b^2, \dots)} \Big|_{b^+=0}$$

- “Soft factor”  $\tilde{\mathcal{S}}$  required to subtract divergences of Wilson line  $\mathcal{U}$
- $\tilde{\mathcal{S}}$  is typically a combination of vacuum expectation values of Wilson line structures
- Here, will consider only ratios in which soft factors cancel

## Decomposition of $\Phi$ into TMDs

All leading twist structures:

$$\Phi^{[\gamma^+]} = f_1 - \left[ \frac{\epsilon_{ij} k_i S_j}{m_H} f_{1T}^\perp \right] \text{odd}$$

$$\Phi^{[\gamma^+ \gamma^5]} = \Lambda g_1 + \frac{k_T \cdot S_T}{m_H} g_{1T}$$

$$\Phi^{[i\sigma^i \gamma^5]} = S_i h_1 + \frac{(2k_i k_j - k_T^2 \delta_{ij}) S_j}{2m_H^2} h_{1T}^\perp + \frac{\Lambda k_i}{m_H} h_{1L}^\perp + \left[ \frac{\epsilon_{ij} k_j}{m_H} h_1^\perp \right] \text{odd}$$

## TMD Classification

All leading twist structures:

$H$ $\downarrow$	$q \rightarrow$	U	L	T
U	$f_1$		$h_1^\perp$	← Boer-Mulders (T-odd)
L		$g_1$	$h_{1L}^\perp$	
T	$f_{1T}^\perp$	$g_{1T}$	$h_1 \quad h_{1T}^\perp$	

↑  
Sivers (T-odd)

## Decomposition of $\tilde{\Phi}$ into amplitudes

$$\tilde{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \hat{\zeta}, \mu) \equiv \frac{1}{2} \langle P, S | \bar{q}(0) \Gamma \mathcal{U}[0, \eta v, \eta v + b, b] q(b) | P, S \rangle$$

Decompose in terms of invariant amplitudes; at leading twist,

$$\begin{aligned} \frac{1}{2P^+} \tilde{\Phi}_{\text{unsubtr.}}^{[\gamma^+]} &= \tilde{A}_{2B} + im_H \epsilon_{ij} b_i S_j \tilde{A}_{12B} \\ \frac{1}{2P^+} \tilde{\Phi}_{\text{unsubtr.}}^{[\gamma^+ \gamma^5]} &= -\Lambda \tilde{A}_{6B} + i[(b \cdot P)\Lambda - m_H(b_T \cdot S_T)] \tilde{A}_{7B} \\ \frac{1}{2P^+} \tilde{\Phi}_{\text{unsubtr.}}^{[i\sigma^{i+} \gamma^5]} &= im_H \epsilon_{ij} b_j \tilde{A}_{4B} - S_i \tilde{A}_{9B} \\ &\quad - im_H \Lambda b_i \tilde{A}_{10B} + m_H[(b \cdot P)\Lambda - m_H(b_T \cdot S_T)] b_i \tilde{A}_{11B} \end{aligned}$$

(Decompositions analogous to work by Metz et al. in momentum space)

## Relation between Fourier-transformed TMDs and invariant amplitudes $\tilde{A}_i$

Invariant amplitudes directly give selected  $x$ -integrated TMDs in Fourier ( $b_T$ ) space (showing just the ones relevant for Sivers, Boer-Mulders shifts), up to soft factors:

$$\tilde{f}_1^{[1](0)}(b_T^2, \hat{\zeta}, \dots, \eta v \cdot P) = 2\tilde{A}_{2B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P) / \tilde{S}(b^2, \dots)$$

$$\tilde{f}_{1T}^{\perp[1](1)}(b_T^2, \hat{\zeta}, \dots, \eta v \cdot P) = -2\tilde{A}_{12B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P) / \tilde{S}(b^2, \dots)$$

$$\tilde{h}_1^{\perp[1](1)}(b_T^2, \hat{\zeta}, \dots, \eta v \cdot P) = 2\tilde{A}_{4B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P) / \tilde{S}(b^2, \dots)$$



## Generalized shifts

Form ratios in which soft factors, ( $\Gamma$ -independent) multiplicative renormalization factors cancel

Boer-Mulders shift:

$$\langle k_y \rangle_{UT} \equiv m_H \frac{\tilde{h}_1^{\perp[1](1)}}{\tilde{f}_1^{[1](0)}} = \frac{\int dx \int d^2 k_T k_y \Phi[\gamma^+ + s^j i \sigma^{j+} \gamma^5](x, k_T, P, \dots)}{\int dx \int d^2 k_T \Phi[\gamma^+ + s^j i \sigma^{j+} \gamma^5](x, k_T, P, \dots)} \Big|_{s_T=(1,0)}$$

Average transverse momentum of quarks polarized in the orthogonal transverse (“ $T$ ”) direction in an unpolarized (“ $U$ ”) hadron; normalized to the number of valence quarks. “Dipole moment” in  $b_T^2 = 0$  limit, “shift”.

**Issue:**  $k_T$ -moments in this ratio singular; generalize to ratio of Fourier-transformed TMDs at *nonzero*  $b_T^2$ ,

$$\langle k_y \rangle_{UT}(b_T^2, \dots) \equiv m_H \frac{\tilde{h}_1^{\perp[1](1)}(b_T^2, \dots)}{\tilde{f}_1^{[1](0)}(b_T^2, \dots)}$$

(remember singular  $b_T \rightarrow 0$  limit corresponds to taking  $k_T$ -moment). “Generalized shift”.

## Generalized shifts from amplitudes

Now, can also express this in terms of invariant amplitudes:

$$\langle k_y \rangle_{UT}(b_T^2, \dots) \equiv m_H \frac{\tilde{h}_1^{\perp[1](1)}(b_T^2, \dots)}{\tilde{f}_1^{[1](0)}(b_T^2, \dots)} = m_H \frac{\bar{A}_{4B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}{\bar{A}_{2B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}$$

Analogously, Sivers shift (in a polarized hadron):

$$\langle k_y \rangle_{TU}(b_T^2, \dots) = -m_H \frac{\bar{A}_{12B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}{\bar{A}_{2B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}$$


Worm-gear ( $g_{1T}$ ) shift:

$$\langle k_x \rangle_{TL}(b_T^2, \dots) = -m_N \frac{\bar{A}_{7B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}{\bar{A}_{2B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}$$

Generalized tensor charge (no  $k$ -weighting) :

$$\frac{\tilde{h}_1^{[1](0)}}{\tilde{f}_1^{[1](0)}} = - \frac{\bar{A}_{9B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P) - (m_N^2 b^2 / 2) \bar{A}_{11B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}{\bar{A}_{2B}(-b_T^2, 0, \hat{\zeta}, \eta v \cdot P)}$$

## Lattice setup





- Evaluate directly  $\bar{\Phi}_{\text{unsubtr.}}^{[\Gamma]}(b, P, S, \hat{\zeta}, \mu)$   
 $\equiv \frac{1}{2} \langle P, S | \bar{q}(0) \Gamma \mathcal{U}[0, \eta v, \eta v + b, b] q(b) | P, S \rangle$
- Euclidean time: Place entire operator at one time slice, i.e.,  $b, \eta v$  purely spatial
- Since generic  $b, v$  space-like, no obstacle to boosting system to such a frame!
- **Parametrization of correlator in terms of  $\tilde{A}_i$  invariants** permits direct translation of results back to original frame; form desired  $\tilde{A}_i$  ratios.
- Use variety of  $P, b, \eta v$ ; here  $b \perp P, b \perp v$  (lowest  $x$ -moment, kinematical choices/constraints)
- Extrapolate  $\eta \rightarrow \infty, \hat{\zeta} \rightarrow \infty$  numerically.





## Results: Sivers shift

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Sivers shift

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Sivers shift

Dependence on staple extent; sequence of panels at different  $|b_T|$



## Results: Sivers shift



Dependence of SIDIS limit on  $|b_T|$







## Results: Sivers shift

Dependence of SIDIS limit on  $\hat{\zeta}$





## Results: Boer-Mulders shift

Dependence of SIDIS limit on  $|b_T|$





## Results: Transversity

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Transversity

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Transversity

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Transversity

Dependence on staple extent; sequence of panels at different  $|b_T|$





## Results: Transversity

Dependence of SIDIS/DY limit on  $|b_T|$



## Results: Transversity



Dependence of SIDIS/DY limit on  $\hat{\zeta}$





## Results: $g_{1T}$ worm gear shift


Dependence of SIDIS/DY limit on  $|b_T|$



Approaching the light cone


## Results: Boer-Mulders shift (pion)

Dependence of SIDIS limit on  $\hat{\zeta}$ ; open symbols: contribution  $\tilde{A}_4$  only




## Results: Boer-Mulders shift (pion)

Dependence of SIDIS limit on  $\hat{\zeta}$ ; fit function  $a + b/\hat{\zeta}$





## Momentum smearing - preliminary results in a nucleon

Dependence of SIDIS limit on  $\hat{\zeta}$



## Advertisement: Dependence of Sivers shift on momentum fraction $x$



## Conclusions and Outlook

- Continued exploration of transverse quark dynamics using bilocal quark operators with staple-shaped gauge link structures. Soft factors, multiplicative renormalizations are canceled by constructing appropriate ratios of Fourier-transformed TMDs.
- Progress on challenges posed by physical pion mass limit,  $\hat{\zeta} \rightarrow \infty$  limit, discretization effects.
- Currently analyzing initial data on the dependence of the Sivers shift on momentum fraction  $x$ .
- Plan to explore further TMD/GTMD observables (longitudinal polarization, quark orbital angular momentum, spin-orbit coupling, twist-3).