# Pion valence structure and form-factors from lattice QCD at physical point

Xiang Gao BNL & THU

9th Workshop of the APS Topical Group on Hadronic Physics



## Pion valence quark PDF

Pion play a central role in the study of the strong interactions.

$$m_{\pi} \approx 140 \text{ MeV} \xrightarrow{\text{chiral limit}}{m_{q}=0} 0$$

- Critical ingredient for understanding the dynamical chiral symmetry breaking in QCD.
- Quarks and gluons in massless NG bosons.
- ...





3



**Quasi PDF:** • X. Ji, PRL 110 (2013); SCPMA57 (2014);

$$\tilde{q}(x) \equiv \int \frac{dz}{4\pi} e^{-ixP_z z} \langle P | \tilde{O}_{\Gamma}(z,\epsilon) | P \rangle,$$
  
$$\tilde{O}_{\Gamma}(z,\epsilon) = \overline{\psi}(0) \Gamma W_{\hat{z}}(0,z) \psi(z)$$

## Equal-time correlators and QCD factorization

Short distance Factorization in coordinate space:

- V. Braun et al., EPJC 55 (2008)
- A. V. Radyushkin, PRD 96 (2017)
- Y. Ma et al., PRL 120 (2018)
- T. Izubuchi et al., PRD 98 (2018)



### **Ratio scheme renormalization:**

$$\mathcal{M}(z, P_z, P_z^0) = \frac{\langle P_z | \tilde{O}_{\Gamma}(z, a) | P_z \rangle}{\langle P_z^0 | \tilde{O}_{\Gamma}(z, a) | P_z^0 \rangle}$$

$$\tilde{O}_{\Gamma}(z,\mu) = Z_{\psi,z} e^{\delta m|z|} \tilde{O}$$

Hadron state independent



## Pion valence quark PDF: NLO results

## **Boosted pion state on the lattice**



4

Moments of pion valence quark PDF



### **Pion valence quark PDF**



### **Improvement:**

- Matching formula beyond one-loop. •
- Computation with physical pion mass.
- Extract PDFs information from chiral fermions.



Pion valence quark PDF: NNLO 5

### **Improvement:**

- Matching formula beyond one-loop.
- Computation with physical pion mass.
- Extract PDFs information from chiral fermions.

### • NNLO matching

• Li, Ma and Qiu, PRL 126 (2021)

$$C_n(z^2\mu^2) = 1 + \alpha_s(\mu)C_n^{(1)}(z^2\mu^2) + \alpha_s^2(\mu)C_n^{(2)}(z^2\mu^2) + \mathcal{O}(\alpha)$$

$$= 1 + \frac{\alpha_s(\mu)C_F}{2\pi} \left[ \left( \frac{3+2n}{2+3n+n^2} + 2H_n \right) \ln(z_0^2 \mu^2) + \dots \right] + z_0^2 = z^2$$

When  $\ln(z_0^2 \mu^2)$  become large, one may need to include the **DGLAP** evolution:

$$\left[\frac{\partial}{\partial \ln \mu^2} + \beta(\alpha_s(\mu))\frac{\partial}{\partial \alpha_s} - \gamma_n\right]C_n^{evo} = 0$$

- A. V. Radyushkin, PLB 781 (2018)
- BNL, ANL, arXiv: 2102.01101



- Clear z<sub>0</sub> dependence can be observed at LO.
- Moments evolved from  $1/z_0$  to  $\mu$  from NNLO are consistent with **NLO** with current statistics but more flat, and agree with the **DGLAP** improved case.



Pion valence quark PDF: Improvement 6

### **Improvement:**

- Matching formula beyond one-loop.
- Computation with physical pion mass.
- Extract PDFs information from chiral fermions.

## • Physical pion mass

Ratio scheme renormalized matrix elements



## • Chiral fermion





## Pion valence quark PDF: Moments



matching coefficients.

• To stabilize the fit and extract higher moments, we perform **combine analysis** with data in range  $[3a, z_{max}]$  using **NNLO** 



## Pion valence quark PDF: Moments

8

## Moments: NNLO matching, physical point, chiral fermion



- The mass dependence is mild for pion valence PDF.
- spacings.

Chiral fermion shows good agreement with Wilson-Clover + HISQ fermion with fine lattice

![](_page_7_Picture_8.jpeg)

9

## Pion valence quark PDF

![](_page_8_Figure_2.jpeg)

Preliminary results of the large-x behavior from model  $x^{\alpha}(1-x)^{\beta}(1+t\sqrt{x}+sx)$ :

$$\beta = 1.07(37)(29),$$

which shows good agreement with JAM18, xFitter.

More improvement:

- Resummation in perturbative matching. For example, NLO+NLL threshold resummation (BNL, ANL 21 arXiv: 2102.01101).
- More statistics and large momentum to extract higher moments.

![](_page_8_Picture_10.jpeg)

Pion form factor and charge radius 10

![](_page_9_Figure_1.jpeg)

 $(p_1 + p_2)^{\mu} F_{\pi}(Q^2)$ 

## Pion elastic electromagnetic form factor $F_{\pi}(Q^2)$

- The 3-D generalized PDFs (GPDs) combine the information contained in **PDFs** and **form** factors.
- The mean charge radius is related to form factors at low  $Q^2$  .
- EIC facility will allow higher  $Q^2$  up to 30  $GeV^2$ , make contact with pQCD.

![](_page_9_Picture_9.jpeg)

### Pion form factor and charge radius 11

![](_page_10_Figure_1.jpeg)

$$r_{\pi}^2 = -6 \frac{dF_{\pi}(Q^2)}{dQ^2}|_{Q^2=0}$$

$$\frac{1}{2}, t = -Q^2$$

![](_page_10_Picture_4.jpeg)

![](_page_11_Picture_0.jpeg)

- We studied pion valence quark PDF with multiple lattices and pion mass.
- The mass dependence of pion valence PDF is mild.
- The usage of Wilson-Clover fermion didn't bring trouble to the determination of pion valence structure.
- We calculate the pion electromagnetic form factors which show good agreement with experimental data.

## Summary

![](_page_11_Picture_7.jpeg)