

# IX-APS TOPICAL GROUP ON HADRONIC PHYSICS



## The structure of nuclei from Drell-Yan and Sidis data

Ignazio Scimemi, Work in progress with M. Bury, F. Hautmann, S. Leal Gomez, A. Vladimirov, P. Zurita

# TMD extraction

## Factorization

$$\frac{d\sigma}{dQ^2 dy dq_T^2} = \sigma_0 \sum_{f_1, f_2} \int \frac{d^2 \mathbf{b}}{4\pi} e^{i(\mathbf{b} \cdot \mathbf{q}_T)} H_{f_1 f_2}(Q, \mu) F_{f_1 \leftarrow h_1}(x_1, \mathbf{b}; \mu, \zeta_1) F_{f_2 \leftarrow h_2}(x_2, \mathbf{b}; \mu, \zeta_2).$$

Scale-independent  
Factors!

$$\frac{d\sigma}{dQ^2 dy dq_T^2} = \sigma_0 \sum_{f_1, f_2} \int \frac{d^2 \mathbf{b}}{4\pi} e^{i(\mathbf{b} \cdot \mathbf{q}_T)} H_{f_1 f_2}(Q, Q) \cancel{R[\mathbf{b}; (Q, Q^2)]}^2 F_{f_1 \leftarrow h_1}(x_1, \mathbf{b}) F_{f_2 \leftarrow h_2}(x_2, \mathbf{b})$$

Separation of NP effects  
in evolution kernel and TMD  
 $\zeta$ -prescription

E288(200)		CMS (7TeV)
E288(300)		CMS (8TeV)
E288(400)	CDF (run1) CDF (run2)	ATLAS (7TeV)
E605	D0 (run1) D0 (run2)	ATLAS (8TeV)
E772	D0 (run2)	ATLAS (8TeV)
		LHCb (7TeV)
		LHCb (8TeV)
		LHCb (13TeV)

## Data selection

Statistical methods:  
replicas of exp

$$V_{ij} = (\sigma_{i,\text{stat}}^2 + \sigma_{i,\text{unc}}^2) \delta_{ij} + \sum_{l=1}^k \sigma_{i,\text{corr}}^{(l)} \sigma_{j,\text{corr}}^{(l)}$$

## RESULTS

# TMD extraction

## Factorization

$$\frac{d\sigma}{dQ^2 dy dq_T^2} = \sigma_0 \sum_{f_1, f_2} \int \frac{d^2 \mathbf{b}}{4\pi} e^{i(\mathbf{b} \cdot \mathbf{q}_T)} H_{f_1 f_2}(Q, \mu) F_{f_1 \leftarrow h_1}(x_1, \mathbf{b}; \mu, \zeta_1) F_{f_2 \leftarrow h_2}(x_2, \mathbf{b}; \mu, \zeta_2).$$

## Scale-independent Factors!

$$\frac{d\sigma}{dQ^2 dy dq_T^2} = \sigma_0 \sum_{f_1, f_2} \int \frac{d^2 \mathbf{b}}{4\pi} e^{i(\mathbf{b} \cdot \mathbf{q}_T)} H_{f_1 f_2}(Q, Q) \{R[\mathbf{b}; (Q, Q^2)]\}^2 F_{f_1 \leftarrow h_1}(x_1, \mathbf{b}) F_{f_2 \leftarrow h_2}(x_2, \mathbf{b})$$

3 independent functions.  
All functions are scale independent in

$\zeta$ -prescription

# Unpolarized TMDPDF

$\zeta$ -prescription *JHEP* 08 (2018) 003

$$\lim_{b \rightarrow 0} F_{1,f \leftarrow h}(x,b) = \sum_{f'} \int_x^1 \frac{dy}{y} C_{f \leftarrow f'} \left( \frac{x}{y}, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) f_{1,f' \leftarrow h}(y, \mu_{\text{OPE}}),$$
$$\lim_{b \rightarrow 0} D_{1,f \rightarrow h}(z,b) = \sum_{f'} \int_z^1 \frac{dy}{y} C_{f \rightarrow f'} \left( \frac{z}{y}, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) \frac{d_{1,f' \rightarrow h}(y, \mu_{\text{OPE}})}{y^2},$$

PDF ARE PART  
OF THE MODEL

NNLO

TMDPDF: T. Gehrmann et al. *JHEP* 06 (2014) 155,  
TMDPDF and TMDFF: M.G. Echevarria et al. *Phys. Rev.* D93 (2016) 011502, *JHEP* 09 (2016) 004

NNNLO TMDPDF: M. X. Luo et al. *Phys. Rev. Lett.* 124 (2020) 9, 092001, M. Ebert et al. *JHEP* 09 (2020) 146

TMDFF: M. Ebert et al. e-Print:2012.07853

F

# Unpolarized TMDPDF

$\zeta$ -prescription *JHEP* 08 (2018) 003

$$\lim_{b \rightarrow 0} F_{1,f \leftarrow h}(x, b) = \sum_{f'} \int_x^1 \frac{dy}{y} C_{f \leftarrow f'} \left( \frac{x}{y}, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) f_{1,f' \leftarrow h}(y, \mu_{\text{OPE}}),$$

$$\lim_{b \rightarrow 0} D_{1,f \rightarrow h}(z, b) = \sum_{f'} \int_z^1 \frac{dy}{y} C_{f \rightarrow f'} \left( \frac{z}{y}, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) \frac{d_{1,f' \rightarrow h}(y, \mu_{\text{OPE}})}{y^2},$$

PDF ARE PART  
OF THE MODEL

NNLO

TMDPDF: T. Gehrmann et al. *JHEP* 06 (2014) 155,

TMDPDF and TMDFF: M.G. Echevarria et al. *Phys. Rev.* D93 (2016) 011502, *JHEP* 09 (2016) 004

NNNLO TMDPDF: M. X. Luo et al. *Phys. Rev. Lett.* 124 (2020) 9, 092001, M. Ebert et al. *JHEP* 09 (2020) 146

TMDFF: M. Ebert et al. e-Print:2012.07853

$$F_{1,f \leftarrow h}(x, b) = \int_x^1 \frac{dy}{y} \sum_{f'} C_{f \leftarrow f'} \left( y, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) f_{1,f' \leftarrow h} \left( \frac{x}{y}, \mu_{\text{OPE}} \right) f_{\text{NP}}(x, b),$$

$$D_{1,f \rightarrow h}(z, b) = \frac{1}{z^2} \int_z^1 \frac{dy}{y} \sum_{f'} y^2 C_{f \rightarrow f'} \left( y, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}}) \right) d_{1,f' \rightarrow h} \left( \frac{z}{y}, \mu_{\text{OPE}} \right) D_{\text{NP}}(z, b)$$

NNLO-PDF

NLO-FF

Bottleneck!!!

$$f_{\text{NP}}(x, b) = \exp \left( -\frac{\lambda_1(1-x) + \lambda_2 x + x(1-x)\lambda_5 b^2}{\sqrt{1+\lambda_3 x^{\lambda_4} b^2}} \right)$$

$$D_{\text{NP}}(x, b) = \exp \left( -\frac{\eta_1 z + \eta_2(1-z) b^2}{\sqrt{1+\eta_3(b/z)^2} z^2} \right) \left( 1 + \eta_4 \frac{b^2}{z^2} \right),$$

Ansatz for Artemide in SV19

# The evolution

$$R(b, Q, \mu) = \left( \frac{Q^2}{\zeta_\mu[\mathcal{D}(b, \mu)]} \right)^{-2\mathcal{D}(b, \mu)}$$


$$\mathcal{D}(b, \mu) = \mathcal{D}_{\text{pert}}(b, \mu) + d_{\text{NP}}(b)$$

N3LO:

Y. Li, H.X. Zhu, Phys. Rev. Lett. 118, 022004  
(2017)

A. Vladimirov, Phys. Rev. Lett. 118 (2017) 6,  
062001

Consistency of extractions  
Of the evolution kernel  
In DY and DY+SIDIS



# Unpolarized TMD and data

---

Fit from DY:

D'Alesio et al. JHEP11 (2014) 098 ([Evolution kernel](#))

Bacchetta et al, JHEP06 (2017)081

I. S., A. Vladimirov Eur. Phys. J. C. 78 (2018) 2, 89 ([Artemide](#))

I. S., A. Vladimirov JHEP06 (2019) 028

V. Bertone, I.S., A. Vladimirov JHEP10 (2019)090 ([pion-proton](#))

A.Vladimirov JHEP10 (2019)090 ([pion-proton](#))

V. Bertone et at. JHEP07(2020) 117 ([Nangaparbat](#)) ([Pavia19](#))

M. Bury et al. In progress

Fit from DY+SIDIS: I. S., A.Vladimirov JHEP06 (2020) 137([SV19](#))

**W-production: D. Gutierrez-Reyes, S. Leal-Gomez, I.S. e-print 2011.05351 [hep-ph]**

# Open questions


---

- ✿ Which data are sensitive to TMD?
- ✿ Do TMD fits depend on the choice of PDF sets?
- ✿ Do we have a unique  $f_{NP}$  for all sets of PDF?

# Data sets distributions


High Energy (only DY):  
LHC+TeVatron

Low Energy (DY+SIDIS)




# Forgotten data at LHC?

For TMD we need LHC data with low boson mass: ex. LHCb-CONF-2012-013 ; CERN-LHCb-CONF-2012-013




Please address trigger problem  
at ATLAS, CMS


# TMD validity range



SV17, DY



Pavia19, DY




SV19, SIDIS

The range of validity of TMD is independent of models, implementation, perturbative order, experiment type

In SIDIS       $q_T \simeq p_\perp/z \rightarrow \delta = \frac{q_T}{Q} \simeq \frac{p_\perp}{zQ} < 0.25$

HERA data excluded, EIC necessary

# TMD validity range



*Terra Incognita (JLAB)*

$$q_T/Q \lesssim 0.25$$

Resummation region

Non-perturbative TMD region

$q_T/Q \lesssim 0.25 \text{ & } q_T/\Lambda \gg 1$  TMD factorization valid and dominated by perturbative effects

$q_T/Q \lesssim 0.25 \text{ & } q_T/\Lambda \sim 1$  TMD factorization valid and dominated by non-perturbative effects

# TMD and PDF

Do TMD fit quality depend on PDF set choice?  
In principle the TMD is independent but ...


$$F_{f \leftarrow h}(x, b) = \sum_{f'} \int_x^1 \frac{dy}{y} C_{f \leftarrow f'}(y, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}})) f_{f' \leftarrow h} \left( \frac{x}{y}, \mu_{\text{OPE}} \right) f_{\text{NP}}(x, b)$$

↑  
Matching (Wilson)  
coefficient NNLO

↑  
PDF

↑  
Gaussian?  
Exponential?


- Every PDF set has its own error band
- Spread among different sets



# Differences among PDF sets are visible in TMD physics



DY fit from I.S., A. Vladimirov JHEP 06 (2020) 137

PDF set	$\chi^2/N_{pt}$	Parameters for $\mathcal{D}$	Parameters for $f_1$	
HERA20	0.97	$B_{NP} = 2.29 \pm 0.43$ $c_0 = (2.22 \pm 0.93) \cdot 10^{-2}$	$\lambda_1 = 0.324 \pm 0.029$ $\lambda_2 = 13.2 \pm 2.9$	$\lambda_3 = (3.56 \pm 1.59) \cdot 10^2$ $\lambda_4 = 2.05 \pm 0.26$ $\lambda_5 = -10.4 \pm 3.5$
NNPDF31	1.14	$B_{NP} = 1.86 \pm 0.30$ $c_0 = (2.96 \pm 1.04) \cdot 10^{-2}$	$\lambda_1 = 0.253 \pm 0.032$ $\lambda_2 = 9.0 \pm 3.0$	$\lambda_3 = (3.47 \pm 1.16) \cdot 10^2$ $\lambda_4 = 2.48 \pm 0.15$ $\lambda_5 = -5.7 \pm 3.4$
MMHT14	1.34	$B_{NP} = 1.55 \pm 0.29$ $c_0 = (4.70 \pm 1.77) \cdot 10^{-2}$	$\lambda_1 = 0.198 \pm 0.040$ $\lambda_2 = 26.4 \pm 4.9$	$\lambda_3 = (26.8 \pm 13.2) \cdot 10^3$ $\lambda_4 = 3.01 \pm 0.17$ $\lambda_5 = -23.4 \pm 5.4$
PDF4LHC	1.53	$B_{NP} = 1.93 \pm 0.47$ $c_0 = (3.66 \pm 2.09) \cdot 10^{-2}$	$\lambda_1 = 0.218 \pm 0.041$ $\lambda_2 = 17.9 \pm 4.5$	$\lambda_3 = (9.26 \pm 8.38) \cdot 10^2$ $\lambda_4 = 2.54 \pm 0.17$ $\lambda_5 = -15.5 \pm 4.7$
CT14	1.59	$B_{NP} = 2.35 \pm 0.61$ $c_0 = (2.27 \pm 1.33) \cdot 10^{-2}$	$\lambda_1 = 0.277 \pm 0.029$ $\lambda_2 = 24.9 \pm 2.9$	$\lambda_3 = (12.4 \pm 3.2) \cdot 10^3$ $\lambda_4 = 2.67 \pm 0.13$ $\lambda_5 = -23.8 \pm 2.9$



# TMD uncertainties in W production

D. Gutierrez-Reyes,  
S. Leal-Gomez, I.S.  
arXiv: 2011.05351



Scale uncertainty  
(More study in the future)

Replicas for  
NNPDF31\_nnlo\_118


PDF sets

# TMD uncertainties in W production


D. Gutierrez-Reyes,  
 S. Leal-Gomez, I.S.  
 arXiv: 2011.05351

$\chi^2/N$  includes PDF, data and theoretical error

	CDF $\sqrt{s} = 1.8$ TeV	D0 $\sqrt{s} = 1.8$ TeV	ATLAS	CMS $e\nu$	CMS $\mu\nu$
Number of points	10	10	2	4	4
NNPDF31	0.540	1.485	0.463	1.674	3.165
HERA20	0.469	1.591	0.271	1.563	3.721



# TMD uncertainties from a single PDF set



I.S., A. Vladimirov *JHEP* 06 (2020) 137

Experimental uncertainty  
 PDF uncertainty (replicas, NNPDF31\_nnlo\_118)

# TMD and PDF sets: bias removing

• Spread among different sets

$$F_{f \leftarrow h}(x, \mathbf{b}) = \sum_{f'} \int_x^1 \frac{dy}{y} C_{f \leftarrow f'}(y, \mathbf{L}_{\mu_{\text{OPE}}}, a_s(\mu_{\text{OPE}})) f_{f' \leftarrow h} \left( \frac{x}{y}, \mu_{\text{OPE}} \right) f_{NP}(x, b)$$

SV19 ansatz:

$$\boxed{f_{NP}(x, b) = \exp \left( -\frac{\lambda_1(1-x) + \lambda_2 x + x(1-x)\lambda_5}{\sqrt{1+\lambda_3 x^{\lambda_4}} b^2} b^2 \right)}$$
$$D_{NP}(x, b) = \exp \left( -\frac{\eta_1 z + \eta_2(1-z)}{\sqrt{1+\eta_3(b/z)^2}} \frac{b^2}{z^2} \right) \left( 1 + \eta_4 \frac{b^2}{z^2} \right)$$

BHLSVZ21 ansatz:

$$f_{NP}(x, b) = \exp \left( -\frac{\lambda_1(1-x) + \lambda_2 x + x(1-x)\lambda_5}{\sqrt{1+\lambda_3 x^{\lambda_4}} b^2} b^2 \right)$$
$$D_{NP}(x, b) = \exp \left( -\frac{\eta_1 z + \eta_2(1-z)}{\sqrt{1+\eta_3(b/z)^2}} \frac{b^2}{z^2} \right) \left( 1 + \eta_4 \frac{b^2}{z^2} \right)$$

*Top Secret:  
Preliminary*

The non-perturbative ansatz used in previous fits is too rigid:  
We need flavor dependence of the ansatz to compensate the differences in different PDF sets


# TMD and PDF sets: preliminary results

---

PDF	$\chi^2/N$
NNPDF31	0.97
HERA20	0.90
CT18	0.98
MSHT20	0.88


The spread in the fit quality does not depend on PDF sets

# TMD uncertainties from PDF sets



**The goodness of a set depends on  $x$**

# TMD uncertainties from PDF sets



---

# Conclusions

---


- Great progress in the extraction of unpolarized TMD: Higher order perturbative extractions available. The critical point is the inclusion of collinear functions in TMD extractions (NNLO fragmentation functions are needed)
- W production is included in Artemide. W and DY data at LHC can be largely improved looking at neglected regions of phase space
- JLAB and EIC are/will be exploring fundamental regions of phase space
- PDF set dependence of TMD is removed with flavor dependent models

Back slides

# Non perturbative TMD effects at LHC

■  $R_\sigma = 2 \frac{d\sigma_{\text{test}} - d\sigma_{\text{TMD}}}{d\sigma_{\text{test}} + d\sigma_{\text{TMD}}}$

■ Scales uncertainty



F. Hautmann, I.S., A. Vladimirov *Phys.Lett.B* 806 (2020) 135478



# Non perturbative TMD effects at LHC

We have tested several simple models for LHC with/without NP effects in TMD.

A non-perturbative part on evolution kernel is always necessary and it is present in every code. We tested several possibilities.

Models with an fNP different from 1 give better agreement with LHC data.

Case	$B_{NP}$	$g_K$	$\lambda_1 (f_{NP} = \exp -\lambda_1 b^2)$	$\chi^2/dof$	$\chi^2/dof(\text{norm.})$
1	5.5 (max)	$0.116 \pm 0.002$	$10^{-3}(\text{fixed})$	3.29	3.04
2	$2.2 \pm 0.4$	$0.032 \pm 0.006$	$0.29 \pm 0.02$	1.50	1.28
Case	$B_{NP}$	$c_0$	$\lambda_1$	$\chi^2/dof$	$\chi^2/dof(\text{norm.})$
3	1. (min)	$0.016 \pm 0.001$	$10^{-3}(\text{fixed})$	2.21	1.99
4	$3.0 \pm 1.5$	$0.04 \pm 0.02$	$0.27 \pm 0.04$	1.61	1.36
Case	$B_{NP}$	$g_K^*$	$\lambda_1$	$\chi^2/dof$	$\chi^2/dof(\text{norm.})$
5	$1.34 \pm 0.01$	$0.16 \pm 0.01$	$10^{-3}(\text{fixed})$	1.70	1.52
6	$2.43 \pm 0.66$	$0.05 \pm 0.02$	$0.24 \pm 0.04$	1.49	1.28



F. Hautmann, I.S., A. Vladimirov *Phys.Lett.B* 806 (2020) 135478

# Correlation of parameters

