




UCLA Mani L. Bhaumik Institute
for Theoretical Physics

Nuclear TMDs and 3D imaging in nuclei (arXiv:2107.12401)

John Terry


In collaboration with Mishary Alrashed, Daniele Paolo Anderle,
Zhong-bo Kang, and Hongxi Xing

July 29, 2021




Overview of Global Analyses

Factorization Theorems




Central idea of a global analysis

- Calculate the perturbative physics at high order.
- We take an ansatz for the NP physics.
- Use experimental data constrain parameterization.



Nuclear Modified PDFs


Four Decades of the EMC Effect



This effect is still not understood from first principles.

Effective understanding of the nuclear modifications.

- Eskola, Kolhinen, Ruuskanen
1998: Kept DGLAP evolution equations unchanged.
- Found that global experimental data can be described by modifying NP parameterization.
- Indication that nuclear modifications may be non-perturbative in nature.




Nuclear Modified FFs

Extractions

Sassot, Stratmann, Zurita 2009 (SSZ)

Zurita 2020 (LIKEN)




Data from SIDIS + pA → h collisions.

- No modification to the DGLAP evolution.
- Accounted for nuclear medium effects in the non-perturbative parameterization.

$$D_{i/A}^H(z, Q_0^2) = [W_i \otimes D_i^H](z, Q_0^2)$$

$$W_q(y, A, Q_0^2) = n_q y^\alpha (1-y)^\beta + n'_q \delta(1-\varepsilon-y)$$



Nuclear Holography in Three-Dimensions

From collinear distributions to TMDs




Drell-Yan Measurements

- $R_{AB} = \frac{d\sigma_A}{dq_{\perp}} / \frac{d\sigma_B}{dq_{\perp}}$
 - E866
 - E772
 - Prelim. RHIC
- $d\sigma/dq_{\perp}$ (p Pb)
 - ATLAS
 - CMS

SIDIS Measurements

- Multiplicity ratio $R_h^A = M_h^A/M_h^D$.
 - HERMES 2007
 - Prelim. JLab
 - Planned JLab
 - Possible EIC.



Parameterization

Cross section

$$\frac{d\sigma^A}{dx dQ^2 dz d^2 P_{h\perp}} = \sigma_0 H(Q) \sum_q e_q^2 \int_0^\infty \frac{b db}{2\pi} J_0 \left(\frac{b P_{h\perp}}{z} \right) f_{q/n}^A(x, b; Q) D_{h/q}^A(z, b; Q)$$

TMDs

$$f_{q/n}^A(x, b; Q) = \left[C_{q \leftarrow i} \otimes f_{i/n}^A \right] (x, \mu_{b_*}) \exp \left\{ -S_{\text{pert}}(\mu_{b_*}, Q) - S_{\text{NP}}^f(b, Q, A) \right\}$$

$$D_{h/q}^A(z, b; Q) = \frac{1}{z^2} \left[\hat{C}_{i \leftarrow q} \otimes D_{h/i}^A \right] (z, \mu_{b_*}) \exp \left\{ -S_{\text{pert}}(\mu_{b_*}, Q) - S_{\text{NP}}^D(b, z, Q, A) \right\}$$

Our assumptions

- Perturbative information is left unchanged by the nuclear medium.
 $C_{q \leftarrow i}$, $\hat{C}_{i \leftarrow q}$, and S_{pert} are unchanged.
- Non-perturbative information is modified.
 $f_{i/n}^A$, $D_{h/i}^A$, S_{NP}^D , and S_{NP}^f are altered.

Parameterization continued

TMDs

$$f_{q/n}^A(x, b; Q) = \left[C_{q \leftarrow i} \otimes f_{i/n}^A \right] (x, \mu_{b_*}) \exp \left\{ -S_{\text{pert}}(\mu_{b_*}, Q) - S_{\text{NP}}^f(b, Q, A) \right\}$$

$$D_{h/q}^A(z, b; Q) = \frac{1}{z^2} \left[\hat{C}_{i \leftarrow q} \otimes D_{h/i}^A \right] (z, \mu_{b_*}) \exp \left\{ -S_{\text{pert}}(\mu_{b_*}, Q) - S_{\text{NP}}^D(b, z, Q, A) \right\}$$

Collinear Distributions


We use the EPPS16 parameterization for $f_{i/n}^A$ (NLO).

We use the LIKEN parameterization for $D_{h/i}^A$ (NLO).

Perturbative order in our analysis

Work at NLO+NNLL for the TMDs.

We take the non-perturbative parameterization



$$S_{\text{NP}}^f(b, Q, A) = S_{\text{NP}}^f(b, Q) + a_N \left(A^{1/3} - 1 \right) b^2$$

$$S_{\text{NP}}^D(z, b, Q, A) = S_{\text{NP}}^D(z, b, Q) + b_N \left(A^{1/3} - 1 \right) \frac{b^2}{z^2}$$

Fitting Procedure and Data

Kinematic cuts


SIDIS

Drell-Yan

$$P_{h+} < 0.7 \text{ GeV} \quad z < 0.7$$

$$q_{\perp}/Q < 0.3$$


Description of the experimental data




$\chi^2/d.o.f = 1.045$ 126 points with $a_N = 0.0171 \pm 0.003$ and $b_N = 0.0144 \pm 0.001$

3D imaging in the nucleus

$$R_{u/p}^{\text{Au}}(x, k_\perp, Q_0) = \frac{f_{u/p}^{\text{Au}}(x, k_\perp, Q_0)}{f_{u/p}(x, k_\perp, Q_0)}$$




$$\mathcal{R}_{\pi^+/u}^{\text{Xe}}(z, p_\perp, Q_0) = \frac{D_{\pi^+/u}^{\text{Xe}}(z, p_\perp, Q_0)}{D_{\pi^+/u}(z, p_\perp, Q_0)}$$



Summary and Outlook

- There has been tremendous progress over the past few decades in extracting nuclear modified collinear distributions.
- We have performed the first extractions of the nuclear modified TMDPDF and TMDFFs.
- This allows us to quantify for the first time the nuclear broadening of these distributions.
- We expect that additional data from JLab, RHIC, the LHC, as well as future measurements at the EIC could help further constrain these distributions.



Thank you to the audience and the organizers!