#### 2021 EIC UG Meeting Early Career Workshop

#### **Multiple Parton Scattering and Gluon Saturation in Dijet Production at EIC**

arXiv:2104.04520

Yuan-Yuan Zhang (CCNU) **Collaborator : Xin-Nian Wang (LBL)** 



### Contents

- Introduction: Jet Transport Cofficient  $\hat{q}$  in cold nuclei
- Single Scattering Dijet & Double Scattering Dijet Cross Section
- Summary

• Nuclear Modification of Dijet in e+A (  $\Delta \phi$ ,  $|y_{l_a} - y_l|$ ,  $R_A$  dependence)



#### in QGP and Cold Nuclei -average pT transfer squared per unit length

6

5

2

0

MARTINI

HT-BW

HT-M

 $\hat{q}_N/T_{eff}^3$ (DIS)

0.1

**Electron Ion Collision** 

 $\hat{q} \approx 0.015 \text{ GeV}^2/fm$ P. Ru et al, arXiv:2004.00027 NB Chang et al, PRC 89.3 (2014): 034911 E Wang, XN Wang, PRL 89.16 (2002): 162301

Yuan-Yuan Zhang

0.2



JET Collaboration, PRC 90, 014909

Introduction

# **Existing efforts on** $\hat{q}$ in cold nuclei

High Twist Approach

- Use the pT broadening of final-state hadron (DIS) to extract  $\hat{q}$ 0  $\hat{q} \approx 0.015 \text{ GeV}^2/fm$ P. Ru et al, arXiv:2004.00027
- Use the suppression of leading hadron (DIS) to extract  $\hat{q}$ 0  $\hat{q} \approx 0.02 \text{ GeV}^2/fm$ NB Chang et al, PRC 89.3 (2014): 034911

Generalized High Twist Approach Zhang, Y. Y., Qin, G. Y., & Wang, X. N. (2019). PRD, 100(7), 074031.

- Relax  $l_{\perp}, l_{q\perp} \gg k_{\perp}$ , no  $k_{\perp}$  /twist expansion  $\rightarrow$  medium-induced radiation spectra 0
- Radiation spectra contain medium gluon TMD pdf  $\phi(x_G, k_{\perp})$  or TMD  $\hat{q}(k_{\perp})$ 0  $\hat{q} = d^2 k_{\perp} \hat{q} \left( k_{\perp} \right)$

Yuan-Yuan Zhang

$$= C \int d^2 k_{\perp} \rho \phi \left( x_g, k_{\perp} \right)$$

Introduction





# Dijet to probe gluon TMD pdf or TMD $\hat{q}(k_{\perp})$



- Generalized High Twist Approach deals with hard splitting Second scattering directly probe gluon TMD pdf
- 0 0
- Further include initial quark transverse momentum  $v_{\perp}$ 0

Yuan-Yuan Zhang

Introduction



## Dijet in e+A: Single scattering



$$\frac{d\sigma_{eA}^{o}}{dx_B dQ^2 dz d^2 l_\perp d^2 l_{q\perp}} = \frac{2\pi\alpha_{em}^2}{Q^4} \sum_q e_q^2 [1]$$

0  $\Delta_F(\vec{b}_{\perp}) = \left| dy_0^- \hat{q}_F(y_0^-, \vec{b}_{\perp}) \right| \text{ depend on } \hat{q}$ Yuan-Yuan Zhang

multiple soft interaction (eikonalized as gauge link)  $\rightarrow$  quark pT broadening pT broadening embedded in effective  $q_N(x, \vec{v}_{\perp}, \vec{b}_{\perp})$ , gaussian broadening with width Liang, Z. T., Wang, X. N., & Zhou, J. (2008). PRD, 77(12), 125010.

Single & Double Scattering Dijet







# Dijet in e+A : Double scattering

Under two-parton correlation factorization

 $T_{qg}^{A} = \rho_{A}(y_{0}^{-}) \otimes \rho_{A}(y_{1}^{-}) \otimes$ 

$$\frac{d\hat{\sigma}_{eA}^{D}}{dx_{B}dQ^{2}dzd^{2}l_{\perp}d^{2}l_{q\perp}} = \frac{2\pi\alpha_{em}^{2}}{Q^{4}}\sum_{q}e_{q}^{2}[1+(1-\frac{Q^{2}}{x_{B}s})^{2}]\frac{\alpha_{s}}{2\pi}\frac{1+z^{2}}{1-z}\frac{2\pi\alpha_{s}}{N_{c}}\int\frac{d^{2}k_{\perp}}{(2\pi)^{2}}\int d^{2}b_{\perp}dy_{0}^{-}dy_{1}^{-}$$
$$\rho_{A}(y_{0}^{-},\overrightarrow{b}_{\perp})\rho_{A}(y_{1}^{-},\overrightarrow{b}_{\perp})q_{N}(x_{B},\overrightarrow{v}_{\perp},\overrightarrow{b}_{\perp})\frac{\phi_{N}(x_{G},\overrightarrow{k}_{\perp})}{k_{\perp}^{2}}\left[\mathcal{N}_{g}^{\mathsf{nonLPM}}+\mathcal{N}_{g}^{\mathsf{qLPM}}+\mathcal{N}_{g}^{\mathsf{qLPM}}\right]$$

intial quark  $q_N(x, \vec{v}_{\parallel}, \vec{b}_{\parallel})$  contain pT broadening, depend on  $\hat{q}$  indirectly 0

depend on  $\phi(x_G, k_{\perp}) \sim \hat{q}(k_{\perp})$  directly 0

Yuan-Yuan Zhang

Single & Double Scattering Dijet



$$\bigotimes q_N(x_B, \overrightarrow{v}_{\perp}) \bigotimes \phi_N(x_G, \overrightarrow{k}_{\perp})$$



- harder than gauge link gluons



# Different contribution in double scattering

Contribution divided by how gluon radiated, understand from central cut diagrams

$$\mathcal{N}_{g}^{\mathsf{qLPM}} = \frac{1}{N_{c}} (\cdots) \left[ 1 - \cos(\frac{y_{1}^{-} - y_{0}^{-}}{\tau_{gf}}) \right] \qquad \qquad \tau_{qf} = \frac{2q^{-}z(1-z)}{[\vec{l}_{\perp} - (1-z)\vec{v}_{\perp}]^{2}}$$

$$\mathcal{N}_{g}^{\mathsf{gLPM}} = C_{A}(\cdots) \left[ 1 - \cos(\frac{y_{1}^{-} - y_{0}^{-}}{\tau_{gf}}) \right] \qquad \qquad \tau_{gf} = \frac{2q^{-}z(1-z)}{[\vec{l}_{\perp} - (1-z)\vec{v}_{\perp} - \vec{k}_{\perp}]^{2}}$$

$$\mathcal{N}_g^{\mathsf{nonLPM}} = C_F(\cdots) \qquad \qquad \mathsf{no LP}$$

Yuan-Yuan Zhang

Single & Double Scattering Dijet

#### PM interference





### A simple model for gluon saturation in $\phi(x_G, k_{\perp}) \sim \hat{q}(k_{\perp})$

Simple model to include saturation

$$\phi_N(x_G, k_{\perp}, \mu^2) = \begin{cases} \phi_N^0 \text{ at } Q_s, & k_{\perp} < Q_s; \\ \\ \phi_N^0(\mu^2 = k_{\perp}^2), & k_{\perp} > Q_s, \end{cases}$$
 TMDlib Package  
F Hautmann et al., EPJC 74 (2014)

Calculate saturation scale self-consistently

$$Q_s^2(x_B, Q^2, b_{\perp}) = C \int dy_0^- \rho(y_0^-, b_{\perp}) \int \frac{d^2 k_{\perp}}{(2\pi)^2} \alpha_s(\mu) d\mu$$

Scale in  $\alpha_s(\mu)$  and  $\phi_N(x_G, k_\perp, \mu^2)$  is  $\mu^2 = k_\perp^2$ 

Yuan-Yuan Zhang

Single & Double Scattering Dijet

 $\phi_N(x_G, k_{\perp}, \mu^2)$ 







### $\hat{q}_F$ from our simple saturation model

From definition of saturation scale, relation between quark qhat & gluon qhat

$$Q_s^2 = \int dy_0^- \hat{q}_A(y_0^-, b_\perp, k_\perp), \, \hat{q}_F = \frac{4}{9} \hat{q}_A$$

For kinematics:  $x_R = 0.1 - 0.4, Q^2 = 2 - 6 \text{ GeV}^2$ 

Suppression of leading hadron method give  $\hat{q}_F^0 \approx 0.02 - 0.03$  GeV<sup>2</sup>/fm NB Chang et al, PRC 89.3 (2014): 034911

Single & Double Scattering Dijet

Yuan-Yuan Zhang

Our simple model give  $\hat{q}_F^0 \approx 0.014 - 0.025$  GeV<sup>2</sup>/fm

pT broadening of final-state hadron method give  $\hat{q}_F^0 \approx 0.015$  GeV<sup>2</sup>/fm P. Ru et al, arXiv:2004.00027



## **Nuclear Modification Ratio**

Ratio of dijet cross section in e+A and e+p

$$R_{eA}^{S(D)}(l_{\perp}, l_{q\perp}, \Delta \phi, z) = \frac{d\hat{\overline{\sigma}}_{eA}^{S(D)}}{d\mathscr{P}} / A \frac{d\hat{\overline{\sigma}}_{eA}}{d\mathscr{P}} / A \frac{d\hat{\overline{\sigma}}$$

Don't distinguish quark jet from gluon jet  $\hat{\overline{\sigma}} \equiv \hat{\sigma}(l_{\perp}, l_{a\perp}, \Delta\phi, z) + \hat{\sigma}(l_{a\perp}, l_{\perp}, \Delta\phi, 1 - z)$ 

Kinematics for calculation:

 $E_{\rho} = 10 \text{ GeV}, E_N = 100 \text{ GeV}, x_R = 0.2, Q^2 = 200 \text{ GeV}^2, A = 208$ 

Kinematic constraints from approximations, experiments

Yuan-Yuan Zhang





# Dijet Spectrum

 $d\sigma_{e\mathrm{A}}^{S(D)}$  $dx_R dQ^2 dz d^2 l_\perp d^2 l_{q\perp}$ 



• Dijet spectra from e+p and e+A scattering peak at  $\Delta \phi = \pi$ 

• Double scattering dijet Xsection small, can be negative due to non-LPM term

Yuan-Yuan Zhang









Nuclear Modification of Dijet

Yuan-Yuan Zhang

 $|\vec{l}_{\perp} + \vec{l}_{q\perp}| = 2l_{\perp}^2(1 + \cos\Delta\phi)$ for  $l_{\perp} = l_{a\perp}$ 

nonLPM: finite(relative contribution increase with z increase)

#### qLPM: negligible(suppress by color and LPM factor)

gLPM: dominant



11

# Azimuthal angle $\Delta \phi$ dependence : $R_{eA}^{S}$ & $R_{eA}^{S+D}$



 $Q_{\rm s} = 1.2 \,\,{\rm GeV}$ 

• Single scattering dominates in dijet Xsection,  $R_{eA}^{S}$  dominate in  $R_{eA}^{S+D}$ 

•  $Q_s$  artificially increase, peak in  $R_{eA}^S$  moves,  $R_{eA}^D$  contribution increase

Yuan-Yuan Zhang

Nuclear Modification of Dijet



 $Q_{\rm s} = 2 \,\,{\rm GeV}$ 

 $\Delta \phi$ 



### **Rapidity gap** $|y_{l_a} - y_l|$ dependence



o  $R_{\rho A}^{S}$  only due to quark pT broadening, independent of rapidity gap  $|y_{l_{a}} - y_{l}|$ 

o  $R_{eA}^D$  dominant - gLPM term with LPM suppression factor  $1 - \cos \frac{y_{10}}{2}$  $au_{gf}$  $|y_{l_a} - y_l| \uparrow$ ,  $z \downarrow$ ,  $\tau_{gf} \downarrow$ , when  $2R_A/\tau_{gf} \ge 2\pi$ , LPM suppression disappear, increased incoherent contribution Yuan-Yuan Zhang











### Nuclear size $R_A$ dependence



•  $R_{\rho A}^{S}$  linear in  $R_{A}$ 

Yuan-Yuan Zhang

Nuclear Modification of Dijet



14

Summary



- $\bullet$
- Nuclear modification in single and double scattering depend on  $\phi(x_G, k_{\perp})$
- LPM effect and gluon saturation embedded in  $\phi(x_G, k_{\perp})$  bring unique features in  $\Delta \phi$ ,  $|y_{l_a} - y_l|$ ,  $R_A$  dependence of nuclear modification ratio

Yuan-Yuan Zhang



Use dijet correlation in e+A to probe gluon TMD pdf  $\phi(x_G, k_1)$  or TMD  $\hat{q}(k_1)$ 



### Thanks for your attention!