

Diquarks as a QCD Breakthrough into Nuclear Physics (via Short-Range Correlated Nucleon Pairs)

*Jennifer Rittenhouse West
Lawrence Berkeley National Lab
EIC Early Career Workshop 2022*



Overview: Fundamental QCD effects in nuclei

- Fundamental QCD degrees of freedom: color-charged quarks, gluons, SU(3)
- fQCD underlies all of nuclear physics but often unnecessary to descend to that level - Effective field theory sufficient
- Experimental puzzle: 1983 EMC effect, mysterious quark behavior in nuclei \implies possible fQCD effects on nuclear scales
- Diquark & Hexadiquark solutions proposed to affect structure functions F_2 - short-range QCD physics appears in nuclei





Image credit: Brookhaven National Lab

EMC effect: Deep Inelastic Scattering

- Lepton scatters from target, exchanging virtual photon with 4-momentum q^2 given by: $Q^2 \equiv -q^2 = 2EE'(1 - \cos \theta)$
- γ^* strikes quark: We know the fraction of nucleon momentum carried by the struck quark via the Bjorken scaling variable $x_B = \frac{Q^2}{2M_p\nu}$ where $\nu = E - E'$, M_p =mass of proton, lepton masses neglected
- EMC plots: Ratio of structure functions vs. momentum fraction carried by struck quark x_B



Adapted from *Nuclear & Particle Physics* by B.R. Martin, 2003

Differential cross section for DIS:

$$\frac{d\sigma}{dx dy} (e^- p \rightarrow e^- X) = \sum_f x e_f^2 \left[q_f(x) + \bar{q}_f(x) \right] \cdot \frac{2\pi\alpha^2 s}{Q^4} (1 + (1 - y)^2)$$

where $y = \frac{\nu}{E}$ is the fraction of ℓ^- energy transferred to the target. $F_2(x)$ is the **nucleon structure function**, defined as:



$$F_2(x_B) \equiv \sum_f x_B e_f^2 \left(q_f(x_B) + \bar{q}_f(x_B) \right)$$

in terms of quark distribution functions $q_f(x)$: probability to find a quark with momentum $x_i \in [x, x + dx]$.

EMC effect: Distortion of nuclear structure functions

Plotting ratio of $F_2(x_B) \equiv \sum_f x_B e_f^2 (q_f(x_B) + \bar{q}_f(x_B))$ vs. x_B

- Predicted $F_2(x_B)$ ratio in complete disagreement with theory
- Why should quark behavior - confined in nucleons at QCD energy scales ~ 200 MeV - be so affected when nucleons embedded in nuclei, $BE \geq 2.2$ MeV?
- Mystery has not been solved to this day.



“THE RATIO OF THE NUCLEON STRUCTURE FUNCTIONS F_2^N FOR IRON AND DEUTERIUM “
The European Muon Collaboration, J.J. AUBERT et al. 1983

EMC effect experiments & explanations

POSSIBLE EXPLANATIONS

- Mean field effects involving the whole nucleus
- Local effects, e.g., 2-nucleon correlations

Simple mean field effects inconsistent with the EMC effect in light nuclei - MC of ${}^9\text{Be}$ \implies clustering
Seely *et al.*, 2009.

“This one new bit of information has reinvigorated the experimental and theoretical efforts to pin down the underlying cause of the EMC effect.” Malace *et al.*, 2014


Short-range N-N correlated pairs (SRC) may cause EMC effect (first suggested in *Ciofi & Liuti 1990, 1991*).
Neutron-proton pairs later found to dominate SRC
(CLAS collaboration & others)

DOZENS OF EXPERIMENTS

CONFIRM EMC EFFECT

Target	Collaboration/ Laboratory
${}^3\text{He}$	JLab HERMES
${}^4\text{He}$	JLab SLAC NMC
${}^6\text{Li}$	NMC
${}^9\text{Be}$	JLab SLAC NMC
${}^{12}\text{C}$	JLab SLAC NMC EMC
${}^{14}\text{N}$	HERMES BCDMS
${}^{27}\text{Al}$	Rochester-SLAC-MIT SLAC NMC
${}^{40}\text{Ca}$	SLAC NMC EMC
${}^{56}\text{Fe}$	Rochester-SLAC-MIT SLAC NMC BCDMS
${}^{64}\text{Cu}$	EMC
${}^{108}\text{Ag}$	SLAC
${}^{119}\text{Sn}$	NMC EMC
${}^{197}\text{Au}$	SLAC
${}^{207}\text{Pb}$	NMC


New model: Diquark formation proposed to create short-range correlations (SRC), modifying quark behavior in the NN pair



Malace, Gaskell, Higinbotham & Cloet,
Int.J.Mod.Phys.E 23 (2014)

Overview: Fundamental QCD dynamics in NN pairs

New model: Diquark formation proposed to create short-range correlations (SRC), modifying quark behavior in the NN pair



Short-range QCD potentials act on distance scales < 1 fm. Strong NN overlap can bring valence quarks within range.

What are SRC?

Short-range correlated nucleon-nucleon pairs

- Nuclei consist of protons and neutrons
~80% of which are organized into shells/
LRC
- Nuclear shell model is a “description of nuclei of atoms by analogy with the Bohr atomic model of electron energy levels.
- It was developed independently in the late 1940s by the American physicist **Maria Goeppert Mayer** and the German physicist **J. Hans D. Jensen**, who shared the Nobel Prize for Physics in 1963 for their work.”
- *William L. Hosch, www.britannica.com*


~20% of nucleons are in short-range correlated pairs - not shells/LRC

- SRC have very high relative momentum - nearly all nucleons above the Fermi momentum of the nucleus, $k_F \sim 250 \text{ MeV}/c$, are in SRC

Guide to the Nuclear Wallchart*
*You don't need to be a Nuclear Physicist to understand Nuclear Science.

The Shell Model

One such model is the Shell Model, which accounts for many features of the nuclear energy levels. According to this model, the motion of each nucleon is governed by the average attractive force of all the other nucleons. The resulting orbits form "shells," just as the orbits of electrons in atoms do. As nucleons are added to the nucleus, they drop into the lowest-energy shells permitted by the Pauli Principle which requires that each nucleon have a unique set of quantum numbers to describe its motion




When a shell is full (that is, when the nucleons have used up all of the possible sets of quantum number assignments), a nucleus of unusual stability forms. This concept is similar to that found in an atom where a filled set of electron quantum numbers results in an atom with unusual stability—an inert gas. When all the protons or neutrons in a nucleus are in filled shells, the number of protons or neutrons is called a "magic number." Some of the magic numbers are 2, 8, 20, 28, 50, 82, and 126. For example, ^{116}Sn has a magic number of protons (50) and ^{54}Fe has a magic number of neutrons (28). Some nuclei, for example ^{40}Ca and ^{208}Pb , have magic numbers of both protons and neutrons; these nuclei have exceptional stability and are called "doubly magic." Magic numbers are indicated on the chart of the nuclides.

www2.lbl.gov/abc/wallchart/chapters/06/1.html

What is a diquark?

- Strong force described by special unitary group $SU(3)_C$, local symmetry of the strong interaction \equiv QCD
- QCD \implies Diquark creation: Quark-quark bond with single gluon exchange & group theory transformation into a fundamentally different object:

$$3_C \otimes 3_C \rightarrow \bar{3}_C$$



Like quarks and gluons, diquarks carry color charge. They cannot be seen directly due to color confinement. Only 1_C (red+green+blue or red-antired etc.) directly detected.

Therefore there is no direct evidence for diquarks. Work in progress for diquark detection experimental proposals (e.g., diquark jets from DIS increase Λ production)

Strong indirect evidence exists (baryon mass splittings, Regge slopes).

What are diquark-induced short-range correlations (SRC)?


Diquark-induced SRC

What causes the “short-range” part of short-range NN correlations?

- Quantum fluctuations in separation distance between 2 nucleons
- *or*
- Quantum fluctuations in relative momentum between 2 nucleons

What causes the “correlation” in SRC?

- Diquark forms across nucleons
- Valence quarks from different nucleons “fall into” short-range QCD potential $V(r)$
- Highly energetically favorable $[ud]$ diquark created, a spin-0, isospin-0 qq combination



Why spin-0 $[ud]$ diquark formation?

There are 4 options for diquarks created out of valence quarks in the proton and neutron:

- Spin-0, Isospin-0 $[ud]$
- Spin-1, Isospin-1 (ud)
- Spin-1, Isospin-1 (uu)
- Spin-1, Isospin-1 (dd)

The scalar $[ud]$ is lower in mass by nearly 200 MeV.

What about a spin-0, isospin-1 $[ud]'$? Doesn't work due to spin-statistics constraints on the diquark wave function:

$$\Psi_{[ud]'} \propto \psi_{\text{color}} \psi_{\text{spin}} \psi_{\text{iso}} \psi_{\text{space}}$$

\uparrow
 Antisymmetric


\uparrow
 Symmetric, L=0

What are the requirements for a diquark to form in the nuclear environment?
(not inside of a single baryon)

Diquark formation across N-N pairs


Requirements for diquark induced SRC:

1. Nucleon-Nucleon wavefunctions must **STRONGLY** overlap
2. Attractive short-range QCD potential between valence quarks
3. Significant binding energy for diquark to form (much stronger than nuclear binding energies - comparable to confinement scale)




1. SRC 3D-overlap for relative momenta 400 MeV/c & 800 MeV/c

- **Plot 1:** According to the ^{12}C measurements from 2021 CLAS, NN tensor force dominates at 400 MeV/c relative momenta. Natural unit conversion gives 0.49 fm = 400 MeV/c.



- **Plot 2:** Tensor-scalar transition momenta - according to the ^{12}C measurements from 2021 CLAS, NN scalar force is in effect at 800 MeV/c relative momenta. Natural unit conversion gives 0.25 fm = 800 MeV/c.



2. Quark-quark potential in QCD: $V(r)$ calculation

- The $SU(3)_C$ invariant QCD Lagrangian:

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4}F^{\mu\nu a}F_{\mu\nu}^a + \bar{\Psi}_f \left(i\gamma^\mu D_\mu - m \right) \Psi_f$$

where covariant derivative $D_\mu = \partial_\mu - ig_s A_\mu^a t^a$ acts on quark fields, t^a are the 3x3 traceless Hermitian matrices (e.g. the 8 Gell-Mann matrices), g_s the strong interaction coupling, $\alpha_s \equiv \frac{g_s^2}{4\pi}$.

- QCD potential for states in representations R and R' is given by:

$$V(r) = \frac{g_s^2}{4\pi r} t_R^a \otimes t_{R'}^a$$

- To compute $V(r)$ for a $3_c \otimes 3_c \rightarrow \bar{3}_c$, we use the definition of the scalar $C_2(R)$, $t_R^a t_R^a \equiv C_2(R) \mathbf{1}$, the *quadratic Casimir operator* (NB: R_f is the final state representation):

$$V(r) = \frac{g_s^2}{4\pi r} \cdot \frac{1}{2} \cdot \left(C_2(R_f) - C_2(R) - C_2(R') \right)$$


- Diquarks combine 2 fundamental representation quarks into an anti-fundamental, $3_C \otimes 3_c \rightarrow \bar{3}_C$:

$$V(r) = -\frac{2}{3} \frac{g_s^2}{4\pi r} \implies \text{Diquark is bound!}$$



$$q\bar{q} : V(r) = -\frac{4}{3} \frac{g_s^2}{4\pi r}$$

Diquark induced N-N correlation:



Compare to color singlet attractive potential:

3. Diquark binding energy: Color hyperfine structure

Use Λ^0 baryon to find binding energy of $[ud]$:

$$\text{B.E.}_{[ud]} = m_u^b + m_d^b + m_s^b - M_{\Lambda^0}$$

Spin-spin interaction contribute to hadron mass;
QCD hyperfine interactions:

$$1. M_{(\text{baryon})} = \sum_{i=1}^3 m_i + a' \sum_{i<j} (\sigma_i \cdot \sigma_j) / m_i m_j$$

$$2. M_{(\text{meson})} = m_1 + m_2 + a (\sigma_1 \cdot \sigma_2) / m_1 m_2$$

(de Rujula, Georgi & Glashow 1975, Gasirowicz & Rosner 1981, Karliner & Rosner 2014)

Effective masses of light quarks are found using Eq.1 and fitting to measured baryon masses:

$$m_u^b = m_d^b \equiv m_q^b = 363 \text{ MeV}, \quad m_s^b = 538 \text{ MeV}$$

$$\text{B.E.}_{[ud]} = m_u^b + m_d^b + m_s^b - M_{\Lambda} = 148 \pm 9 \text{ MeV}$$

Relevant diquark-carrying baryons: Λ , Σ^+ , Σ^0 , Σ^-




TABLE I: Diquark properties

Diquark	Binding Energy (MeV)	Mass (MeV)	Isospin I	Spin S
$[ud]$	148 ± 9	578 ± 11	0	0
(ud)	0	776 ± 11	1	1
(uu)	0	776 ± 11	1	1
(dd)	0	776 ± 11	1	1

Uncertainties calculated using average light quark mass errors
 $\Delta m_q = 5 \text{ MeV}$ [37]

TABLE II: Relevant $SU(3)_C$ hyperfine structure baryons [28]

Baryon	Diquark-Quark content	Mass (MeV)	$I (J^P)$
Λ	$[ud]s$	1115.683 ± 0.006	$0 \left(\frac{1}{2}^+ \right)$
Σ^+	$(uu)s$	1189.37 ± 0.07	$1 \left(\frac{1}{2}^+ \right)$
Σ^0	$(ud)s$	1192.642 ± 0.024	$1 \left(\frac{1}{2}^+ \right)$
Σ^-	$(dd)s$	1197.449 ± 0.030	$1 \left(\frac{1}{2}^+ \right)$

$I (J^P)$ denotes the usual isospin I , total spin J and parity P quantum numbers, all have $L=0$ therefore $J = S$

“Diquark Induced Nucleon-Nucleon Correlations and the EMC Effect,”
JRW, arXiv:2009.06968

Diquark formation across N-N pairs

Requirements for diquark induced SRC:

1. Nucleon-Nucleon wavefunctions must **STRONGLY** overlap
2. Attractive short-range QCD potential between valence quarks
3. Significant binding energy for diquark to form (much stronger than nuclear binding energies - comparable to confinement scale)



What are the implications of NN diquark formation?
Quark flavor dependence of low mass [*ud*] will affect the np vs. pp SRC!

Diquark formation prediction for A=3 SRC: Isospin

Nucleon wavefunction : $|N\rangle = \alpha |qqq\rangle + \beta |q[qq]\rangle$

Scalar [ud] diquark formation for nucleons with 3-valence quark internal structure

$|N\rangle \propto |qqq\rangle$:

$${}^3H : 2n + p \rightarrow 4u, 5d \implies np \supset [ud] \times 10 \implies 60\% n-p, 40\% n-n$$

$$\implies nn \supset [ud] \times 4$$

$${}^3He : 2p + n \rightarrow 5u, 4d \implies np \supset [ud] \times 10 \implies 60\% n-p, 40\% p-p$$

$$\implies pp \supset [ud] \times 4$$

Scalar diquark formation for nucleons in quark-diquark internal configuration $|N\rangle \propto |q[qq]\rangle$:

$${}^3H : u [ud] + u [ud] + d [ud] \implies 100\% n-p$$

$${}^3He : d [ud] + d [ud] + u [ud] \implies 100\% n-p$$

The number of possible diquark combinations in A = 3 nuclei with nucleons in the 3-valence quark configuration is found by simple counting arguments. First, the 9 quarks of 3He with nucleon location indices are written as:

$$\begin{aligned} N_1 : p &\supset u_{11} u_{12} d_{13} \\ N_2 : p &\supset u_{21} u_{22} d_{23} \\ N_3 : n &\supset u_{31} d_{32} d_{33} \end{aligned} \quad (21)$$

where the first index of q_{ij} labels which of the 3 nucleons the quark belongs to, and the second index indicates which of the 3 valence quarks it is. Diquark induced SRC requires the first index of the quarks in the diquark to differ, $[u_{ij}d_{kl}]$ with $i \neq k$. The 4 possible combinations from $p-p$ SRC are listed below.

$$u_{11}d_{23} \quad u_{12}d_{23} \quad (22)$$

$$u_{21}d_{13} \quad u_{22}d_{13} \quad (23)$$

Short-range correlations from $n-p$ pairs have 10 possible combinations,

$$\begin{aligned} u_{11}d_{32} \quad u_{12}d_{32} \\ u_{11}d_{33} \quad u_{12}d_{33} \\ u_{21}d_{32} \quad u_{22}d_{32} \\ u_{21}d_{33} \quad u_{22}d_{33} \\ u_{31}d_{13} \quad u_{31}d_{23} \end{aligned} \quad (24)$$

which gives the number of $p-p$ combinations to $n-p$ combinations in this case as $\frac{2}{5}$.

Combining these results yields the following inequality for the isospin dependence of N-N SRC:

$${}^3He : 0 \leq \frac{\mathcal{N}_{pp}}{\mathcal{N}_{np}} \leq \frac{2}{5} \quad (25)$$

where \mathcal{N}_{NN} is the number of SRC between the nucleon flavors in the subscript.

The same argument may be made for 3H due to the quark-level isospin-0 interaction, to find

$${}^3H : 0 \leq \frac{\mathcal{N}_{nn}}{\mathcal{N}_{np}} \leq \frac{2}{5}. \quad (26)$$

JRW, arXiv:2009.06968

Combine into isospin dependent SRC ratio predictions :

$${}^3He : 0 \leq \frac{N_{pp \text{ SRC}}}{N_{np \text{ SRC}}} \leq \frac{2}{5}, \quad {}^3H : 0 \leq \frac{N_{nn \text{ SRC}}}{N_{np \text{ SRC}}} \leq \frac{2}{5}, \quad \text{Maximum 40\%!}$$

Diquark formation induced SRC inequality tentatively confirmed: JLab experiment E12-11-112 A=3 mirror nuclei results

Preliminary results from JLab: $\frac{\mathcal{N}_{nn}}{\mathcal{N}_{np}} = \frac{1}{4.23} \sim 0.24$

Individual nucleon wavefunctions at lowest order are dominated by two Fock states with unknown coefficients; the 3 valence quark configuration and the quark-diquark configuration,

$$|N\rangle = \alpha|qqq\rangle + \beta|q[qq]\rangle, \quad (27)$$

where square brackets indicate the spin-0 $[ud]$ diquark. The full A=3 nuclear wavefunction is given by

$$|\Psi_{A=3}\rangle \propto (\alpha|qqq\rangle + \beta|q[qq]\rangle)(\alpha|qqq\rangle + \beta|q[qq]\rangle) \quad (28)$$

$$(\gamma|qqq\rangle + \delta|q[qq]\rangle)$$

where the proton and the neutron are allowed to have different weights for each valence quark configuration. This expands out to

$$|\Psi_{A=3}\rangle \propto \alpha^2\gamma|qqq\rangle^3 + 2\alpha\beta\gamma|qqq\rangle^2|q[qq]\rangle \quad (29)$$

$$\alpha^2\delta|qqq\rangle^2|q[qq]\rangle + \beta^2\gamma|qqq\rangle|q[qq]\rangle^2 +$$

$$2\alpha\beta\delta|qqq\rangle|q[qq]\rangle^2 + \beta^2\delta|q[qq]\rangle^3,$$

with mixed terms demonstrating that it is not straightforward to map the $\frac{\mathcal{N}_{pp}}{\mathcal{N}_{np}}$ ratio to precise coefficients for each nucleon's Fock states. A perhaps reasonable simplification is to assume that the proton and the neutron have the same coefficients for their 2-body and 3-body valence states, i.e. to set $\gamma = \alpha$ and $\delta = \beta$ in Eq. 28. In this case, the nuclear wavefunction reduces to

$$|\Psi_{A=3}\rangle \propto \alpha^3|qqq\rangle^3 + 3\alpha^2\beta|qqq\rangle^2|q[qq]\rangle \quad (30)$$

$$+ 3\beta^2\alpha|qqq\rangle|q[qq]\rangle^2 + \beta^3|q[qq]\rangle^3.$$

JRW, arXiv:2009.06968

Isospin dependent SRC ratio inequalities from diquark induced SRC :

$${}^3\text{He} : \quad 0 \leq \frac{N_{pp \text{ SRC}}}{N_{np \text{ SRC}}} \leq 0.4$$

$${}^3\text{H} : \quad 0 \leq \frac{N_{nn \text{ SRC}}}{N_{np \text{ SRC}}} \leq 0.4$$

⇒ Nucleon wavefunction MAY contain both $|qqq\rangle$ and $|q[ud]\rangle$
with approximately equal coefficients

⇒ Diquark formation – single gluon exchange & $SU(3)_C$ transformation –
favored over quark and diquark exchange.

Nuclear structure functions $F_2(x_B)$ from the diquark model?

Likely wrong: Diquark formation modification of F_2 from Fermi motion of quarks in 2-nucleon SRC

Recall quark momentum distribution functions $q(x_B)$: $F_2(x_B) \equiv \sum_f x_B e_f^2 \left(q_f(x_B) + \bar{q}_f(x_B) \right)$

Fermi energy : $E_F = \frac{p_F^2}{2m}$

Fermi momentum : $p_F = \sqrt{2mE_F} \propto m^{\frac{1}{2}}$

- Diquarks lower the mass of the system
- Effective masses of quarks in nucleons:
 $m_u = m_d = 363 \text{ MeV}$
- $[ud]$ diquark mass: $m_{[ud]} = 578 \text{ MeV}$
- Therefore each quark loses 75 MeV and its Fermi

momentum is depleted: $m_{\text{final}} = \sqrt{m_q - \frac{BE}{2}}$

Momentum ratio of quark in diquark to free quark :

$$\frac{p_{\text{final}}}{p_{\text{initial}}} = \sqrt{\frac{m_f}{m_i}} \approx 0.89$$

Diquark structures in nuclei: X17 anomaly

New work on arXiv →

Effects of Hexadiquark Fock state in ${}^4\text{He}$ nuclear wavefunction

Based on “QCD hidden-color hexadiquark in the core of nuclei,” *jrw, Brodsky, de Teramond, Goldhaber & Schmidt*

Nucl.Phys.A 1007 (2021), 2004.14659

jennifer@lbl.gov



Quantum Chromodynamics Resolution of the ATOMKI Anomaly in ${}^4\text{He}$ Nuclear Transitions

Valery Kubarovsky,¹ Jennifer Rittenhouse West,^{2,3} and Stanley J. Brodsky⁴

¹Thomas Jefferson National Accelerator Laboratory, Newport News, VA 23606, USA

²Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

³EIC Center at Thomas Jefferson National Accelerator Laboratory, Newport News, VA 23606, USA

⁴SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94309, USA

(Dated: Wednesday 29th June, 2022)

Recent observations of the angular correlation spectra in the decays ${}^4\text{He}^* \rightarrow {}^4\text{He} + e^+e^-$ and ${}^8\text{Be}^* \rightarrow {}^8\text{Be} + e^+e^-$ have been suggested as due to the creation and subsequent decay to an electron-positron pair of a new light particle with a mass of ~ 17 MeV. In this work, we present a calculation of the invariant mass $m_{e^+e^-}$ spectrum of the electromagnetic transition of an excited state of helium and estimate the differential and total width of the decay. We investigate the possibility that the source of the signal is an e^+e^- pair created by a new electromagnetic decay of ${}^4\text{He}$ caused by a proposed 12-quark hidden-color Fock state in the ${}^4\text{He}$ nuclear wavefunction, the “hexadiquark.” We find that we can fit the shape of the signal with the QCD Fock state at excitation energy $E^* \simeq 17.9$ MeV and a Gaussian form factor for the electromagnetic decay. We address the physical issues with the fit parameters using properties of the hexadiquark state. In light of this work, we emphasize the need for independent experimental confirmation or refutation of the ATOMKI results as well as further experiments to detect the proposed new excitation of ${}^4\text{He}$.

Introduction

Observations by the ATOMKI collaboration of anomalous angular correlations in electron-positron pairs produced in the nuclear decays ${}^4\text{He}^* \rightarrow {}^4\text{He} + e^+e^-$ [1, 2] and ${}^8\text{Be}^* \rightarrow {}^8\text{Be} + e^+e^-$ [3] have been attributed to the creation and subsequent decay of a new light particle to an e^+e^- pair of a new light particle with mass of ~ 17 MeV, dubbed the X17 or simply X. Recently, the same group reported observations of the lepton pair in the off-resonance region of ${}^7\text{Li}(p, e^+e^-){}^8\text{Be}$ direct proton-capture reactions [4]. Our work in this article will focus on the ${}^4\text{He}$ experiment in which the observed invariant mass $m_{e^+e^-}$ of the lepton pair was found to be $m_X = 16.94 \pm 0.12(\text{stat.}) \pm 0.21(\text{syst.})$ MeV. The Feynman diagram for this transition is shown in Fig. 1. The signal has generated a great deal of theoretical interest in both the particle and nuclear physics communities [5–25].

The light-front Fock state expansion of QCD has led to new perspectives for the nonperturbative eigenstructure of hadrons, including the quark-antiquark structure of mesons, the quark-diquark structure of baryons (such as the $|u[ud]\rangle$ composition of the proton) and the diquark-antidiquark structure of tetraquarks. In the case of nuclear physics, the color-singlet $|[ud][ud][ud][ud][ud][ud]\rangle$ “hexadiquark” Fock state has the same quantum numbers as the ${}^4\text{He}$ nucleus. The existence of the hexadiquark Fock state in the eigensolution of the ${}^4\text{He}$ nuclear eigenstate [27] can explain the anomalously large binding energy of the α particle. QCD also predicts orbital and radial excitations of the hexadiquark and thus novel excitations of ${}^4\text{He}$ beyond the standard excitations predicted by nucleonic degrees of freedom. The excitation energy of the hexadiquark can be below the energy required to produce hadronic decays such as $p + {}^3\text{H}$ and thus have evaded detection.

iv:submit/4378584 [hep-ph] 29 Jun 2022

arXiv:2206.14659

such as $p + {}^3\text{H}$ and thus have evaded detection. The excitation energy of the hexadiquark can be below the energy required to produce hadronic decays the hexadiquark and thus novel excitations of ${}^4\text{He}$ beyond the standard excitations predicted by nucleonic degrees explain the anomalously large binding energy of the α particle. QCD also predicts orbital and radial excitations of ${}^4\text{He}$ nuclei. The existence of the hexadiquark Fock state in the eigensolution of the ${}^4\text{He}$ nuclear eigenstate [27] can explain the anomalously large binding energy of the α particle. QCD also predicts orbital and radial excitations of the hexadiquark and thus novel excitations of ${}^4\text{He}$ beyond the standard excitations predicted by nucleonic degrees of freedom. The excitation energy of the hexadiquark can be below the energy required to produce hadronic decays such as $p + {}^3\text{H}$ and thus have evaded detection.

Summary

- Diquark formation proposed to cause short-range correlated nucleon pairs
- Diquark-induced SRC may be a viable explanation for the EMC effect


- Direct calculations of F_2 ideal; Complete $F_2 \forall$ nuclear targets missing for 39 years:

$$F_2(x) \equiv \sum_f x_B Q_f^2 \left(q_f(x_B) + \bar{q}_f(x_B) \right)$$

- Approach the problem obliquely & look for indirect evidence of NN diquarks like the SRC flavor inequality

- Aim: Calculate the ~20% SRC from QCD - this needs both imaging and dynamics of nucleons

- *Pipe dream: Shell model from QCD*



Fin

Jennifer Rittenhouse West
Berkeley Lab & EIC Center @JLab
EIC Early Career Workshop #2!
25 July 2022



backup slides

Nuclear physics: Diquark formation should account for
pieces of the known NN potential

Argonne v_{18} nucleon-nucleon potential: $V = \sum_{n=1}^{18} V_n(r) O_n$

51

ACCURATE NUCLEON-NUCLEON




FIG. 7. Tensor and tensor-isospin parts of the potential. Also shown are the OPE contribution to the tensor-isospin potential, and for comparison an OPE potential with a monopole form factor containing a 900 MeV cutoff mass.

Wiringa, Stoks, Schiavilla 1995




FIG. 6. Central, isospin, spin, and spin-isospin components of the potential. The central potential has a peak value of 2031 MeV at $r = 0$.

The Argonne V18 potential has the form

$$V = \sum_{n=1}^{18} V_n(r) O_n \quad (1.1)$$

where $V_n(r)$ are rotationally-invariant coefficient functions of the relative coordinate of the nucleons and the O_n are the eighteen spin-isospin operators given in Table 1.,

Table 1: Argonne V18 spin-isospin operators in coordinate-space

Term	spin-isospin Operator in r-space
O_1	\mathbf{I}
O_2	$(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_3	$(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)$
O_4	$(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_5	$S_{12} = 3(\boldsymbol{\sigma}_1 \cdot \hat{\mathbf{r}})(\boldsymbol{\sigma}_2 \cdot \hat{\mathbf{r}}) - \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2$
O_6	$S_{12}(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_7	$(\mathbf{L} \cdot \mathbf{S})$
O_8	$(\mathbf{L} \cdot \mathbf{S})(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_9	$(\mathbf{L} \cdot \mathbf{L})$
O_{10}	$(\mathbf{L} \cdot \mathbf{L})(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_{11}	$(\mathbf{L} \cdot \mathbf{L})(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)$
O_{12}	$(\mathbf{L} \cdot \mathbf{L})(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_{13}	$(\mathbf{L} \cdot \mathbf{S})^2$
O_{14}	$(\mathbf{L} \cdot \mathbf{S})^2(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2)$
O_{15}	$T_{12} = (3\tau_{1z}\tau_{2z} - \boldsymbol{\tau} \cdot \boldsymbol{\tau})$
O_{16}	$(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)T_{12}$
O_{17}	$S_{12}T_{12}$
O_{18}	$(\tau_{1z} + \tau_{2z})$

S. Veerasamy and W. N. Polzou 2011


“The NN potential is written as a sum of an electromagnetic (EM) part, a one-pion-exchange (OPE) part, and an intermediate- and short-range phenomenological part.”

$$v(NN) = v^{\text{EM}}(NN) + v^{\pi}(NN) + v^R(NN)$$

The short and intermediate range pieces are a sum of central + tensor + (others, L^2 , spin-orbit, quadratic spin-orbit) \implies diquark potentials within tensor piece

Argonne v_{18} nucleon-nucleon potential: $V = \sum_{n=1}^{18} V_n(r) O_n$

\exists a dip in tensor $V(r)$ at $r = 0.1$ fm




Wiringa, Stoks, Schiavilla 1995

- $V(0.1) \approx -10$ MeV
- QCD short-range potentials ~ 200 MeV
- The scale problem is here too!
- Solution: In a strongly overlapping N-N pair, the diquark $V(r)$ is not the only potential acting in the system - linear combination of QCD potentials just like the nuclear V_{total}
- Valence quark q_1 from N_1 , fluctuating into a short distance from valence quark q_2 from N_2 , senses the diquark potential as well as its "home" color singlet potential:

$$V(r_{q_1-q_2}) = -\frac{2}{3} \frac{g_s^2}{4\pi r_{12}} \text{ vs. } V(r_{q_1-dq_1}) = -\frac{4}{3} \frac{g_s^2}{4\pi r_{11}}$$

AV18: Argonne v_{18} nucleon-nucleon potential



McKenna et al. 2011

- N-N system - nucleons in quark-diquark configuration - analogous to the hydrogen molecule
- Complicated problem. Proof of principle only.

2D 2-body approximation. Set V to 10 MeV, solve for separation distances:

$$V_{\text{total}} = -\frac{2}{3} \frac{g_s^2}{4\pi r_{12}} - \frac{4}{3} \frac{g_s^2}{4\pi r_{11}} = -10 \text{ MeV}$$

- For separation distance between quarks in the newly formed diquark:

$$r_{q_1-q_2} = 0.1 \text{ fm}$$

- Solving for separation distance between q_1 and $[u_1 d_1]$:

$$r_{q_1-dq_1} \approx 0.65 \text{ fm}$$

Reasonable. Work in progress.

