Disentangling Long and Short Distances in Momentum-Space TMDs

Zhiquan Sun Markus Ebert, Johannes Michel, Iain Stewart arXiv: 2201.07237

Correlation in Partonic and Hadronic Interactions (CPHI) March 10, 2022 Duke University



Massachusetts Institute of Technology





Outline

• Motivation

$$\int d^2 \vec{k}_T f^{\text{TMD}}(x, k_T) \stackrel{?}{=} f^{\text{coll}}(x)$$

- Challenging to move between position and momentum space
- Challenging to separate perturbative and NP physics in TMD models

Mathematical Formalism

- Truncated functionals: model-independent perturbative baseline
- Boundary terms: systematic corrections from NP region

Applications

- Compare cumulative TMDPDF to collinear PDF
- Constrain OPE coefficients

Conclusions

Cumulative TMDPDF

• What's the k_T -integral of the TMDPDFs?

$$\int d^2 \vec{k}_T \ f^{\text{TMD}}(x, k_T, \mu, \zeta) \stackrel{?}{=} f^{\text{coll}}(x, \mu)$$

$$\uparrow$$
naively yes :)

Verified at bare level at 1-loop: *Echevarria, Idilbi, & Scimemi (1111.4996)*



Renormalization breaks the naive expectation

$$\mu \frac{d}{d\mu} \int d^2 \vec{k}_T f^{\text{TMD}}(x, k_T, \mu, \zeta) \neq \mu \frac{d}{d\mu} f^{\text{coll}}(x, \mu)$$

renormalization says no :(

Observed to hold for certain NP models at NLO (also for g_1, h_1): Bacchetta & Prokudin (1303.2129)

TMD Data

Drell-Yan cross section:



$$\frac{d\sigma}{dQdYd^2q_T} = H(Q,\mu) \sum_i \int d^2 \vec{b}_T \ e^{i\vec{q}_T \cdot \vec{b}_T} f_i(x_a, b_T, \mu, \zeta_a) \ f_{\bar{i}}(x_b, b_T, \mu, \zeta_b) \times \left[1 + \mathcal{O}(\frac{q_T^2}{Q^2})\right]$$

• Measurements are done in momentum space (q_T)

CMS: 1909.04133 ATLAS: 1912.02844



900

TMD Factorization

• TMD factorization of Drell-Yan:

$$\frac{d\sigma}{dQdYd^2q_T} = H(Q,\mu) \sum_{i} \int d^2 \vec{b}_T \ e^{i\vec{q}_T \cdot \vec{b}_T} f_i(x_a, b_T, \mu, \zeta_a) \ f_{\vec{i}}(x_b, b_T, \mu, \zeta_b) \times \left[1 + \mathcal{O}(\frac{q_T^2}{Q^2})\right]$$
Hard virtual corrections
Hard virtual momentum of the partons

- Factorization is most easily written in position space (b_T)
 - μ = Renormalization scale

 Q, q_{7}

 ζ = Collins-Soper parameter

p



TMDPDFs have both perturbative and nonperturbative parts, and usually:

$$f_i(x, b_T, \mu, \zeta) = f_{\text{pert, }i}(x, b^*(b_T), \mu, \zeta) \cdot f_{\text{NP}}(x, b_T, \zeta)$$
Calculated with expansion in $\alpha_s(1/b_T)$ Intrinsic + NP evolution

 The perturbative part can be computed by matching onto collinear PDFs:

$$\begin{split} f_{\text{pert, }i}^{\text{TMD}}(x, b_T, \mu, \zeta) &= \sum_j \int_x^1 \frac{dz}{z} C_{ij}(\frac{x}{z}, b_T, \mu, \zeta) f_j^{\text{coll}}(z, \mu) \\ &= f_i^{\text{coll}}(x, \mu) + \alpha_s C_{ij}^{(1)} \otimes f_j^{\text{coll}}(x, \mu) + \mathcal{O}(\alpha_s^2) \end{split}$$

• The NP part looks like: $f_{\text{NP}}(x, b_T, \zeta) = 1 + b_T^2 \left(\Lambda_i^{*(2)}(x) + \gamma_{\zeta,i}^{*(2)} L_{\zeta} \right) + \mathcal{O}(b_T^4)$

TMDPDFs have both perturbative and nonperturbative parts, and usually:

$$f_i(x, b_T, \mu, \zeta) = f_{\text{pert, }i}(x, b^*(b_T), \mu, \zeta) \cdot f_{\text{NP}}(x, b_T, \zeta)$$



Collins & Soper (Nucl. Phys. B 197 (1982) 446) Bacchetta et al (1703.10157) Scimemi & Vladimirov (1803.11089)

- $b^*(b_T)$ shields the Landau pole
- $b_T \ll 1/\Lambda_{\text{QCD}}$: $b^*(b_T) \to b_T, f_{\text{NP}} \to 1$

 $f_{\rm pert}$ dominates

• $b_T \gg 1/\Lambda_{\text{QCD}}$: $b^*(b_T) \rightarrow \text{constant}$

$f_{\rm NP}$ dominates

- Different models of $f_{\rm NP}$ are used for fitting to data
- $b^*(b_T)$ shields the Landau pole and is <u>coupled</u> to f_{NP} $f_{TMD}(x, b_T, \mu, \zeta) = f_{pert}(x, b_A^*(b_T), \mu, \zeta) \cdot f_{NP}^A(x, b_T, \zeta)$ $= f_{pert}(x, b_B^*(b_T), \mu, \zeta) \cdot f_{NP}^B(x, b_T, \zeta)$

 $b_A^*(b_T) \neq b_B^*(b_T) \Rightarrow f_{NP}^A(x, b_T) \text{ and } f_{NP}^B(x, b_T) \text{ are not comparable!}$

• The perturbative and nonperturbative effects are mixed up!

 b^* known to be incompatible with leading renormalon in TMD PDF and γ_{ζ} : *Scimemi & Vladimirov (1609.06047)*

use either $b_{CS}^*(b_T)$ or $b_{Pavia}^*(b_T)$:

- b^* prescriptions makes different $f_{\rm NP}$ not comparable
- For example, take the same $f_{NP}(b_T) = e^{-(0.5 \text{GeV } b_T)^2}$,

0.08 $f_{
m NP} = \exp[-\Lambda^2 b_T^2]$ $b_{\rm CS}^*(b_T) = \frac{b_T}{\sqrt{1 + (b_T/b_{\rm max})^2}} = b_T \left(1 + \mathcal{O}(b_T^2)\right)$ $b_{
m max}=1.123~{
m GeV}^{-1}$ $egin{aligned} \Lambda &= 0.5 \; ext{GeV} \ ---f_{ ext{pert}}^{ ext{LL}}(b_{ ext{CS}}^*) \cdot f_{ ext{NP}} \ ---f_{ ext{pert}}^{ ext{LL}}(b_{ ext{Pavia}}^*) \cdot f_{ ext{NP}} \end{aligned}$ 0.06 $\Lambda = 0.5 \text{ GeV}$ $b_T f_{
m pert}^{
m LL}(Q,b_T)$ Collins & Soper (Nucl. Phys. B 197 (1982) 446) $b_{\text{Pavia}}^{*}(b_{T}) = b_{\max}\left(1 - \exp(-\frac{b_{T}^{4}}{b^{4}})\right)^{\frac{1}{4}} = b_{T}\left(1 + \mathcal{O}(b_{T}^{4})\right)$ 0.02 Bacchetta et al (1703.10157) 0.00L 1 2 3 $b_T ~[{
m GeV}^{-1}]$

• Goal: extract nonperturbative physics without b^* contamination

Momentum Space

• Measurements are in q_T space: Fourier transform

$$\frac{d\sigma}{dq_T} = 2\pi q_T \int_0^\infty \frac{d^2 \vec{b}_T}{(2\pi)^2} e^{i\vec{q}_T \cdot \vec{b}_T} \sigma(b_T)$$

$$q_T \operatorname{spectrum} = q_T \int_0^\infty db_T \ b_T \int_0^{2\pi} \frac{d\phi}{2\pi} \ e^{iq_T b_T \cos\phi} \sigma(b_T) = q_T \int_0^\infty db_T \ b_T \ J_0(q_T b_T) \sigma(b_T)$$

- For perturbative q_T , integral still includes nonperturbative b_T !
- Intuition: perturbative q_T should be dominated by perturbative $b_T \sim 1/q_T$

Momentum Space

- Intuition: perturbative q_T should be dominated by perturbative b_T
- Goal: make this intuition manifest



Truncated Functionals

- Want to approximate S[f] using <u>perturbative</u> $b_T \leq b_T^{\text{cut}}$
- Can use $S_{\leq}[f]$, but need to systematically account for $S_{\geq}[f]$

$$S_{>}[f](q_T, b_T^{\text{cut}}) = q_T \int_{b_T^{\text{cut}}}^{\infty} db_T \ b_T J_0(q_T b_T) f(b_T)$$

Assumption:

a) $|f(b_T \to \infty)| < b_T^{-\rho},$ $|f'(b_T \to \infty)| < b_T^{-\rho-1}, \rho > \frac{1}{2}$

b) $f(b_T)$ differentiable at $b_T^{\rm cut}$



$$= -b_T^{\text{cut}} J_1(q_T b_T^{\text{cut}}) f(b_T^{\text{cut}}) - \int_{b_T^{\text{cut}}}^{\infty} db_T \ b_T J_1(q_T b_T) f'(b_T)$$
asymptotic form
$$q_T \gg 1/b_T^{\text{cut}} > \Lambda_{\text{QCD}}$$

$$= \sqrt{\frac{2b_T^{\text{cut}}}{\pi q_T}} \cos\left(q_T b_T^{\text{cut}} + \frac{\pi}{4}\right) \ f(b_T^{\text{cut}}) + \mathcal{O}[(b_T^{\text{cut}} q_T)^{-\frac{3}{2}}]$$

$$J_0(x \to \infty) = \sqrt{\frac{2}{\pi x}} \cos(x - \frac{\pi}{4}) + \mathcal{O}(x^{-\frac{3}{2}})$$
$$J_1(x \to \infty) = -\sqrt{\frac{2}{\pi x}} \cos(x + \frac{\pi}{4}) + \mathcal{O}(x^{-\frac{3}{2}})$$

Truncated Functionals

Perturbative region

• Define a systematic series to approximate S[f] using $b_T \leq b_T^{cut}$

Leading order:
$$S^{(0)}[f](q_T) \equiv S_{<}[f](q_T) = q_T \int_0^{b_T^{\text{cut}}} db_T \ b_T J_0(q_T b_T) f(b_T)$$

• Define $S^{(1)}[f]$ to include leading boundary contribution from $S_{>}[f]$

$$S^{(1)}[f](q_T) \equiv S^{(0)}[f] + \sqrt{\frac{2b_T^{\text{cut}}}{\pi q_T}} \cos\left(q_T b_T^{\text{cut}} + \frac{\pi}{4}\right) f(b_T^{\text{cut}}) \quad \longleftarrow \text{First correction!}$$

$$S[f](q_T) = S^{(1)}[f](q_T, b_T^{\text{cut}}) + \frac{1}{q_T} \mathcal{O}[(b_T^{\text{cut}}q_T)^{-\frac{1}{2}}]$$

Truncated Functionals

 $b_T^{\text{cut}} \xrightarrow{\text{nonperturbative}}$ perturbative Systematically add on power corrections J_{TMD} so $S^{(n)}[f] \to S[f]$ $S^{(0)}[f](q_T, b_T^{\text{cut}}) = \int_0^{b_T^{\text{cut}}} db_T \ b_T J_0(q_T b_T) f(b_T),$ $S^{(1)}[f](q_T, b_T^{\text{cut}}) = S^{(0)}[f] + \sqrt{\frac{2b_T^{\text{cut}}}{\pi q_T}} f(b_T^{\text{cut}}) \cdot \cos(b_T^{\text{cut}} q_T + \frac{\pi}{4})$ $1/\Lambda_{\text{QCD}} \ \vec{b_{\tau}}$ $S^{(2)}[f](q_T, b_T^{\text{cut}}) = S^{(1)}[f] - \sqrt{\frac{2b_T^{\text{cut}}}{\pi q_T}} \left(\frac{3 f(b_T^{\text{cut}})}{8 b_T^{\text{cut}} q_T} + \frac{f'(b_T^{\text{cut}})}{q_T}\right) \cdot \cos(b_T^{\text{cut}} q_T - \frac{\pi}{4})$ $S^{(3)}[f](q_T, b_T^{\text{cut}}) = S^{(2)}[f] + \sqrt{\frac{2b_T^{\text{cut}}}{\pi q_T}} \left(\frac{15 \ f(b_T^{\text{cut}})}{128 \ b_T^{\text{cut}^2} q_T^2} - \frac{7 \ f'(b_T^{\text{cut}})}{8 \ b_T^{\text{cut}} q_T^2} - \frac{f''(b_T^{\text{cut}})}{q_T^2}\right) \cdot \cos(b_T^{\text{cut}} q_T + \frac{\pi}{4})$

$$S[f](q_T) = S^{(n)}[f] + \frac{1}{q_T} \mathcal{O}[(b_T^{\text{cut}}q_T)^{-n+\frac{1}{2}}]$$

Power Correction to Functionals

- Toy function $f = \exp[-\lambda_1 \ln^2(b_T Q)] \exp[-\lambda_2 b_T^2]$
- Errors of truncated functionals follow expected power law $S[f](q_T) = S^{(n)}[f] + \frac{1}{a_T} \mathcal{O}[(b_T^{\text{cut}}q_T)^{-n+\frac{1}{2}}]$



Perturbative Input

- Power expand toy function and use only "perturbative" input $f^{(0)}$ $f = \exp[-\lambda_1 \ln^2(b_T Q)](1 - \lambda_2 b_T^2 + \mathcal{O}(b_T^4))$ $f^{(0)}$
- "Errors" of truncated functionals identify missing quadratic term



Cumulative Functionals

- Investigate $\int d^2 \vec{k}_T f^{\text{TMD}}(x, k_T, \mu, \zeta) \stackrel{?}{=} f^{\text{coll}}(x, \mu)$
- <u>Cumulative distribution</u> with a UV cutoff:

$$\int_{|k_{T}| \le k_{T}^{\text{cut}}} d^{2}\vec{k}_{T} f(k_{T}) = \int_{|k_{T}| \le k_{T}^{\text{cut}}} d^{2}\vec{k}_{T} \int \frac{d^{2}b_{T}}{(2\pi)^{2}} e^{+i\vec{k}_{T}\cdot\vec{b}_{T}} f(b_{T})$$

$$= \int^{k_{T}^{\text{cut}}} dk_{T} k_{T} \int_{0}^{\infty} db_{T} b_{T} J_{0}(b_{T}k_{T}) f(b_{T})$$

$$= k_{T}^{\text{cut}} \int_{0}^{\infty} db_{T} J_{1}(b_{T}k_{T}^{\text{cut}}) f(b_{T})$$

$$\underbrace{K[f](k_{T}^{\text{cut}})}$$

• Approximate using perturbative region:

$$K^{(0)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) = k_T^{\text{cut}} \int_0^{b_T^{\text{cut}}} db_T J_1(b_T k_T^{\text{cut}}) f(b_T)$$

Cumulative Functionals

• Systematically add on power corrections so $K^{(n)}[f] \rightarrow K[f]$

$$\begin{split} K^{(0)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) &= k_T^{\text{cut}} \int_0^{b_T^{\text{cut}}} db_T J_1(b_T k_T^{\text{cut}}) f(b_T) \\ K^{(1)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) &= K^{(0)}[f] + f(b_T^{\text{cut}}) \cdot \frac{\cos(b_T^{\text{cut}} k_T^{\text{cut}} - \frac{\pi}{4})}{\sqrt{\frac{\pi}{2}} (b_T^{\text{cut}} k_T^{\text{cut}})^{1/2}} \\ K^{(2)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) &= K^{(1)}[f] - \left(\frac{f(b_T^{\text{cut}})}{8 b_T^{\text{cut}} k_T^{\text{cut}}} - \frac{f'(b_T^{\text{cut}})}{k_T^{\text{cut}}}\right) \cdot \frac{\cos(b_T^{\text{cut}} k_T^{\text{cut}} + \frac{\pi}{4})}{\sqrt{\frac{\pi}{2}} (b_T^{\text{cut}} k_T^{\text{cut}})^{1/2}} \\ K^{(3)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) &= K^{(2)}[f] - \left(\frac{9f(b_T^{\text{cut}})}{128 b_T^{\text{cut}^2} k_T^{\text{cut}^2}} - \frac{5f'(b_T^{\text{cut}})}{8 b_T^{\text{cut}} k_T^{\text{cut}^2}} + \frac{f''(b_T^{\text{cut}})}{k_T^{\text{cut}^2}}\right) \cdot \frac{\cos(b_T^{\text{cut}} k_T^{\text{cut}} - \frac{\pi}{4})}{\sqrt{\frac{\pi}{2}} (b_T^{\text{cut}} k_T^{\text{cut}})^{1/2}} \end{split}$$

$$K[f](k_T^{\text{cut}}) = K^{(n)}[f](k_T^{\text{cut}}, b_T^{\text{cut}}) + \mathcal{O}[(b_T^{\text{cut}}k_T^{\text{cut}})^{-n-\frac{1}{2}}]$$

Perturbative Baseline

• Leading NP correction for TMDPDF:

$$f_{i}(x, b_{T}, \mu, \zeta) = \frac{f_{i}^{(0)}(x, b_{T}, \mu, \zeta)}{\int} \left\{ 1 + b_{T}^{2} \left(\Lambda_{i}^{(2)}(x) + \frac{1}{2} \gamma_{\zeta, i}^{(2)} L_{\zeta} \right) \right\} + \mathcal{O} \left[(\Lambda_{\text{QCD}} b_{T})^{4} \right]$$
Perturbative, use matching Intrinsic From NP evolution computed to N3LL

 $L_{\zeta} = \ln(b_T^2 \zeta / b_0^2)$

• Pair with the K[f] functionals to get the TMDPDF cumulant

$$\int^{k_T^{\text{cut}}} d^2 \vec{k}_T f_i(k_T) = K^{(3)}[f_i^{(0)}](k_T^{\text{cut}}) + \Lambda_i^{(2)}K^{(3)}[b_T^2 f_i^{(0)}] + \frac{1}{2}\gamma_{\zeta,i}^{(2)}K^{(3)}[b_T^2 L_\zeta f_i^{(0)}] + \mathcal{O}\Big[\left(k_T^{\text{cut}}b_T^{\text{cut}}\right)^{-\frac{7}{2}}, \left(\frac{\Lambda_{\text{QCD}}}{k_T^{\text{cut}}}\right)^4\Big]$$

$$+ \mathcal{O}\Big[\left(k_T^{\text{cut}}b_T^{\text{cut}}\right)^{-\frac{7}{2}}, \left(\frac{\Lambda_{\text{QCD}}}{k_T^{\text{cut}}}\right)^4\Big]$$

$$\int_{Truncating functional} \text{Truncating OPE}$$

Cumulant of TMDPDFs

- Approximate the cumulant using $K^{(3)}[f^{(0)}_{TMD}]$ and normalize to f^{coll}
- Deviation is small! $\int_{0}^{k_T^{\text{cut}}} d^2 \vec{k}_T f^{\text{TMD}}(x, k_T, \mu = \sqrt{\zeta} = k_T^{\text{cut}}) = f^{\text{coll}}(x, \mu = k_T^{\text{cut}}) \text{ YES!}$



- Model-independent confirmation of previous results at $\mu = k_T^{cut}$ Bacchetta & Prokudin (1303.2129)
- Central value consistent with intuitive result within percents
- $\Delta_{\rm res}$ is perturbative uncertainties estimated by scale variations

Cumulant of TMDPDFs

• Deviation is small!

$$\int d^2 \vec{k}_T f^{\text{TMD}}(x, k_T, \mu = \sqrt{\zeta} = k_T^{\text{cut}}) = f^{\text{coll}}(x, \mu = k_T^{\text{cut}}) \quad \textbf{YES!}$$

• NP uncertainty much smaller than perturbative uncertainty



Cumulant of TMDPDFs

• Deviation is small!

$$d^{2}\vec{k}_{T} f^{\text{TMD}}(x,k_{T},\mu=\sqrt{\zeta}=k_{T}^{\text{cut}})=f^{\text{coll}}(x,\mu=k_{T}^{\text{cut}}) \quad \textbf{YES!}$$

• Test the observation as a function of x and keep k_T^{cut} fixed



- Central value can differ from zero by 2%
- Small deviation supported by modelbased global fits

Scimemi & Vladimirov (1912.06532) Bacchetta et al: 1912.07550

 \boldsymbol{x}

Impact of Evolution Effects



- Intuitive expectation is robust in the vicinity of $\mu = \sqrt{\zeta} = k_T^{\rm cut}$
- For $\mu = k_T^{\rm cut}$, the ζ evolution is negligible*
- Sizable corrections from evolution away from these regions, due to the cusp anomalous dimension
- Evolution effect matters, but at the natural scale $\mu = k_T^{\rm cut}$ the intuition is valid

 $d^{2}\vec{k}_{T} f^{\text{TMD}}(x,k_{T},\mu=k_{T}^{\text{cut}},\zeta) = f^{\text{coll}}(x,\mu=k_{T}^{\text{cut}})$

Conclusions

- Perturbative and nonperturbative physics in TMDPDFs are usually hard to disentangle because of b^* prescriptions
- Systematically improvable truncated functionals let us directly compute linear impact of $\mathcal{O}(b_T^2)$ NP parameters without b* prescription
 - Model-independent estimate of NP uncertainties on integral of TMD PDF
 - Can also apply to cross section: extract NP parameters in linearized fit to experimental data in the future, complementary to model-based global fits (no time to cover — ask me if you want to know more!)
- Demonstrate that integrating the unpolarized TMDPDF over $[0, k_T^{\text{cut}}]$ gives the collinear PDF to the percent level (when renormalization scale $\mu = k_T^{\text{cut}}$)!

Thank you!

Back-up Slides

Drell-Yan Cross Section

• Leading NP correction for cross section in b_T space:

$$\sigma(b_T) = \sigma^{(0)}(b_T) \left\{ 1 + b_T^2 \left(2\overline{\Lambda}^{(2)} + \gamma^{(2)}_{\zeta,q} L_{Q^2} \right) \right\} + \mathcal{O} \left[(\Lambda_{\text{QCD}} b_T)^4 \right]$$
Perturbative Intrinsic to TMDPDF From evolution From evolution

• Pair with the S[f] functionals to get momentum space spectrum

Truncating functional Truncating OPE

 $\Rightarrow \begin{array}{l} \textbf{Useful for estimating uncertainties or} \\ \textbf{putting model-independent constraints on } \gamma^{(2)}_{\zeta,q} \text{ and } \overline{\Lambda}^{(2)} \end{array}$

Interpret as Uncertainty

- $\Delta_{
 m cut}$ is from varying the choice of $b_T^{
 m cut}$ in $\int_0^{\circ_T} db_T$
- Evaluating impact from each NP parameter individually
- Compare with SV and Pavia global fits



Constraints on NP Parameters

- Solid line: mock data generated using b^* and $f_{\rm NP}$
- Dashed line: signal template using our functionals
- Linear signal, easy to scale





30





32



More on NP effects



Evolution effects + perturbative

