

## Conclusions

- Neither M-scaling nor F-scaling in the particle generation particularly narrow  $p_T$  bins.
- Intermittency and hence scale invariant fluctuations not present.
- For wide  $p_T$  bins, F-scaling observed with  $\nu \sim 1.7-1.9 > 1.304$ , the theoretical value predicted by GL theory for second-order phase transition.
- Scaling exponent is independent of centrality cut for wide  $p_T$  bins.
- Angantyr overestimates the value of scaling exponent compared with ALICE data for Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV.

## Intermittency analysis of charged hadrons generated in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and 5.02 TeV using PYTHIA8/Angantyr

Salman K. Malik and Dr. Ramni Gupta

Department of Physics, University of Jammu, J&K, India

### Introduction

- At  $\mu_B > 0$  (baryonic chemical-potential), experimental approach to study phase-diagram is via event-by-event fluctuations.
- Non-monotonous increase in fluctuations near phase transition and critical point.
- At critical point, correlation length increases rapidly, and the system becomes scale-invariant.
- Scale invariance is given by the behaviour of moments  $\rightarrow$  **Normalized Factorial Moments** (NFM) in our case.

### Intermittency


NFMs are given by: [Bialas:1988](#), [Bialas:1985](#):

$$F_q(M) = \frac{\frac{1}{N} \sum_{e=1}^N \frac{1}{M} \sum_{m=1}^M f_q(n_{me})}{\left( \frac{1}{N} \sum_{e=1}^N \frac{1}{M} \sum_{m=1}^M f_1(n_{me}) \right)^q}$$

where,  $f_q(n_{me}) = \prod_{j=0}^{q-1} (n_{me} - j)$   
 $q$  (order of the moment)  $\geq 2$

Scaling of  $F_q$  with the number of bins  $M$ :

- $F_q(M) \propto (M^D)^{\phi_q} \rightarrow$  **M-scaling**
- Scaling of different orders of factorial moments with the decreasing number of bins  $M$  is called *intermittency*



M-scaling depends on different critical parameters of the system than **F-scaling**, given as:

- $F_q(M) \propto F_2(M)^{\beta_q}$ , where
- $\beta_q \propto (q-1)^\nu$

$\nu$ , **Scaling exponent** is independent of the critical parameters of the system. Predictions:

$\rightarrow$  Theoretical predictions: 1.304 in GL theory for second-order PT [Hwa:1992](#)

$\rightarrow$  1.0 in 2D Ising model [Hwa:1992](#).

### PYTHIA8/Angantyr

- Extrapolates pp dynamics, to heavy ion collisions, retaining as much as possible from pp.
- It does not assume a hot thermalised medium and is developed with the motivation that differences between the model and experimental results may show some effects of collective behaviour.
- Angantyr gives a good description of general final state properties, in p-Pb and Pb-Pb, Xe-Xe collisions. [Bierlich:2018xfw](#).
- Intermittency analysis and more specifically, the value of  $\nu$  (Scaling exponent) is already calculated with **AMPT**, **EPOS3** and in a recent **QM 2022 poster** for ALICE data.








Fig: Charged particle pseudorapidity density distribution compared with ALICE and ATLAS data points.

### Observations


Intermittency analysis has been performed for various  $p_T$  bins ( $0.4 \leq p_T \leq 1.0$ ,  $0.4 \leq p_T \leq 0.6$ ,  $0.6 \leq p_T \leq 0.8$ ,  $0.8 \leq p_T \leq 1.0$ ) for  $\sqrt{s_{NN}} = 2.76$  & 5.02 TeV with different centralities and  $|\eta| < 0.8$ . **2.76 TeV**  $\sim 2M$  events, 0-5 % centrality



### 5.02 TeV $\sim 1M$ events, 0-5 % centrality



Figures show M-scaling for various  $p_T$  bins for two different energies and F-scaling,  $\nu$  for  $0.4 \leq p_T \leq 1.0$  GeV/c. For narrow width  $p_T$  bins, no F-scaling is observed, and hence  $\nu$  is not calculated:



**HUGS 2022**

Email: [salman.khurshid.malik@cern.ch](mailto:salman.khurshid.malik@cern.ch)