

Charge Symmetry Violation in valence quark distribution extraction


via. Precise Measurement of $\frac{\pi^+}{\pi^-}$ Ratios in SIDIS

Shuo Jia

shuojia@temple.edu

Temple University & Argonne National Lab

June 16, 2022




Motivation

Charge symmetry Violation

Charge symmetry (CS) is one special kind of **isospin symmetry**.


$$\delta d(x) = d^p(x) - u^n(x),$$

$$\delta u(x) = u^p(x) - d^n(x).$$



Lattice limits

CSV limits from lattice calculation:



arXiv:1512.04139v1

Experimental Limits

CSV(x) contains $\delta d - \delta u$, where

$$\delta d(x) = d^p(x) - u^n(x), \delta u(x) = u^p(x) - d^n(x).$$

Theoretical limits

Model by Sather:

$$\delta d(x) \sim 2 - 3\%, \delta u(x) \sim 1\%$$

E. Sather, Phys. Lett. B274, 433 (1992)

Model by Rodionov, Thomas and Londergan
 $\delta d(x)$ could reach up to 10% at high x

E. N. Rodionov, A. W. Thomas and J. T. Londergan, Mod. Phys. Lett. A 9, 1799 (1994)

Phenomenological limits

CSV parameterization


$$\delta u_v = \delta d_v = k(1-x)^4 x^{0.5} (x - 0.0909)$$

The form has to satisfy the normalization condition

$$\int dx \delta uv(x) = \int dx \delta dv(x) = 0$$

k was varied in the global fit: 90% CL

Eur. Phys. J.35(2004)325



Formalism

Semi-inclusive Deep Inelastic Scattering

Charge symmetry Violation

$$\delta d(x) = d^p(x) - u^n(x), \delta u(x) = u^p(x) - d^n(x).$$

$$CSV(x) = \delta d - \delta u$$

Londergan, Pang and Thomas PRD54(1996)3154

$$R_{meas}^D(x, z) = \frac{4N^{D\pi^-}(x, z) - N^{D\pi^+}(x, z)}{N^{D\pi^+}(x, z) - N^{D\pi^-}(x, z)} = \frac{4R_Y(x, z) - 1}{1 - R_Y(x, z)} \quad (1)$$

where $N^{D\pi^\pm}(x, z)$ is the yield of π^\pm electroproduction on a deuterium target

Factorization

$$N^{Nh} = \sum_i e_i^2 q_i^N(x) D_i^h(z)$$

Impulse Approximation

$$N^{D\pi^\pm}(x, z) = N^{p\pi^\pm}(x, z) + N^{n\pi^\pm}(x, z)$$

Formalism No. 1

Londergan, Pang and Thomas PRD54(1996)3154


$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$

$$R(x, z) = \frac{5}{2} + R_{meas}^D, A(x) = \frac{-4}{3(u_v + d_v)}, \text{known } B(x, z) = \frac{5}{2} + R_{sea-S}^D(x, z) + R_{sea-NS}^D(x), \text{known}$$

$$D(z) = \frac{1 - \Delta(z)}{1 + \Delta(z)}, \Delta(z) = D_u^{\pi^-}(z) / D_u^{\pi^+}(z), CSV(x) = (\delta d - \delta u)$$

1. Calculate $CSV(x)$ for each (Q^2, x) setting from different Fragmentation Functions
2. Extract simultaneously $D(z)$ and $CSV(x)$ from each (Q^2, x) setting

Experiment Overview




4 z measurements (0.4,0.5,0.6,0.7) for each x, Q^2 setting.

$$Q^2 = 4.0 \text{ GeV}^2, x = 0.35, 0.4, 0.45, 0.5$$

$$Q^2 = 4.75 \text{ GeV}^2, x = 0.45, 0.5, 0.55, 0.6$$

$$Q^2 = 5.5 \text{ GeV}^2, x = 0.5, 0.55, 0.6, 0.65$$

Electron: HMS, pion: SHMS
Fall 2018 and Spring 2019




- 10.6 GeV beam, LD₂(10 cm), LH₂(10 cm), Al-dummy
- HMS angle 13°-21°, 4.4-6.4 GeV, electrons
- SHMS angle 11°-21°, 1.7-4.5 GeV, π^+ / π^-

$W^2 > 4 \text{ GeV}^2$ and $W'^2 > 2.8 \text{ GeV}^2$ for SIDIS region

Data Analysis

$$\langle x \rangle = 0.5, \langle Q^2 \rangle = 5 \text{ GeV}^2, \langle z \rangle = 0.5$$



Data yield

$$Y_{corr}^D = \frac{N_{\text{pions}}}{Q \epsilon_t \epsilon_{LT} \epsilon_{PID}}$$

Radiative correction: $RC = \frac{Y_{SIMC,noradia}}{Y_{SIMS,radia}}$

Backgrounds from SIMC:

Y_{exc} : Exclusive radiative backgrounds

$$D(e, e' \pi^\pm) n(p) \gamma$$

Y_{delta} : Delta radiative backgrounds




$$D(e, e' p) \pi$$

Y_ρ : Diffractive ρ $D(e, e' \rho \rightarrow \pi^+ \pi^-)$

$$Y_D(x, z) = RC(Y_{corr}^D - 0.245 Y_{Dummy} - Y_{exc} - Y_{delta})$$

R_{meas}^D from data

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z), R(x, z) = \frac{5}{2} + R_{meas}^D(x, z)$$



$R_{meas}^D(x, z)$ after standard diffractive ρ backgrounds subtraction projected on z axis.



All variables are bin center corrected.

For each of (Q^2, x, z) , weighted average are taken for the overlap of the different group of runs

Formalism No.1

Model Dependence

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$



$$A(x) = \frac{-4}{3(u_v + d_v)}$$

$$B(x, z) = \frac{5}{2} + R_{sea_S}^D(x, z) + R_{sea_NS}^D(x)$$

Calculate CSV from formula No.1

Different Fragmentation Functions


$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$

$$CSV(x) = \frac{B(x, z) - D(z) R(x, z)}{A(x)}$$

$$D(z) = \frac{1 - \Delta(z)}{1 + \Delta(z)}, \Delta(z) = D_u^{\pi^-}(z) / D_u^{\pi^+}(z)$$

Fragmentation Functions:

- JAM20SIDIS
- fDSS LO
- fDSS NLO



Extract simultaneously from formula No.1

Fragmentation ratio and CSV extraction

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$

$$\Delta(z) \equiv \frac{D_u^{\pi^-}(z)}{D_u^{\pi^+}(z)} = z^\alpha (1-z)^\beta$$

$$CSV x \equiv \delta d - \delta u = x^a (1-x)^b (x-c)$$

constrain: $\int_0^1 CSV(x) dx = 0$


$$c = \frac{\int_0^1 x^{(a+1)} (1-x)^b}{\int_0^1 x^a (1-x)^b} = \frac{B(a+2, b+1)}{B(a+1, b+1)}, B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$$

$$R_{fit}^D(x, z) = \frac{B(x, z) - A(x) CSV(x)}{D(z)} - \frac{5}{2}$$

Diffractive ρ : $Y_{corr}^{D\pi^-} = Y^{D\pi^-} + \gamma Y_{\pi^-}^\rho$


Extract simultaneously from formula No.1

Results after standard ρ background subtraction



$$\Delta(z) \equiv D_u^{\pi^-}(z)/D_u^{\pi^+}(z) = z^\alpha(1-z)^\beta$$

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$



$$CSVx \equiv \delta d - \delta u = x^a(1-x)^b(x-c)$$

From the fitting result $\Delta(z)$, CSV can be calculated for each kinematic point. Weighted average are taken for overlap

New Formalism

Proposal equation

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$

$$CSV(x) = \delta d - \delta u$$

This equation used simple expansion.

New equation


Assume $T = \frac{\bar{u} + \bar{d}}{u + d}$, $S = \frac{2(s + \bar{s})}{u + d}$ and $D_s^+ = D_s^- = D_u^-$, $\delta d = -a\delta u$, $CSV = \delta d - \delta u$

$$Y = \frac{(4 + T + S)\Delta(z) + (4T + 1) + (4a\Delta(z) - 1)\delta u}{(4T + 1 + S)\Delta(z) + (4 + T) + (4a - \Delta(z))\delta u}$$

1. Calculate δu from existing FFs using the Yield ratio.
2. Fit the Yield ratio to extract simultaneously $\Delta(z)$ and δu

Calculate CSV from the new Formula

$$Y = \frac{(4 + T + S)\Delta(z) + (4T + 1) + (4a\Delta(z) - 1)\delta u}{(4T + 1 + S)\Delta(z) + (4 + T) + (4a - \Delta(z))\delta u}$$



Extract simultaneously from new formula

$$Y = \frac{(4 + T + S)\Delta(z) + (4T + 1) + (4a\Delta(z) - 1)\delta u}{(4T + 1 + S)\Delta(z) + (4 + T) + (4a - \Delta(z))\delta u}$$

$$\Delta(z) \equiv \frac{D_u^{\pi^-}(z)}{D_u^{\pi^+}(z)} = z^\alpha(1 - z)^\beta$$

$$\delta u(x) = kx^a(1 - x)^b(x - c)$$

In progress...

H_2 runs are added to constrain fragmentation functions



$$Y_{H_2} = \frac{(4u + \bar{d})\Delta(z) + (4\bar{u} + d) + (s + \bar{s})\Delta(z)}{(4u + \bar{d}) + (4\bar{u} + d)\Delta(z) + (s + \bar{s})\Delta(z)}$$

H₂ runs results

Factorization test from H₂ runs

$$\frac{\sigma_p^{\pi^+} - \sigma_p^{\pi^-}}{\sigma_d^{\pi^+} - \sigma_d^{\pi^-}} = \frac{4u + 4\bar{u} + d + \bar{d}}{5(u + \bar{u} + d + \bar{d})}$$



$$\frac{\sigma_p^{\pi^+} + \sigma_p^{\pi^-}}{\sigma_d^{\pi^+} + \sigma_d^{\pi^-}} = \frac{4u_v - d_v}{3(u_v + d_v)}$$



(predictions are from JAM20 PDFs)

Some other quantity by our D_2 runs

$$\frac{N_D^{\pi^+} - N_D^{\pi^-}}{N_D^{\pi^+} + N_D^{\pi^-}} = \frac{3(u_v + d_v)(D_u^+ - D_u^-)}{5(u + d + \bar{u} + \bar{d})(D_u^+ + D_u^-)}$$



(predictions are from JAM20 PDFs and FFs)

Future work

- Improve and update ρ background subtraction
- Systematic Uncertainty
- Finalize the analysis cuts and data set, cross check with Hem's ratio
- Fit with more data from PtSIDIS experiment

Acknowledgement

I would like to acknowledge my advisor Dr. Zein-Eddine Meziani. And thanks to Dr. Dipankar Dutta, Dr. Dave Gaskell, Dr. Peter Bosted and Dr. Whitney Armstrong for their help. Also I thank my collaborator Hem Bhatt.


This work is supported in part by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No DE-FG02-94ER4084 and US DOE contract DE-AC02-06CH11357.

Thank you!

Backups

Data Analysis

Kinematic Cut



4 z measurements (0.4,0.5,0.6,0.7)
for each x, Q^2 setting.

$$Q^2 = 4.0 GeV^2, x = 0.35, 0.4, 0.45, 0.5$$

$$Q^2 = 4.75 GeV^2, x = 0.45, 0.5, 0.55, 0.6$$

$$Q^2 = 5.5 GeV^2, x = 0.5, 0.55, 0.6, 0.65$$

DIS cut: Invariant mass squared $W^2 = (P + k - k')^2 = M^2 + \frac{1-x}{x} Q^2$


SIDIS cut: Mass of the unobserved final state squared

$$W'^2 = (P + k - k' - P_h)^2 = M^2 + Q^2 \frac{1-x}{x} + M_h^2 - 2 \cdot (z + 1) \frac{Q^2}{2 \cdot M x}$$

DIS: $W^2 > 4 GeV^2$ and SIDIS: $W'^2 > 2.8 GeV^2$

Model dependence for Formular No.1

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$




$$A(x) = \frac{-4}{3(u_v + d_v)}$$

$A(x)$ is calculated from Parton Distribution Function. Plot is calculated from JAM20 PDF.

Model dependence for Formular No.1

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$



cteq6 PDF and JAM20 FF

$$B(x, z) = \frac{5}{2} + R_{sea_S}^D(x, z) + R_{sea_NS}^D(x)$$

$$R_{sea_NS}^D = \frac{5(\bar{u}^p(x) + \bar{d}^p(x))}{[u_v^p(x) + d_v^p(x)]}$$

$$R_{sea_S}^D = \frac{\Delta_s(z)[s(x) + \bar{s}(x)]/(1 + \Delta(z))}{[u_v^p(x) + d_v^p(x)]}$$

$$\Delta_s(z) = \frac{D_s^-(z) + D_s^+(z)}{D_u^+(z)}$$

Diffraction ρ background subtraction

$$D(z) R(x, z) + A(x) CSV(x) = B(x, z)$$

Assumption:

- 1, The ρ decay into pions are charge symmetric: $N_{\pi^-}^\rho = N_{\pi^+}^\rho$
- 2. The ρ subtraction for π^+ runs and π^- runs are same

$$Y_{corr}^{D\pi^-} = Y^{D\pi^-} + \gamma Y_{\pi^-}^\rho$$

$$Y_{corr}^{D\pi^+} = Y^{D\pi^+} + \gamma Y_{\pi^-}^\rho$$

$$\delta R_\rho^D = \delta R_\rho^D(\gamma, Y_{\pi^-}^\rho, Y^{D\pi^+}, Y^{D\pi^-})$$

For standard ρ subtraction, γ is -1