


E07-007 & E08-025 Overview

Carlos Muñoz Camacho

Institut de Physique Nucléaire, Orsay, IN2P3/CNRS

DVCS Collaboration Meeting
December 19–20, 2013

Hall A DVCS program



Handbag diagram


High Q^2
Perturbative QCD

Non-perturbatif
GPDs

Limite de Bjorken :

$$Q^2 = \begin{cases} -q^2 & \rightarrow \infty \\ \nu & \rightarrow \infty \end{cases} \quad x_B = \frac{Q^2}{2M\nu} \text{ fixed}$$

Hall A DVCS program



High Q^2
Perturbative QCD

Non-perturbative
GPDs

Handbag diagram

PAC 39 Report, June 2012

"The new data will be of unprecedented statistical precision, but the statistics will go to waste if the data cannot be interpreted with commensurate accuracy. [One] example is the GPD program, where it is pointless to display new asymmetries before scaling has been established in a given channel"



E07-007/E08-025

Goal:

- Measure DVCS cross sections as a function of Q^2 and different beam energies, for both LH2 and LD2.
 - Separation of DVCS² from interference BH-DVCS interference terms
 - Rosenbluth separation of π^0 electroproduction cross section
-
- Same setup for both experiments
 - Only target change from LH2 to LD2
 - Data taken: Oct-Dec 2010

Motivation: results from E00-110


Q^2 variation of $\text{Im}\mathcal{C}^I$



BH much smaller than total cross section $\Rightarrow \text{BH} \cdot \text{DVCS}$
interference alone cannot explain the difference



E07-007: Rosenbluth-like DVCS²- \mathcal{I} separation in Hall A

- Clean separation of BH-DVCS interference term from pure DVCS²
- Scaling test on the real part of the DVCS amplitude
- Rosenbluth separation of σ_L/σ_T for $ep \rightarrow ep\pi^0$



DVCS detector package

- 208-channel PbF_2 electromagnetic calorimeter
- DVCS stand of top of BigBite stand
(moving cart: 1.1 m → 5.5 m from target)
- CH shielding in front of calorimeter



DVCS electronics and DAQ

1 GHz sampling based on the **Analog Ring Sampler (ARS)**


Chip developed by IRFU/CEA-Saclay

Front-end electronics by LPC/Clermont-Ferrand

Calorimeter trigger:

- **VALIDATES** if 1 set of 2×2 blocks over threshold (~ 500 ns).



1	2	3	4	5	6	7	8	9	10	11
12	13	14	15	16	17	18	19	20	21	22
23	24	25	26	27	28	29	30	31	32	33
34	35	36	37	38	39	40	41	42	43	44
45	46	47	48	49	50	51	52	53	54	55
56	57	58	59	60	61	62	63	64	65	66
67	68	69	70	71	72	73	74	75	76	77
78	79	80	81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96	97	98	99
100	101	102	103	104	105	106	107	108	109	110
111	112	113	114	115	116	117	118	119	120	121
122	123	124	125	126	127	128	129	130	131	132



$t(ns)$

- Each sample in 12-bit flash ADCs

DVCS setup in Hall A



Running conditions for E07-007 & E08-025

Data analysis

- Beam line
 - Polarimetry (Compton + Møller) data (**FINISHED**)
 - Beam charge monitors (BCM) calibration (**FINISHED**)
- HRS
 - Optics and mispointing checks (**FINISHED**)
 - Drift chambers time offsets (**FINISHED**)
- Calorimeter
 - Elastic (energy) calibrations (**FINISHED**)
 - Waveform analysis of PMT pulses (**FINISHED**)
 - Time (calorimeter + HRS) corrections (**FINISHED**)
 - Clustering and calibration optimizations (**FINISHED**)
- MC simulation
 - GEANT4 written from scratch based on previous GEANT3 (**FINISHED**)
 - Validating results by comparing to old simulation (**FINISHED**)

All basic detector calibrations + MC are finalized

TODO: Physics analysis

- E07-007: preliminary cross-section results available
(but cross-check needed!)
- E08-025: final checks on number of counts underway
(π^0 contamination, LH2 subtraction, etc)
- π^0 cross section...
- Rosenbluth separation (Charles' talk)