


# Vector Meson Domestic Zoo

- Some *vector mesons* can, compared to other mesons, be measured to very high precision.
- This stems from fact that *vector mesons* have **same** quantum numbers as *photon*.

$$I^G(J^{PC}) = 0^-(1^{--})$$



Name  
PDG  
particle data group

Quark  
Content  $\Gamma$   
(MeV)

$\rho^+(770)$   $u\bar{d}$  148

$\rho^0(770)$   $\frac{1}{\sqrt{2}}(u\bar{u} - d\bar{d})$  149

$\omega(782)$   $\frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d})$  8.5

$K^{*+}(892)$   $u\bar{s}$  51

$K^{*0}(892)$   $d\bar{s}$  47

$\phi(1020)$   $s\bar{s}$  4.3

$D^{*+}(2010)$   $c\bar{d}$  0.083

*Open Charm*

$D^{*0}(2007)$   $c\bar{u}$  < 2.1

$J/\psi(1S)(3097)$   $c\bar{c}$  0.093

*Charmonium*

$\psi'(2S)(3686)$   $c\bar{c}$  0.284

$Y(1S)(9460)$   $b\bar{b}$  0.052

*Quarkonium*



SLAC



- We will focus on **5 vector mesons** from  $\bar{q}q$  *Nonet* which **widths** are **narrow** enough to study *meson photoproduction* @ **threshold** & where data are available.



# Vector Meson – Nucleon SL

$$|\alpha_{Vp}| = B_V \cdot h_{Vp}$$




$$B_V^2 = \frac{\alpha \cdot m_V \cdot k}{12\pi \cdot \Gamma(V \rightarrow e^+e^-)}$$

$$h_{Vp} = \sqrt{b_1}$$



Jefferson Lab JLab6 JLab12 JLab20



IS, S. Prakhov, Ya. Azimov *et al*, Phys Rev C **91**, 045207 (2015)  
 IS, D. Epifanov, & L. Pentchev, Phys Rev C **101**, 042201 (2020)  
 IS, L. Pentchev, & A.I. Titov, Phys Rev C **101**, 045201 (2020)  
 Meng-Lin Du, V. Baru, Feng-Kun Guo, Ch. Hanhart, U.-G. Meissner,  
 A. Nefediev, & IS, Eur Phys J C **80**, 1053 (2020)  
 L. Pentchev & IS, Eur Phys J A **57**, 56 (2021)  
 IS, W.J. Briscoe, L. Pentchev, & A. Schmidt, Phys Rev C **104**, 074028 (2021)  
 Xiao-Yun Wang, Fancong Zeng, & IS, Phys Rev C **106**, 015202 (2022)




# Total Cross Sections for Vector Meson Photoproduction off Proton

- Traditionally,  $\sigma_t$  behavior of near-threshold binary *inelastic* reaction

$$m_a + M_b < m_c + M_d$$

is described as series of *odd* powers in  $q$  (*even* powers in case of *elastic*).

$$\sigma_t(q) = b_1 \cdot q + b_3 \cdot q^3 + b_5 \cdot q^5$$



- Our *assumption* is that there is no *VN bound state* below experimental  $q_{min}$ .

- Linear* term is determined by two independent *S*-waves only with total spin  $1/2$  &/or  $3/2$ .
- Contributions to *cubic* term come from both *P*-wave amplitudes & *W* dependence of *S*-wave amplitudes,
- Fifth-order* term arises from *D*-waves & *W* dependencies of *S*- & *P*-waves.

A2

$$b_1 = (4.42 \pm 0.14) \times 10^{-2} \mu\text{b}/(\text{MeV}/c)$$

IS, S. Prakhov, Ya. Azimov *et al*, Phys Rev C **91**, 045207 (2015)

clas

$$b_1 = (3.40 \pm 1.15) \times 10^{-4} \mu\text{b}/(\text{MeV}/c)$$

IS, L. Pentchev, & A.I. Titov, Phys Rev C **101**, 045201 (2020)

GLUEX

$$b_1 = (0.46 \pm 0.16) \times 10^{-6} \mu\text{b}/(\text{MeV}/c)$$

IS, D. Epifanov, & L. Pentchev, Phys Rev C **101**, 042201 (2020)

Quasidata

$$b_1 = (0.37 \pm 0.04) \times 10^{-9} \mu\text{b}/(\text{MeV}/c)$$

IS, W.J. Briscoe, L. Pentchev, & A. Schmidt, Phys Rev C **104**, 074028 (2021)

- Dramatic differences in hadronic factors

$$h_{Vp} = \sqrt{b_1}$$


as slopes ( $b_1$ ) of  $\sigma_t$  @ threshold as function of  $q$  varies significantly from  $\omega$  to  $\phi$  to  $J/\psi$ .

- Therefore, such big difference in *Scattering Length* is determined mainly by *hadronic factor*  $h_{Vp}$ .



# Vector Meson – Nucleon $SL$


- Most *theoretical* calculations using gluonic *van der Waals* interaction disagree with our *phenomenological* results. Specifically, they do not consider **VM young** effect



- Such big difference in  $SL$ s of  $Vp$  systems is determined mainly by hadronic factor  $h_{vp}$ , & reflects strong weakening of interaction in  $\bar{b}b-p$  &  $\bar{c}c-p$  systems compared to that of *light*  $\bar{q}q-p$  ( $q = u, d$ ) configurations.
- Interaction in  $\bar{s}s-p$  has intermediate strength that is manifested in intermediate value of  $\phi p$   $SL$ .

- Such small value of  $\phi p$   $SL$  compared to typical *hadron* size of **1 fm**, indicates that proton is more transparent for  $\phi$ -meson compared to  $\omega$ -meson, & is much less transparent than for  $J/\psi$ -meson.

$$|\alpha_{\psi p}| \ll |\alpha_{\psi' p}| < |\alpha_{J/\psi p}| \ll |\alpha_{\phi p}| \ll |\alpha_{\omega p}|$$



- $p \rightarrow V$  coupling  $\bar{q}q$  is proportional to  $\alpha_s$  & *separation* of corresponding quarks.
- This *separation* (in zero approximation) is proportional to  $1/m_V$ .



Courtesy of Misha Ryskin, July 2020



9/5/2022

JLUO Meeting on NSAC Long Range Plan, September 2022

Igor Strakovsky 4

