

What do lineshapes of resonances teach us about their nature?

Christoph Hanhart


Forschungszentrum Jülich

Based on

- F.-K. Guo et al. “Hadronic molecules,” *Rev. Mod. Phys.* 90(2018)015004
- I. Matuschek et al., “On the nature of near-threshold bound and virtual states,”
Eur. Phys. J. A 57 (2021) no.3, 101
- V. Baru et al., “Effective range expansion for narrow near-threshold resonances,”
[arXiv:2110.07484](https://arxiv.org/abs/2110.07484)
- C. Hanhart et al., “Lineshapes for composite particles with unstable constituents,”
PRD81 (2010), 094028


We want employ **line shapes** to distinguish

e.g. Quarkonia or Tetraquarks



→ **Compact** object formed from $\bar{Q}Q$ or (Qq) and $(\bar{Q}\bar{q})$

and



Hadronic-Molecules

→ **Extended** object made of $(\bar{Q}q)$ and $(Q\bar{q})$

$$\text{Bohr radius} = 1/\gamma = 1/\sqrt{2\mu E_b}$$

$$\gg 1 \text{ fm} \gtrsim \text{confinement radius}$$

for **near threshold states**

Tool: The Weinberg compositeness criterion

Weinberg Phys.Rev.137(1965)B672, Baru et al. (2004)

Expand in terms of **non-interacting** quark and meson states

$$|\Psi\rangle = \begin{pmatrix} \lambda|\psi_0\rangle \\ \chi(\mathbf{p})|h_1 h_2\rangle \end{pmatrix},$$

here $|\psi_0\rangle$ = elementary state and $|h_1 h_2\rangle$ = two-hadron cont., then


$\lambda^2 = |\langle\psi_0|\Psi\rangle|^2$ = probability to find bare state in physical state

→ λ^2 is the quantity of interest!

Crucial observation:

S. Weinberg, Phys. Rev. 130(1963)776; 131, 440 (1963)

Non-pert. hadron-hadron interactions **equivalent** to
pole term + perturbative interaction



⇒ **Dynamical information transferred into coupling**

Therefore: $\hat{H}_{hh} = \hat{H}_{hh}^0 = \vec{p}^2/(2\mu)$ contains **only kinetic terms!**

$$\hat{\mathcal{H}}|\Psi\rangle = E|\Psi\rangle, \quad \hat{\mathcal{H}} = \begin{pmatrix} \hat{H}_c & \hat{V} \\ \hat{V} & \hat{H}_{hh}^0 \end{pmatrix} \longrightarrow \chi(p) = \lambda \frac{f(p^2)}{E - p^2/(2\mu)}$$

introducing the **transition form factor** $\langle \psi_0 | \hat{V} | hh \rangle = f(p^2)$

Therefore

$$|\Psi\rangle = \lambda \begin{pmatrix} |\psi_0\rangle \\ -\frac{f(p^2)}{E_B + p^2/(2\mu)} |h_1 h_2\rangle \end{pmatrix},$$

For the normalization of the physical state we get

$$1 = \langle \Psi | \Psi \rangle = \lambda^2 \left(1 + \int \frac{d^3p}{(2\pi)^3} \frac{f^2(p^2)}{(E_B + p^2/(2\mu))^2} \right)$$

provides **connection between λ^2 and hadronic properties**

\implies **Need to understand the integral**

using
$$\int \frac{f^2(p^2) d^3p}{(p^2/(2\mu) + E_B)^2} = \frac{4\pi^2 \mu^2 f(0)^2}{\sqrt{2\mu E_B}} + \mathcal{O}\left(\frac{\sqrt{E_B \mu}}{\beta}\right)$$

for ***s*-waves**; $1/\beta =$ range of forces; $\mu f(0)^2/(2\pi) = g^2$; $\gamma = \sqrt{2\mu E_B}$

$$1 = \lambda^2 \left(1 + \frac{\mu g^2}{\gamma} + \mathcal{O}\left(\frac{\gamma}{\beta}\right) \right) \implies \boxed{g^2 = \frac{\gamma}{\mu} \left(\frac{1 - \lambda^2}{\lambda^2} \right)}$$

This gives for the residue, $g_{\text{eff(NR)}}^2 = (2\pi/\mu)\lambda^2 g^2$:

$$g_{\text{eff(NR)}}^2 = 2\pi(1 - \lambda^2)\gamma/\mu^2 \leq 2\pi\gamma/\mu^2$$

$(1 - \lambda^2) =$ Quantifies molecular component in physical state

The **structure information** is hidden in the **effective coupling**, extracted from experiment, **independent of the phenomenology** used to introduce the pole(s)

The scattering amplitude is in terms of the previous parameters

$$T_{\text{NR}}(E) = \frac{2\pi}{\mu} \frac{g^2}{E + E_B + g^2(ik + \gamma)}$$

where $k^2 = 2\mu E$ & $g^2 = \infty$ for molecule / $g^2 = 0$ for compact state

The effective range expansion reads:

$$T_{\text{NR}}(E) = -\frac{2\pi}{\mu} \frac{1}{1/a + (r/2)k^2 - ik}$$

and we get from matching coefficients

$$\frac{1}{a} = -\frac{E_B}{g^2} + \gamma \quad \Longrightarrow \quad a = -2 \frac{1 - \lambda^2}{2 - \lambda^2} \left(\frac{1}{\gamma} \right) + \mathcal{O} \left(\frac{1}{\beta} \right)$$


$$r = -\frac{1}{g^2 \mu} \quad \Longrightarrow \quad r = -\frac{\lambda^2}{1 - \lambda^2} \left(\frac{1}{\gamma} \right) + \mathcal{O} \left(\frac{1}{\beta} \right)$$

I. Matuschek et al., EPJA57(2021)3

Assume **attractive interaction**
(bound state $a < 0$, all others $a > 0$)

Molecules: $|a| \gg |r|$ and $|r| \simeq \text{range}$

Compact states: $|a| \ll |r|$ and
 $r < 0$ with $|r| \gg \text{range}$





When a changes sign (r fixed): **Molecule** \rightarrow virtual state
Compact state \rightarrow resonance

Subsummed in **compositness** $\bar{X} = 1/\sqrt{1 + |2r/a|}$

other approaches: Sekihara, Hyodo, Oset, Oller, Nieves, Jido ...
mostly relying on on-shell factorisation of the potential; little about virtual states

$\chi_{c1}(3872)$ also known as $X(3872)$



LHCb, PRD102(2020)092005

C.H. et al., PRD76(2007)034007

Data analysed employing for the rate

$$\Gamma_\rho(E)$$

$$\left| E - E_f + \frac{i}{2} [g_1^2 \sqrt{2\mu_1 E} + g_2^2 \sqrt{2\mu_2 (E - \delta)} + \Gamma_\rho(E) + \Gamma_\omega(E) + \Gamma_0] \right|^2$$

with E_f fixed to -7.18 MeV: $g_1^2 = g_2^2 = g^2 = 0.108 \pm 0.003$ such that

$$-r = 2/(\mu_1 g^2) + \sqrt{\mu_2 / (2\mu_1^2 \delta)} \simeq (3.8 + 1.4) \text{ fm} \gg 1/M_\pi$$

Does this mean $\chi_{c1}(3872)$ is a compact state?

A. Esposito et al., PRD105(2022)L031503



The second term in

$$-r = 2/(\mu_1 g_1^2) + \frac{g_2^2}{g_1^2} \sqrt{\mu_2/(2\mu_1^2 \delta)} \simeq (3.8 + 1.4) \text{ fm} \gg 1/M_\pi$$

comes from **isospin-symmetry**, $g_1^2 = g_2^2 = g^2$ and the expansion

$$ik_2 = \sqrt{2\mu_2(\delta - k_1^2/(2\mu_1))} = \sqrt{2\mu_2\delta} - \frac{1}{2} \sqrt{\frac{\mu_2}{2\mu_1^2\delta}} k_1^2 + \mathcal{O}\left(\left(\frac{k_1^2}{\mu_1\delta}\right)^2\right)$$

which “measures” the **contribution from the charged channel** and does **not have a proper isospin limit** ($\delta \rightarrow 0$). However,

- it scales with g^2 ($\rightarrow \infty$ for molecule)
- we thus see that this contribution is sizable
 \rightarrow **needs to be removed to understand structure**

Thus the quantity **relevant for the Weinberg analysis** is thus


$$-r_{\text{eff.}} = 2/(\mu_1 g^2) \leq 3.8 \text{ fm}$$

$$\Gamma_\rho(E)$$

$$\left| E - E_f + \frac{i}{2} \left[g^2 \left(\sqrt{2\mu_1 E} + \sqrt{2\mu_2 (E - \delta)} \right) + \Gamma_\rho(E) + \Gamma_\omega(E) + \Gamma_0 \right] \right|^2$$

For $E < 0$ the parts in red define E_p , the real part of pole location:

$$E_p = E_f + \frac{g^2}{2} \left(\sqrt{2\mu_1 |E_p|} + \sqrt{2\mu_2 (\delta + |E_p|)} \right) \implies g^2(E_f, E_p)$$



Since $E_p \ll \delta$ one may approximate correlation parameter free

$$g^2(E_f, 0) = -\sqrt{\frac{2}{\mu_2 \delta}} E_f$$

To remove correlation:

Express E_f by E_p

The formula that **should be used in the analysis:**

$$\Gamma_\rho(E)$$

$$\left| E - E_p + \frac{i}{2} \left[g^2 \left(\sqrt{2\mu_1 E} \mp i\gamma_1 + \sqrt{2\mu_2 (E - \delta)} - i\gamma_2 \right) + \Gamma_{\text{inel.}}(E) \right] \right|^2$$

for pole on the physical (unphysical) $D^0 \bar{D}^{*0}$ sheet and

where $\gamma_1 = \sqrt{2\mu_1 |E_p|}$ and $\gamma_2 = \sqrt{2\mu_2 (\delta + |E_p|)}$

The LHCb data only provides lower bound for g

If one allows for $\Delta LL = 1$, one finds $g^2 > 0.1$ and accordingly

$$-r_{\text{eff.}} < 4 \text{ fm} \quad \text{and} \quad \bar{X} = \frac{1}{\sqrt{1 + 2|r_{\text{eff.}}/\Re(a)|}} > 0.94 ,$$

fully **consistent with a molecular interpretation**

Similar numbers emerge for the T_{cc} state ...

→ The formulas were derived neglecting finite range corrections

→ The Wigner bound (causality!) requires $r < R \sim 1/\beta$

E.P. Wigner, Phys.Rev 98(1955)145

⇒ Zero range interactions call for neg. effective ranges

The longest range interaction is the one π exchange, however
in the charm system $\pi D\bar{D}$ can go on-shell

⇒ no fixed sign of potential


We need hadronic EFT to quantify the effects!

pert. pions: Mehen, Valderrama, Mikhasenko, ...; non-pert. pions: Baru, Filin, Du, Guo, C.H., ...

⇒ three-body calculation for T_{cc} : $r_{\text{OPE}} = +0.4 \text{ fm}$

M. L. Du et al., PRD105 (2022)014024.

E. Braaten and M. Lu, PRD76(2007)094028, C. H. et al., PRD81(2010)094028



- $E_b = 0.5 \text{ MeV}$
below nominal
threshold,
- additional width:
 $\Gamma_{\text{inel.}} = 1.5 \text{ MeV}$
- constituent width
 $0, 0.1, 1 \text{ MeV}$


Molecules with **unstable const.** can show peculiar line shapes

Strong rise above nominal threshold, because of

- nearby pole
- with **large residue**

Form depends on
interplay of scales


E. Braaten and M. Lu, PRD76(2007)094028, C. H. et al., PRD81(2010)094028



- $E_b = 0.5 \text{ MeV}$
below nominal threshold,
- additional width:
 $\Gamma_{\text{inel.}} = 1.5 \text{ MeV}$
- constituent width
0, 0.1, 1 MeV

$e^+e^- \rightarrow D^* \bar{D} \pi$ data show strong evidence for $Y(4230)$ as $D_1(2420) \bar{D}$ bound state with $E_b = 55 \text{ MeV}$

$D_1(2420)$ ($\Gamma = 30 \text{ MeV}$)
mixing into $D_1(2430)$ ($\Gamma = 300 \text{ MeV}$)



At present the data on $\chi_{c1}(3872)$ aka $X(3872)$ and T_{cc}^+ are consistent with a **molecular interpretation**, but so far a **sizeable compact component cannot be excluded**.

For more definite statements we need

- Reanalysis of LHCb data **with correlations removed**
- **Combined analysis** of inelastic and elastic channels
- Direct **measurement of line shape** (PANDA?)
- Information on **(iso)spin partner states**

Line shapes carry important structure information

... thank you very much for your attention