

# Achieving High Deuteron Tensor Polarization For Polarized Target Experiments

Allison J. Zec  
(she/her)

Univ. of New Hampshire

2023-09-26



# Deuterons and the Tensor Experiments

# What Deuterons Do That Protons Don't

## Proton

Spin- $\frac{1}{2}$  System



$$m = +\frac{1}{2}$$



$$m = -\frac{1}{2}$$

"Typical" Vector Polarization



-



$$P_z = p_+ - p_-$$

## Deuteron

Spin-1 System



$$m = +1$$



$$m = 0$$



$$m = -1$$

Vector **and** Tensor Polarization


$$\left( \text{m=+1 up} + \text{m=-1 down} \right) - 2 \text{m=0}$$

$$P_{zz} = (p_+ + p_-) - 2p_0$$

J Forest, et al, PRC **54** 646 (1996)

# Tensor Polarization Properties

If...




Then...

$$0 < P_{zz} \leq 1$$

$$P_{zz} = 0$$

$$-2 \leq P_{zz} < 0$$




- $P_z$  ranges from -1 to +1
- $P_{zz}$  ranges from -2 to +1
- In deuterons both  $P_z$  and  $P_{zz}$  can be nonzero simultaneously



# Tensor Polarization Properties

If...




Then...

$$0 < P_{zz} \leq 1$$

$$P_{zz} = 0$$

$$-2 \leq P_{zz} < 0$$



A high-luminosity tensor-polarized target has promise as a novel probe of nuclear physics


# $b_1$ Experiment

- Intended to improve upon HERMES' 2005 data
- Verifications of zero-crossing
  - Implications for Close-Kumano sum rule
- Tensor physics at quark level
- Better understanding of  $b_1$  allows discrimination of different deuteron components by spin (e.g., quarks vs gluons)

Approved by JLab with A-physics rating!

E12-13-011

The Deuteron Tensor Structure Function  $b_1$



K. Slifer *et al*, JLab C12-13-011 **Spokespersons:** K. Slifer, O.R. Aramayo, J.P. Chen, N. Kalantrians, D. Keller, E. Long, P.

Sullivan


# $b_1$ Experiment

- Intended to improve upon HERMES' 2005 data
- Verifications of zero-crossing
  - Implications for Close-Kumano sum rule
- Tensor physics at quark level
- Better understanding of  $b_1$  allows discrimination of different deuteron components by spin (e.g., quarks vs gluons)

Approved by JLab with A-physics rating!

E12-13-011

The Deuteron Tensor Structure Function  $b_1$



K. Slifer *et al*, JLab C12-13-011 **Spokespersons:** K. Slifer, O.R. Aramayo, J.P. Chen, N. Kalantrians, D. Keller, E. Long, P. Sullivan

# $A_{zz}$ Experiment


- First-of-its-kind quasielastic  $A_{zz}$  measurement
- Implications for SRC physics and deuteron wavefunction
- Widest range of  $x$  covered by a single measurement
- Measurement of  $T_{20}$  included!

**Spokespersons:** E. Long, K. Slifer, P. Solvignon, D. Day, D. Keller, D. Higinbotham

Approved by JLab with A-physics rating!

E12-15-005

Quasi-Elastic and Elastic Deuteron Tensor Asymmetries



E. Long *et al*, JLab C12-15-005

# $A_{zz}$ Experiment


- First-of-its-kind quasielastic  $A_{zz}$  measurement
- Implications for SRC physics and deuteron wavefunction
- Widest range of  $x$  covered by a single measurement
- Measurement of  $T_{20}$  included!

**Spokespersons:** E. Long, K. Slifer, P. Solvignon, D. Day, D. Keller, D. Higinbotham

Approved by JLab with A-physics rating!

E12-15-005

Quasi-Elastic and Elastic Deuteron Tensor Asymmetries



E. Long *et al*, JLab C12-15-005

## $b_1$ Systematics Estimates

Source	Systematic
Polarimetry	8.0%
Dilution/Packing Fraction	4.0%
Others	2.1%
Total	9.2%

## $A_{zz}$ Systematics Estimates

Source	$A_{zz}$ Systematic	$T_{20}$ Systematic
Polarimetry	6.0%	6.0%
Dilution Factor	6.0%	2.5%
Packing Fraction	3.0%	3.0%
Others	2.5%	2.5%
Total	9.2%	7.4%

$$A_{zz} = \frac{2}{f P_{zz}} \left( \frac{\sigma_p}{\sigma_0} - 1 \right)$$

## $b_1$ Systematics Estimates

Source	Systematic
<b>Polarimetry</b>	<b>8.0%</b>
<b>Dilution/Packing Fraction</b>	<b>4.0%</b>
Others	2.1%
Total	9.2%

## $A_{zz}$ Systematics Estimates

Source	$A_{zz}$ Systematic	$T_{20}$ Systematic
<b>Polarimetry</b>	<b>6.0%</b>	<b>6.0%</b>
<b>Dilution Factor</b>	<b>6.0%</b>	<b>2.5%</b>
<b>Packing Fraction</b>	<b>3.0%</b>	<b>3.0%</b>
Others	2.5%	2.5%
Total	9.2%	7.4%

$$A_{zz} = \frac{2}{f \overline{P_{zz}}} \left( \frac{\sigma_p}{\sigma_0} - 1 \right)$$


Both experiments require a highly ( $\geq 30\%$ ) tensor-polarized deuterium target with precise measurement of  $P_{zz}$ . How can we achieve that?

## Tensor Targets: Granite State Style



# Dynamic Nuclear Polarization (DNP)


- Using  $\mu$ waves, drive spin transitions of unpaired electrons
- Electrons transfer spin to nuclei
- Nuclear absorption spectrum gives polarimetry info



Above: Characteristic lineshape  
of the proton

C.D. Keith *et al*, NIM A 501

(2003)




Above: Diagram of the energy level transitions in the DNP process.

Adapted from *Annu. Rev. Nucl. Part. Sci.* 1997. 47:67-109

# Deuteron Polarization


- NMR at nuclear spin transition frequency drives further spin transitions
- Proton lineshape from  $-1/2 \leftrightarrow 1/2$  transition
- Deuteron lineshape has  $-1 \leftrightarrow 0$  and  $0 \leftrightarrow 1$  components
  - But NMR only gives the sum of the two
- Signal shape affected by material properties and magnetic field angle



Above: Simulated deuteron lineshape showing the contributions from both the  $-1 \rightarrow 0$  transition and the  $0 \rightarrow 1$  transition.

# ND<sub>3</sub> and Other Target Materials

C. Dulya, *et al*, NIM A **398** (1997)




- Both  $b_1$  and  $A_{zz}$  experiments call for solid ND<sub>3</sub> targets
- Polarization also done with frozen chemically-doped deuterated alcohols
- Lineshape affected by quadrupole splitting of molecule
  - Different for ND<sub>3</sub> vs butanol

*Left:* C-D, O-D bond contribution to the deuteron NMR lineshape in d-butanol

Material	Dopant & method	Polarizable nucleons % by weight
ND <sub>3</sub>	ND <sub>2</sub>	
d-ammonia	Irradiation	~30%
C <sub>4</sub> D <sub>9</sub> OD	TEMPO	
d-butanol	Chemical	23.7%


D. Crabb, W. Meyer, *Annu. Rev. Nucl. Part. Sci* **47** 67-109 (1997)

# UNH Polarized Target Lab




**The UNH polarized target group is hard at work!**

# UNH Polarized Target Lab




**The UNH polarized target group is hard at work!**

# UNH Polarized Target Lab



The UNH polarized target group is hard at work!

# UNH Polarized Target Lab



*Above: UNH LHe refrigerator*


D. M. Aliaga *et al.* NIM 976 (2020) 164277

## Refrigerator and Magnet

1 K LHe  
evaporative  
fridge, with  
2.4 W cooling  
power

5 T Nb-Sn  
superconducting  
solenoidal  
magnet

# UNH Polarized Target Lab



*Above: UNH LHe refrigerator*

D. M. Aliaga *et al.* NIM 976 (2020) 164277

## Refrigerator and Magnet

1 K LHe  
evaporative  
fridge, with  
2.4 W cooling  
power

5 T Nb-Sn  
superconducting  
solenoidal  
magnet


## Microwaves

Solid-state microwaves  
producing  $>100$  mW power  
between 136 and 144 GHz.

Source is movable by  
remote motor control.

System designed by  
Bridge12 inc.

*Below: Microwave  
source and mount.*





## Target Insert

Designed, printed and assembled in-house.  
Also houses overmodal microwave waveguide.


## Microwaves

Solid-state microwaves  
producing  $>100$  mW power  
between 136 and 144 GHz.


Source is movable by  
remote motor control.

System designed by  
Bridge12 inc.

*Right: 3D-printed  
target insert ladder  
with coils wound and  
no target material.*




*Below: Microwave  
source and mount.*



## Target Insert

Designed, printed and assembled in-house.  
Also houses overmodal microwave waveguide.

*Right: 3D-printed target insert ladder with coils wound and no target material.*




*Below left: frozen unirradiated  $\text{NH}_3$ .  
Below right: frozen doped butanol.*



## Target Material

Material, both ammonia and doped alcohols, frozen and stored on-site.

# UNH Polarized Target Lab



*Left: VME crate for NMR control system and analyzer.*

## NMR System

NMR system with LANL design sweeps at deuteron transition frequency ( $\simeq 30$  MHz).

P. McGaughey, *et al*, NIM A **995** (2021) 165045

*Below left: frozen unirradiated  $\text{NH}_3$ .  
Below right: frozen doped butanol.*



## Target Material


Material, both ammonia and doped alcohols, frozen and stored on-site.

# Tensor Polarization Measurements

# Measuring Tensor Polarization


## Vector Polarization Measurement

$$P_z = C(I_+ + I_-) \quad (1)$$



## Tensor Polarization Measurement

$$P_{zz} = C(I_+ - I_-) \quad (2)$$



where  $C$  is a dimensionless calibration constant,  $I_+ = n_+ - n_0$ , and  $I_- = n_0 - n_-$

Figures courtesy of E. Long

# NMR Curve Fitting

- Fit NMR lineshape with procedure from C. Dulya *et al*, NIM A **398** (1997) 109-125
- Includes effects from molecular bond quadrupole terms
- Can naively use peak height ratio  $r$  to estimate polarization

$$\begin{aligned} P_z &= \frac{r^2 - 1}{r + r^2 + 1} \\ P_{zz} &= \frac{r^2 - 2r + 1}{r^2 + r + 1} \end{aligned} \quad (3)$$

- Then compare *ratio* and *area* methods for  $P_{zz}$  measurement consistency

*Right:* Parts of the curve fitting method suggested by C. Dulya *et al*.

$R, A, \eta, \phi$   compacting variables

$$\begin{aligned} \rho^2 &= \sqrt{A^2 + [1 - \epsilon R - \eta \cos(2\phi)]^2} & R &= \frac{\omega - \omega_d}{3\omega_q} \\ \cos(\alpha) &= \frac{1 - \epsilon R - \eta \cos(2\phi)}{\rho^2} & -3 \leq R \leq 3 \end{aligned}$$

functional form of signal 

$$f_\epsilon(R, A, \eta, \phi) = \frac{1}{2\pi\rho} \left\{ 2\cos\left(\frac{\alpha}{2}\right) \left[ \arctan\left(\frac{Y^2 - \rho^2}{2Y\rho\sin(\frac{\alpha}{2})}\right) + \frac{\pi}{2} \right] \right. \\ \left. + \sin\left(\frac{\alpha}{2}\right) \ln\left(\frac{Y^2 + \rho^2 + 2Y\rho\cos(\frac{\alpha}{2})}{Y^2 + \rho^2 - 2Y\rho\cos(\frac{\alpha}{2})}\right) \right\}$$

$\epsilon = \pm 1$

phi average 

$$F_\epsilon \approx \frac{1}{J+1} \sum_{j=0}^J \frac{\sqrt{3}f_\epsilon(R, A, \eta, \phi_j)}{\sqrt{3 - \eta \cos(2\phi_j)}}$$


positive & negative spin flips 

$$\chi''(r, R) \propto \frac{1}{\omega_q} \left\{ \left[ \frac{r^2 - r^{1-3\theta R}}{r^{1-\theta R}} \right] F_+(R) + \left[ \frac{r^{1+3\theta R} - 1}{r^{1+\theta R}} \right] F_-(R) \right\}$$



$\theta = \omega_q / \omega_d$

# Tensor Polarization Measurement (UVA)

- UVA-pioneered tensor enhancement technique
- Additional RF coils drive spin flips
- Manipulates area of NMR curve
- Small frequency range
- UVA lab achieved  $P_{zz}=31.1\pm 8.5\%$  with ssRF technique




Right: ssRF coil schematic. Below: NMR lineshapes without and with ssRF applied. Figures reproduced from D. Keller, et al. NIM A **981** 164504 (2020)




# Tensor Polarization Measurement (UVA)

- UVA-pioneered tensor enhancement technique
- Additional RF coils drive spin flips
- Manipulates area of NMR curve
- Small frequency range
- UVA lab achieved  $P_{zz}=31.1\pm 8.5\%$  with ssRF technique




Right: ssRF coil schematic. Below: NMR lineshapes without and with ssRF applied. Figures reproduced from D. Keller, et al. NIM A **981** 164504 (2020)





# Tensor Polarization (UNH)

- Fit with Dulya procedure closely matches data from recent UNH cooldown
  - C. Dulya et al, NIM A 398 (1997) 109-125
- Reconstruct spin-flip and quadrupole curves from fit parameters
- With reconstruction can do more in-depth polarization analysis
- Fit method works very well for UNH data!
- **New helium reliquifier system will enhance UNH's ability to run and get physics data!**




Above: Curve fit of NMR lineshape from recent target cooldown at UNH.

# Real & Imaginary Fits

- Can now manually set NMR phase angle  $\phi$  during cooldowns
- Fit using a rotation of the absorptive ( $\chi''$ ) and dispersive ( $\chi'$ ) around phase angle:

$$\begin{aligned}\text{Real} &= \chi'' \cos \phi - \chi' \sin \phi \\ \text{Imag} &= \chi'' \sin \phi + \chi' \cos \phi\end{aligned}\quad (4)$$


- Can fit a simultaneous mixture of real and imaginary
- First fits with the new method match data well, look very promising!



Above: Fit of recent cooldown data using real and imaginary parts. Fit is compared with an “imaginary only” signal and then fitted for a phase mistune.

# Real & Imaginary NMR Signals


- Switch from real to imaginary lineshape by tuning phase
- Use fitting for real and imaginary lineshapes differently
- Demonstrated resilience to having phase not tuned perfectly
- Real and imaginary measurements match each other well!



Above: Data from recent UNH cooldown with both real and imaginary line data for both vector and tensor polarization. Figure courtesy of M. McClellan.

# Summary

## Professors



Karl Slifer



Elena Long




Nathaly  
Santiesteban

## Postdocs




Allison Zec



David Ruth

## Graduate Students



Michael McClellan



Zoe Wolters




Anchit Arora

**Thank you  
to the UNH  
PolTarg  
Group and  
our  
collaborators  
at UVA!**

- Tensor target experiments approved!
- Conditionals removed for both experiments
- target development at UNH and UVA

- Improvements with ss-RF for better tensor polarization
- UNH group polarizing more (publications upcoming. . .)
- Exciting new developments upcoming!


# Trans & Nonbinary Physicists



The Trans and Nonbinary Physicists Discord server is an online community for transgender and nonbinary physicists — from enthusiasts to professors! — to socialize, network, and support one another. All are welcome, and so far we have over 200 members from across the world!

**Follow**  
**@transphysicists**  
**on twitter!**

# Trans & Nonbinary Physicists



The Trans and Nonbinary Physicists Discord server is an online community for transgender and nonbinary physicists — from enthusiasts to professors! — to socialize, network, and support one another. All are welcome, and so far we have over 200 members from across the world!

**Follow**  
**@transphysicists**  
**on twitter!**


**Questions, comments, concerns, observations?**

## Backup Slides

# Protons & Deuterons

## Proton

Spin- $\frac{1}{2}$  System



Three valence quarks + gluons and sea quarks


No nucleon-nucleon interactions

$$m = \pm \frac{1}{2}$$

S. Kumano, IOP Proc. Tens. Pol. Targ. (2014)

## Deuteron

Spin-1 System



Proton-Neutron bound state

Simplest nuclear system: nucleon interaction effects

$$m = \pm 1, 0$$




# Elastic Tensor Analyzing Power

For  $1.5 \leq x \leq 2.0$

$$T_{20} \approx \frac{A_{zz}}{\sqrt{2}d_{20}} \quad (5)$$

- Third of three elastic scattering functions of deuteron
- Extracted by measuring  $A_{zz}$  near elastic peak
- Current data doesn't constrain models well at high  $x$

M. Kohl Nucl Phys A **805** (2008)



**Above:**  $T_{20}$  with current measurements and theoretical models.

R. Holt, R. Gilman Rept.Prog.Phys. **75** (2012)


# $b_1$ Kinematics

- Approved for 30 days of physics running + 10.8 days overhead
- 11 GeV beam incident on polarized target
- 9.2% systematic error on  $A_{zz}$
- Forward scattering angles

	$x_{Bj}$	$Q^2$ [GeV <sup>2</sup> ]	$E'_0$ [GeV]	$\theta_{e'}$ [°]
SHMS	0.15	1.21	6.70	7.35
SHMS	0.30	2.00	7.45	8.96
SHMS	0.452	2.58	7.96	9.85
HMS	0.55	3.81	7.31	12.50

E12-13-011

The Deuteron Tensor Structure Function  $b_1$



K. Slifer *et al*, JLab C12-13-011


# $A_{zz}$ Kinematics

- Approved for 34 days of physics running + 10.3 days overhead
  - 25 days 8.8 GeV beam
  - 8 days 6.6 GeV beam
  - 1 day of 2.2 GeV beam
- 9.2% systematic error on  $A_{zz}$ , 7.4% on  $T_{20}$
- Forward scattering angles

	$E_0$ [GeV]	$Q^2$ [GeV <sup>2</sup> ]	$E'_0$ [GeV]	$\theta_{e'}$ [°]
SHMS (S1)	8.8	1.5	8.36	8.2
HMS (H1)	8.8	2.9	7.26	12.2
SHMS (S2)	6.6	0.7	6.35	7.5
HMS (H2)	6.6	1.8	5.96	12.3
SHMS (S3)	2.2	0.2	2.15	10.9
HMS (H3)	2.2	0.3	2.11	14.9

E12-15-005

Quasi-Elastic and Elastic Deuteron  
Tensor Asymmetries





E. Long *et al*, JLab C12-15-005

# From Spin Flips To Lineshape

- Simulation derived from Cohen & Reif model (1957)
- Assume random distribution of quadrupole angles  $\theta$  from  $0^\circ$ - $90^\circ$
- Then calculates transition frequency based on angle and energy level
- Performs 125,000 spin flips into 300 angle bins, giving NMR signal
- Simulation can reproduce results from UVA polarized target lab, as well as previous UNH cooldowns

*Right:* Quadrupole angles randomly distributed in B-field. *Below:* Spin transition frequencies for each angle  $\theta$  and each transition.



## BACKUP: Tensor Polarization Analysis


# Thermal Equilibrium & Enhancement

Deuteron thermal equilibrium (TE) polarization before microwave irradiation:

$$P(1) = \frac{4 \tanh(\frac{g_i \mu_i B}{2k_B T})}{3 + \tanh^2(\frac{g_i \mu_i B}{2k_B T})} \quad (6)$$

Only 0.1% polarization at 5 T and 1 K.

TE signal can be used for calibration if detected. Signal is then enhanced with microwaves.



Above: Deuteron TE signal from CLAS target. From C. Keith *et al*, NIM A 501 (2003). Right: Polarization curve during enhancement.

