




LIGHT MESON DECAYS FROM PHOTON-INDUCED REACTIONS WITH CLAS

10 April 2015 | GHP 2015 | Michael C. Kunkel | IKP-1 | on behalf of the CLAS Collaboration and LMD group

Thomas Jefferson National Laboratory




Continuous Electron Beam Accelerator Facility (CEBAF) at 12 GeV



Aerial View

CEBAF Large Acceptance Spectrometer (CLAS)



CLAS

The g11 and g12 experiments



<i>g11</i> $\gamma p \rightarrow pX$	<i>g12</i> $\gamma p \rightarrow pX$
<i>60 - 65 nA 4.023 GeV e⁻ beam</i> <i>0.803 < E_y < 3.815</i>	<i>60 - 65 nA 5.714 GeV e⁻ beam</i> <i>1.142 < E_y < 5.425</i>
<i>40 cm (2 cm radius) liquid H₂ target</i> <i>placed at CLAS center</i>	<i>40 cm (2 cm radius) liquid H₂ target</i> <i>placed -90cm from CLAS center</i>
<i>Trigger required at least two charged tracks in different sectors</i>	<i>Trigger required at least two charged tracks in different sectors for E_y > 3.6</i>
<i>20x10⁹ productions triggers as 21 TB of raw data</i>	<i>26x10⁹ productions triggers as 128 TB of raw data</i>
	<i>Cherenkov Counters and Electromagnetic Calorimeter in trigger for entire E_y range</i>

CLAS Light Meson Decay (LMD) Program Institutional Contributors

- Old Dominion University, Norfolk, Virginia 23529
- Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188300, Russia
- INFN, Sezione di Genova, 16146 Genova, Italy
- The George Washington University, Washington, DC 20052
- Florida State University, Tallahassee, Florida 32306
- University of South Carolina, Columbia, South Carolina 29208
- Arizona State University, Tempe, Arizona 85287-1504
- Indian Institute of Technology Indore, Khandwa Road, Indore-452017, Madhya Pradesh, India
- Institut für Kernphysik, Forschungszentrum, Jülich, Germany
- Jülich Center for Hadron Physics, Forschungszentrum Jülich, 52425 Jülich, Germany
- Florida International University, Miami, Florida 33199
- Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606
- University of Glasgow, Glasgow G12 8QQ, United Kingdom
- Catholic University of America, Washington, DC 20064
- Institut für Experimentalphysik I, Ruhr Universität Bochum, 44780 Bochum, Germany
- Helmholtz-Institut für Strahlen- und Kernphysik (Theorie) and Bethe
- Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany
- Rensselaer Polytechnic Institute, Troy, New York 12180-3590
- Skobeltsyn Nuclear Physics Institute, 119899 Moscow, Russia
- Department of Physics and Astronomy, University of Georgia, Athens, GA 30602, USA
- Institut für Theoretische Physik II, Ruhr-Universität Bochum, D-44780 Bochum, Germany
- Norfolk State University, Norfolk, VA 23504, USA
- Institute for Advanced Simulation, Forschungszentrum Jülich, 52425 Jülich, Germany

Light Meson Decays in CLAS



CLAS Light Meson Decay (LMD) Program was established to investigate

Meson Decay	Physics	Meson Decay	Physics
$\pi^0 \rightarrow e^+ e^- \gamma$	<i>Heavy photon upper limit</i>	$\eta(') \rightarrow \pi \pi^+ \gamma$	<i>Box anomaly</i>
$\eta(') \rightarrow e^+ e^- \gamma$	<i>Transition Form Factor</i>	$\omega \rightarrow \pi \pi^+ \gamma$	<i>Upper limit branching ratio <3.6x10⁻³</i>
$\omega \rightarrow \pi^0 e^+ e^-$	<i>Transition Form Factor</i>	$\eta, \omega, \Phi \rightarrow \pi \pi^+ \pi^0$	<i>Dalitz plot analysis</i>
$\eta(') \rightarrow \pi^0 e^+ e^-$	<i>C violation</i>	$\eta' \rightarrow \pi \pi^+ \eta$	<i>Dalitz plot analysis/meson mixing</i>
$\eta(') \rightarrow \pi \pi^+ e^+ e^-$	<i>CP violation</i>	$\Phi \rightarrow \pi \pi^+ \eta$	<i>G-parity violation</i>


Box Anomaly of $\eta(') \rightarrow \pi^-\pi^+\gamma$

Njencheu Georgie Mbianda, Moskov Amaryan;

Old Dominion University

Motivation:

- The 2 photon decay of $\pi^0, \eta, \eta' \rightarrow \gamma\gamma$ proceed from the understood triangle or axial anomaly. While radiative decays of $\eta, \eta' \rightarrow \pi^-\pi^+\gamma$ are related to a less understood box anomaly.



- With an analysis of the photon energy distribution of the radiative decays of η and η' , the decay widths are determined by the box anomaly in the chiral limit.
- Radiative decays from CLAS will test the box anomaly, including FSI of the pions. FSI occur because beyond the chiral limit, quarks have mass.

Box Anomaly

$$\frac{d\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma)}{ds_{\pi\pi}} = |AP(s_{\pi\pi}) F_V(s_{\pi\pi})|^2 \Gamma_0(s_{\pi\pi})$$

$$\Gamma_0(s_{\pi\pi}) = \frac{1}{3 \cdot 2^{11} \cdot \pi^3 M_\eta^3} \left(M_\eta^2 - s_{\pi\pi} \right)^3 s_{\pi\pi} \cdot \beta_\pi^3$$

with $\beta_\pi = \sqrt{1 - 4M_\pi^2/s_{\pi\pi}}$.

- the pion vector form factor can be approximated by the polynomial


$$|F_V(s_{\pi\pi})| \approx 1 + (2.12 \pm 0.01)s_{\pi\pi} + (2.13 \pm 0.01)s_{\pi\pi}^2 + (13.80 \pm 0.14)s_{\pi\pi}^3$$

- Expansion around $s_{\pi\pi} = 0$ gives the process specific function

$$P(s_{\pi\pi}) = 1 + \alpha \cdot s_{\pi\pi} + \mathcal{O}(s_{\pi\pi}^2)$$

- from which α can be measured.

Experimental data from WASA-at-COSY(η) and CRYSTAL BARREL(η')




$$s_{\pi\pi} = m^2 - 2E_\gamma m$$

[1] F. Stollenwerk et al., Phys. Lett. B707:184-190, 2012



[2] A. Abele et al. Phys. Lett. B402, 195 (1997).

Radiative Decay



CLAS data yield for $\gamma p \rightarrow p\{\eta, \eta' \rightarrow \pi^+\pi^-\gamma\}$ from g11 data set

CLAS Uncorrected Data


$$\eta \rightarrow \pi^+ \pi^- \gamma$$

$$\eta' \rightarrow \pi^+ \pi^- \gamma$$


Dalitz Plot of $\eta' \rightarrow \pi^- \pi^+ \eta$

Sudeep Ghosh, Anki Roy;

IIT





Motivation:

- Dalitz plot of $\eta' \rightarrow \pi^- \pi^+ \eta$ provides kinematic information of the decay, enabling the studying of low energy dynamics of QCD and heavier mass pseudoscalar mesons.
- The $\eta' \rightarrow \pi^- \pi^+ \eta$ decay has a low Q-value due to relatively heavier decay products, thus helping us to test and limit the effective chiral Lagrangian theory
- $f(X, Y) = N \cdot (1 + a(Y) + b(Y)^2 + c(X) + d(X)^2)$

Parameters	VES	Theory	BESIII	Stat. err. in BESIII	Stat. err. in CLAS
a	-0.127 ± 0.018	-0.116 ± 0.011	-0.047 ± 0.012	± 0.011	± 0.004
b	-0.106 ± 0.032	-0.042 ± 0.034	-0.069 ± 0.021	± 0.019	± 0.006
c	$+0.015$...	$+0.019 \pm 0.012$	± 0.011	± 0.004
d	-0.082 ± 0.019	$+0.010 \pm 0.019$	-0.073 ± 0.013	± 0.012	± 0.004

Dalitz Plot of $\gamma p \rightarrow p\eta' \rightarrow p\pi^-\pi^+\eta$



Dalitz variables for $\pi^-\pi^+\eta$


$$X = \frac{\sqrt{3}(T_{\pi^+} - T_{\pi^-})}{Q}$$

$$Y = \frac{(m_\eta + 2m_\pi)}{m_\pi} \cdot \frac{T_\eta}{Q} - 1$$



where $T_i (i = \pi^+, \pi^-, \eta)$ is the kinetic energy of a given particle in the rest frame of η' and $Q = T_{\pi^+} + T_{\pi^-} + T_\eta$

Data Analysis

Data satisfying the cross-section is fed to the Dalitz plot of 30x30 ($X(-1.5,1.5) \times Y(-1.5,1.5)$) and then a pol-3 background subtraction performed in the $3\sigma(0.946-0.97 \text{ GeV})$ region of every Dalitz bin



Dalitz plot bin with lower statistics



Dalitz plot bin with higher statistics

Dalitz Plot of $\eta \rightarrow \pi^- \pi^+ \pi^0$


JPAC: Adam Szczepaniak, Diane Schott, Peng Guo et. al.

Jlab

Motivation:

- $\eta \rightarrow \pi^- \pi^+ \pi^0$ is sensitive to isospin breaking, which in QCD originates from the mass difference between the up and down quarks.







$$A(s, t) = \sum_J^{J_{max}} (2J + 1) d_{1,0}^J(\theta_s) f_J(s) + \sum_J^{J_{max}} (2J + 1) d_{1,0}^J(\theta_t) f_J(t) + \sum_J^{J_{max}} (2J + 1) d_{1,0}^J(\theta_u) f_J(u)$$



- The isobar model assumes quasi 2-body decay and is insufficient for some channels
- It is important to construct amplitudes which contain all the known physics such as 3-body interactions, coupled channel, unitarity, analyticity, etc.
- The $\eta \rightarrow \pi^- \pi^+ \pi^0$ analysis is building in the three-body interaction (unitarity and analyticity) as a first step for future experimental analysis tools.

Dalitz Plot


Fit after background subtraction



Transition Form Factors


Susan Schamand, Michaela Schever, Michael C. Kunkel;

Institut für Kernphysik, Forschungszentrum Jülich



Motivation:


- In the VMD model the transition form factors provides insight into the meson charge radius.
- For pseudoscalar mesons η and η' , ratio of form factors provides information on mixing angle.
- For vector meson ω there currently exist discrepancy in the measurement of the form factor with VMD model.
- The knowledge of the η form factor is also needed for the interpretation of the g-2 experiment.
- g12 experiment collected world breaking data samples of the $p e^+ e^- X$ reaction using Cherenkov Counters and an Electromagnetic Calorimeter



Transition Form Factors

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}} = \frac{2\alpha}{3\pi q^2} \left(1 - \frac{q^2}{m_P^2}\right)^3 \left(1 - \frac{4m_l^2}{q^2}\right)^{1/2} \left(1 + \frac{2m_l^2}{q^2}\right) |_{\text{Q.E.D}}$$

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}}|_{\text{measured}} = \frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 \Gamma_{P \rightarrow \gamma\gamma}}|_{\text{Q.E.D}} |F(q^2)|^2$$






η' Transition Form Factor

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}} = \frac{2\alpha}{3\pi q^2} \left(1 - \frac{q^2}{m_P^2}\right)^3 \left(1 - \frac{4m_l^2}{q^2}\right)^{1/2} \left(1 + \frac{2m_l^2}{q^2}\right) |_{\text{Q.E.D}}$$

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}}|_{\text{measured}} = \frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 \Gamma_{P \rightarrow \gamma\gamma}}|_{\text{Q.E.D}} |F(q^2)|^2$$


L.G. Landsberg, Electromagnetic decays of light mesons




η Transition Form Factor

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}} = \frac{2\alpha}{3\pi q^2} \left(1 - \frac{q^2}{m_P^2}\right)^3 \left(1 - \frac{4m_l^2}{q^2}\right)^{1/2} \left(1 + \frac{2m_l^2}{q^2}\right) |_{\text{Q.E.D}}$$

$$\frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 d\Gamma_{P \rightarrow \gamma\gamma}}|_{\text{measured}} = \frac{d\Gamma_{P \rightarrow l^+ l^- \gamma}}{dq^2 \Gamma_{P \rightarrow \gamma\gamma}}|_{\text{Q.E.D}} |F(q^2)|^2$$



Recent results the η transition form factor with errors. Image Source: Phys. Rev. C 89, 044608



CLAS projected errors on η transition form factor

Currently seeking applicant for analysis

ω Transition Form Factor



old issue, new interest:

- modern theoretical models
- recent measurements


PHYSICAL REVIEW D 88, 054013 (2013)

Study of $e^+ e^- \rightarrow \omega \pi^0 \rightarrow \pi^0 \pi^0 \gamma$ in the energy range 1.05–2.00 GeV with the SND detector


- with CLAS *g12 experiment*
dilepton final state



Still includes in-peak background




a first peak at the Mee distribution:




Physics Beyond SM

Motivation:



- Scalar Boson






- Vector Boson
(dark/heavy photon)



Physics Beyond SM




Physics Beyond SM




Entries/(1MeV)


Physics Beyond SM





Invariant mass $M(e^+e^-)$



PL B720 (2013) 111-115



Physics Beyond SM




$m_{A'} (\text{GeV})$

Summary

- CLAS LMD program has established a wide range of physics topics related to meson decays
- CLAS LMD program has global participation
- Statistics of CLAS data will enable precise measurements for the LMD physics program including but not limited to
 - Study $\pi^-\pi^+$ FSI within the anomalous decay $\eta(')\rightarrow\pi^-\pi^+\gamma$
 - Dalitz plot variable measurements
 - Transition form factors of pseudoscalar and vector mesons
 - Possibility of upper limit on heavy photon

BACKUP START HERE

CLAS Uncorrected Data



$$\alpha = (1.22 \pm 0.47) \text{GeV}^{-2}$$

- Need to work on MC to improve statistical precision

Tools

Home / Browse / AmpTools


Amp Tools AmpTools
Brought to you by: mashephe

Summary Files Reviews Support Wiki Mailing Lists Tickets ▾ Code

2 Downloads (This Week)
Last Update: 2013-07-18

Download AmpTools_v0.6.2.1.gz

Browse All Files



Description


A utility library for performing amplitude analysis on particle physics data.

[AmpTools Web Site >](#)




- **AmpTools framework:**
- It contains directories:
 - AmpTools/ :includes fitter and plotter
 - Tutorials/ : includes amplitudes, data
- User defined configuration files lets you define:
 - Initial fit parameters
 - Step size
 - Fix parameters
 - Real or imaginary parameters
 - Can call different amplitudes without changing or recompiling code

CLAS Uncorrected Data


$\eta \rightarrow \pi^+ \pi^- \gamma$




$\eta' \rightarrow \pi^+ \pi^- \gamma$



Experimental data from WASA-at-COSY(η) and CRSTAL BARREL(η') with error weighted fits.







$$s_{\pi\pi} = m^2 - 2E_\gamma m$$




$$\alpha = (1.96 \pm 0.27 \pm 0.02) \text{ GeV}^{-2}; \quad \alpha' = (1.80 \pm 0.49 \pm 0.04) \text{ GeV}^{-2}$$

Dalitz Plot



Extraction Cross-Sections

Comparing g11 $\eta \rightarrow \pi^-\pi^+\pi^0$ to g12 $\eta \rightarrow \pi^-\pi^+\pi^0$ using AmpTools



Light Meson Decays in CLAS




CLAS Light Meson Decay (LMD) Program was established to investigate

Meson Decay	Physics	Data Set
$\eta(')\rightarrow\pi\pi^+\gamma$	<i>Box anomaly</i>	$g11, g12$
$\omega\rightarrow\pi\pi^+\gamma$	<i>Upper limit branching ratio</i>	$g11, g12$
$\eta, \omega, \Phi\rightarrow\pi\pi^+\pi^0$	<i>Dalitz plot analysis</i>	$g11, g12$
$\eta'\rightarrow\pi\pi^+\eta$	<i>Dalitz plot analysis/meson mixing</i>	$g11, g12$
$\Phi\rightarrow\pi\pi^+\eta$	<i>G-parity violation</i>	$g11, g12$
$\Phi\rightarrow\omega\gamma$	<i>C violation, rare decay</i>	$g11, g12$
<i>NULL</i>	<i>Invisible decay</i>	$g11, g12$
f_1	<i>isospin symmetry breaking, f_1</i>	$g11, g12$


ω Transition Form Factor

$$\frac{d\Gamma_{\omega \rightarrow l^+ l^- \pi^0}}{dq^2 d\Gamma_{\omega \rightarrow \pi^0 \gamma}} = \frac{\alpha}{3\pi q^2} \left(\left(1 + \frac{q^2}{m_\omega^2 - m_{\pi^0}^2} \right)^2 - \frac{4m_\omega^2 q^2}{m_\omega^2 - m_{\pi^0}^2} \right)^{\frac{3}{2}} \left(1 - \frac{4m_l^2}{q^2} \right)^{1/2} \left(1 + \frac{2m_l^2}{q^2} \right) |Q.E.D|$$


$$\frac{d\Gamma_{\omega \rightarrow l^+ l^- \pi^0}}{dq^2 d\Gamma_{\omega \rightarrow \pi^0 \gamma}}|_{\text{measured}} = \frac{d\Gamma_{\omega \rightarrow l^+ l^- \pi^0}}{dq^2 d\Gamma_{\omega \rightarrow \pi^0 \gamma}}|_{\text{Q.E.D}} |F(q^2)|^2$$



Recent results the ω transition form factor with errors. Image Source: Conference Proceedings



CLAS data yield from $\gamma p \rightarrow p e^+ e^- X$ with
 $M_x^2(p e^+ e^-) = M^2_{\pi^0} \pm 0.01 \text{ GeV}^2$



CLAS data yield from $\gamma p \rightarrow p e^+ e^- X$ with
 $M_x(p) = M_\omega \pm 0.031 \text{ GeV}$