

Quantum dynamics of entanglement and hadronization in jet production in the massive Schwinger model

David Frenklakh




2301.11991, 2404.00087

[Florio, DF, Ikeda, Kharzeev, Korepin, Shi, Yu]

+ work in progress

with A.Florio, S.Griener, D.Kharzeev,

A. Palermo and S. Shi




APS GHP 2025 Workshop

Anaheim

3.14.2025


Background: QCD jets

Jet: a collimated set of hadrons emerging in a high-energy collision.



Background: QCD jets

Jet: a collimated set of hadrons emerging in a high-energy collision.




fragmentation:

q, \bar{q}, g

hadronization:

$\pi, K, p, \Lambda \dots$



detection



Jets often emerge in pairs, e.g. in e^+e^- annihilation.

Background: QCD jets


Jet: a collimated set of hadrons emerging in a high-energy collision.



Jets often emerge in pairs, e.g. in e^+e^- annihilation.

Originating in the same process, the partons possess **quantum entanglement**.

Motivation




How to understand entanglement in jet fragmentation?

Real-time quantum process requires

Real-time quantum simulation

Motivation




How to understand entanglement in jet fragmentation?

Real-time quantum process requires

Real-time quantum simulation

A nice testbed for learning
interesting physics with methods
suitable for quantum simulation

Motivation



How to understand entanglement in jet fragmentation?

Real-time quantum process requires

Real-time quantum simulation

A nice testbed for learning
interesting physics with methods
suitable for quantum simulation

Why Schwinger model?

- Simple enough for a first-principle real-time quantum simulation
- Has a lot of similarity with QCD in 3+1

Outline

- Schwinger model + jets
- Local operators and thermalization
- Entanglement, Schmidt states and hadronization

Schwinger model

Single-flavor (1+1)-dimensional QED:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^2 + \bar{\psi}(i\gamma^\mu\partial_\mu - g\gamma^\mu A_\mu + m)\psi$$

Features include:

- No magnetic field/no dynamical photons
- Linear potential between “quarks” – confinement
- Chiral condensate (spontaneous chiral symmetry breaking at $m=0$)

Schwinger model

Single-flavor (1+1)-dimensional QED:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^2 + \bar{\psi}(i\gamma^\mu\partial_\mu - g\gamma^\mu A_\mu + m)\psi$$

Features include:

- No magnetic field/no dynamical photons
- Linear potential between “quarks” – confinement
- Chiral condensate (spontaneous chiral symmetry breaking at $m=0$)

Massless case is exactly solvable, e.g. by bosonization:

$$\mathcal{L} = \frac{1}{2}(\partial_\mu\phi)^2 - \frac{1}{2}m_B^2\phi^2, \quad m_B^2 = \frac{g^2}{\pi}$$

Schwinger model and jets: history

1974


Vacuum polarization and the absence of free quarks

A. Casher,* J. Kogut,† and Leonard Susskind‡

Massless Schwinger model with external source:

$$j_0^{\text{ext}} = g\delta(z - t), \quad j_1^{\text{ext}} = g\delta(z - t) \quad \text{for } z > 0,$$

$$j_0^{\text{ext}} = -g\delta(z + t), \quad j_1^{\text{ext}} = g\delta(z + t) \quad \text{for } z < 0,$$



Schwinger model and jets: history

1974


Vacuum polarization and the absence of free quarks

A. Casher,* J. Kogut,† and Leonard Susskind‡

Massless Schwinger model with external source:

$$j_0^{\text{ext}} = g\delta(z-t), \quad j_1^{\text{ext}} = g\delta(z-t) \quad \text{for } z > 0,$$

$$j_0^{\text{ext}} = -g\delta(z+t), \quad j_1^{\text{ext}} = g\delta(z+t) \quad \text{for } z < 0,$$




2012

Jet energy loss and fragmentation in heavy ion collisions

Dmitri E. Kharzeev^{1,2} and Frashër Loshaj¹

$$\phi(x) = \theta(t^2 - z^2) [1 - J_0(m\sqrt{t^2 - z^2})]$$

$$j^0 = \partial_z \phi$$



Schwinger model and jets: history

1974

Vacuum polarization and the absence of free quarks

A. Casher,* J. Kogut,† and Leonard Susskind‡

Massless Schwinger model with external source:

$$j_0^{\text{ext}} = g\delta(z-t), \quad j_1^{\text{ext}} = g\delta(z-t) \quad \text{for } z > 0,$$

$$j_0^{\text{ext}} = -g\delta(z+t), \quad j_1^{\text{ext}} = g\delta(z-t)$$




2012

Jet energy loss and fragmentation in heavy ion collisions

Dmitri E. Kharzeev^{1,2} and Frashër Loshaj¹

$$\phi(x) = \theta(t^2 - z^2) [1 - J_0(m\sqrt{t^2 - z^2})]$$

$$j^0 = \partial_z \phi$$



Classical treatment is mostly sufficient
in the exactly solvable **massless** case

However, **massive** fermion case
is not exactly solvable and
inherently **quantum**

The massive Schwinger model on the lattice

Continuum: $H = \int dx \left[\frac{1}{2} E^2 + \bar{\psi} (-i\gamma^1 \partial_1 + g\gamma^1 A_1 + m) \psi \right]$

Temporal gauge
 $A_0 = 0$

The massive Schwinger model on the lattice

Continuum: $H = \int dx \left[\frac{1}{2} E^2 + \bar{\psi} (-i\gamma^1 \partial_1 + g\gamma^1 A_1 + m)\psi \right]$ Temporal gauge
 $A_0 = 0$

Fermion $\psi(an) \rightarrow \frac{1}{\sqrt{a}} \begin{pmatrix} \chi_{2n} \\ \chi_{2n-1} \end{pmatrix}$ Kogut-Susskind

$\{\psi_a(x), \psi_b^\dagger(y)\} = \delta_{ab}\delta(x-y) \rightarrow \{\chi_i, \chi_j^\dagger\} = \delta_{ij}$

N sites encode
 $N/2$ physical sites

The massive Schwinger model on the lattice

Continuum: $H = \int dx \left[\frac{1}{2} E^2 + \bar{\psi} (-i\gamma^1 \partial_1 + g\gamma^1 A_1 + m) \psi \right]$ Temporal gauge
 $A_0 = 0$

Fermion $\psi(a n) \longrightarrow \frac{1}{\sqrt{a}} \begin{pmatrix} \chi_{2n} \\ \chi_{2n-1} \end{pmatrix}$ Kogut-Susskind

$\{\psi_a(x), \psi_b^\dagger(y)\} = \delta_{ab} \delta(x - y) \longrightarrow \{\chi_i, \chi_j^\dagger\} = \delta_{ij}$ N sites encode
 $N/2$ physical sites

Gauge field $E(x = a n) \longrightarrow L_n$

Gauss law $\partial_1 E - g j^0 = 0 \longrightarrow L_n - L_{n-1} - q_n = 0, \quad q_i = \chi_i^\dagger \chi_i + \frac{(-1)^i - 1}{2}$

With **open boundary conditions** the electric field is fully determined by the fermions

Mapping to a spin chain (optional)

X, Y, Z – Pauli matrices

$$X_n \equiv I \underset{1^{\text{st}}}{\otimes} \cdots \otimes I \underset{(n-1)^{\text{th}}}{\otimes} X \underset{n^{\text{th}}}{\otimes} I \underset{(n+1)^{\text{th}}}{\otimes} \cdots \otimes I \text{ etc.}$$

Mapping to a spin chain (optional)

X, Y, Z – Pauli matrices

$$\chi_n = \frac{X_n - iY_n}{2} \prod_{j=1}^{n-1} (-iZ_j),$$

$$\chi_n^\dagger = \frac{X_n + iY_n}{2} \prod_{j=1}^{n-1} (iZ_j),$$

$$X_n \equiv I \otimes \cdots \otimes I \otimes X \otimes I \otimes \cdots \otimes I \text{ etc.}$$

$\underbrace{}_{1^{\text{st}}}$
 $\underbrace{}_{(n-1)^{\text{th}}}$
 $\underbrace{X}_{n^{\text{th}}}$
 $\underbrace{I}_{(n+1)^{\text{th}}}$

Jordan-Wigner transformation

Mapping to a spin chain (optional)

X, Y, Z – Pauli matrices

$$X_n \equiv I \otimes \dots \otimes I \otimes X \otimes I \otimes \dots \otimes I \quad \text{etc.}$$

$\underset{1^{\text{st}}}{I}$
 \dots
 $\otimes \underset{(n-1)^{\text{th}}}{I} \otimes \underset{n^{\text{th}}}{X} \otimes \underset{(n+1)^{\text{th}}}{I} \otimes \dots \otimes I$

$$\chi_n = \frac{X_n - iY_n}{2} \prod_{j=1}^{n-1} (-iZ_j),$$

$$\chi_n^\dagger = \frac{X_n + iY_n}{2} \prod_{j=1}^{n-1} (iZ_j),$$

Jordan-Wigner transformation

Spin chain Hamiltonian:

$$H^L = \frac{1}{4a} \sum_{n=1}^{N-1} (X_n X_{n+1} + Y_n Y_{n+1}) + \frac{m}{2} \sum_{n=1}^N (-1)^n Z_n + \frac{ag^2}{2} \sum_{n=1}^{N-1} L_n^2$$

Kinetic term

Mass term

Nonlocal
electric field term

$$L_n = \sum_{i=1}^n q_i$$

$$q_{n,t} = \frac{\langle Z_n \rangle_t + (-1)^n}{2a}$$

Adding the jets

$$j_{\text{ext}}^0(x, t) = g[\delta(\Delta x - \Delta t) - \delta(\Delta x + \Delta t)]\theta(\Delta t)$$

$$j_{\text{ext}}^1(x, t) = g[\delta(\Delta x - \Delta t) + \delta(\Delta x + \Delta t)]\theta(\Delta t)$$

$$\Delta x \equiv x - x_0$$

$$\Delta t \equiv t - t_0$$

$$H = \int dx \left[\bar{\psi}(-i\gamma^1 \partial_1 + g\gamma^1 A_1 + m)\psi + \frac{1}{2}E^2 + j_{\text{ext}}^1(x, t)A_1 \right]$$



$$H^L(t) = \frac{1}{4a} \sum_{n=1}^{N-1} (X_n X_{n+1} + Y_n Y_{n+1}) + \frac{m}{2} \sum_{n=1}^N (-1)^n Z_n$$

$$+ \frac{ag^2}{2} \sum_{n=1}^{N-1} (L_{\text{dyn},n} + L_{\text{ext},n}(t))^2.$$

$$L_{\text{dyn},n} = \sum_{i=1}^n q_i$$

$$L_{\text{ext},n}(t) = -\theta \left(t - t_0 - \left| x - x_0 + \frac{a}{2} \right| \right)$$

Numerical procedure


Start from the ground state
of the Hamiltonian:

$$H(t = 0)|\Psi(t = 0)\rangle = E_0|\Psi(t = 0)\rangle$$

Switch on the external source
and time evolve:


$$|\Psi_t\rangle = \mathcal{T}e^{-i\int_0^t dt' H(t')} |\Psi_0\rangle$$

$t/a = 1.00$



Numerical time evolution using
classical exact diagonalization or
tensor networks mimics
simulation on a **quantum** device

Screening, chiral condensate and entanglement



$$q_{n,t} = \langle \psi^\dagger(an) \psi(an) \rangle_t = \frac{\langle Z_n \rangle_t + (-1)^n}{2a}$$


$$L_n = L_{\text{dyn},n} + L_{\text{ext},n}(t)$$

$$L_{\text{dyn},n} = \sum_{i=1}^n q_i$$

$$L_{\text{ext},n}(t) = -\theta \left(\frac{t-t_0}{a} - \left| n - \frac{N}{2} \right| \right)$$

$$\nu_{n,t} = \langle \bar{\psi}(an) \psi(an) \rangle_t = \frac{(-1)^n \langle Z_n \rangle_t}{2a}$$

Screening, chiral condensate and entanglement




Electric energy

Chiral condensate

Entanglement entropy



Total charge



Screening, chiral condensate and entanglement

Screening the electric field

Effects of the dynamical pair production:





Screening, chiral condensate and entanglement

Effects of the dynamical pair production:

Screening the electric field

Destroying vacuum condensate



Screening, chiral condensate and entanglement


Effects of the dynamical pair production:

$$\rho_A = \text{Tr}_B \rho, \quad S_{EE} = -\text{Tr}_A(\rho_A \log \rho_A)$$


Screening the electric field

Destroying vacuum condensate

Entangling the jets




Towards thermalization




Towards thermalization

Compare to exact diagonalization



Tensor network methods allow studying much larger system



Towards thermalization



Equilibration towards late times

Towards thermalization

Averaging over the central part



Equilibration towards late times

Thermal expectation values

For any operator

$$\langle \mathcal{O} \rangle_T = \frac{\sum_n e^{-E_n/T} \langle E_n | \mathcal{O} | E_n \rangle}{\sum_n e^{-E_n/T}}$$


where

$$H |E_n\rangle = E_n |E_n\rangle$$


Access the whole spectrum with exact diagonalization

Can also access Gibbs entropy:


$$S = - \sum_n p_n \log p_n, \quad p_n = e^{-E_n/T}$$



Thermalization dynamics



Renyi entropy of the central region




$$S_2(L) = -\log \text{Tr}(\rho_A^2)$$

Study as a function of L


Ground state: “area law” (L -independent)

Typical state, e.g. thermal: “volume law” (linear in L)

E. Bianchi, L. Hackl, M. Kieburg, M. Rigol, and L. Vidmar,
PRX Quantum **3** (2022)




Area and volume laws of entanglement



Adjust by the jet arrival time

area law at early times

Area and volume laws of entanglement



Adjust by the jet arrival time





area law at early times

Rescale by the subsystem size



volume law at late times

Entanglement spectrum



Schmidt decomposition:

$$|\Psi(t)\rangle = \sum_{i=1}^{2^{N/2}} \sqrt{\lambda_i(t)} |\psi_i^L(t)\rangle \otimes |\psi_i^R(t)\rangle$$

$$\rho_L(t) = \sum_{i=1}^{2^{N/2}} \lambda_i(t) |\psi_i^L(t)\rangle \langle \psi_i^L(t)|$$

$$S_{EE}(t) = - \sum_{i=1}^{2^{N/2}} \lambda_i \ln \lambda_i$$

Symmetry-resolved:

$$\sum_{n=1}^{N/2} q_n |\psi_i^L\rangle \equiv Q_L |\psi_i^L\rangle$$

Renyi entropies and entanglement


$$S_\alpha(t) \equiv \frac{\ln \text{Tr}_L(\rho_L(t)^\alpha)}{1 - \alpha} = \frac{\ln \sum_{i=1}^{2^{N/2}} \lambda_i^\alpha}{1 - \alpha}$$

$$\mathcal{E} \equiv \frac{1 - \text{tr} \rho_L^2}{1 - 2^{-N/2}} = \frac{1 - \sum_{i=1}^{2^{N/2}} \lambda_i^2}{1 - 2^{-N/2}}$$

Differentiate between pure state (PS)
and maximally entangled state (MES):

$$S_\alpha[\text{PS}] = 0, \quad \mathcal{E}[\text{PS}] = 0$$


$$S_\alpha[\text{MES}] = \frac{N \ln 2}{2} \forall \alpha, \quad \mathcal{E}[\text{MES}] = 1$$



Fermionic Fock (computational) basis

$$|\mathcal{N}\rangle = |1010\dots 10\rangle$$

Neel state




N=10 example

Fermionic Fock (computational) basis

$$|\mathcal{N}\rangle = |1010\dots 10\rangle$$


Neel state



N=10 example

$$|ij\rangle = X_i X_j |\mathcal{N}\rangle$$


1-pair excitation



$|36\rangle$ example


...

Hadronization in real time



Full state overlap with one-pair states


$$p_{i,j} = |\langle \Psi_t | ij \rangle|^2$$



Full state overlap with one-pair states

$$p_{i,j} = |\langle \Psi_t | ij \rangle|^2$$

$t/a=3$ Nearest neighbor pair - meson




Full state overlap with one-pair states

$$p_{i,j} = |\langle \Psi_t | ij \rangle|^2$$

$t/a=3$ Nearest neighbor pair - meson

$t/a=6$ - Two mesons




Full state overlap with one-pair states

$$p_{i,j} = |\langle \Psi_t | ij \rangle|^2$$

$t/a=3$ Nearest neighbor pair - meson

$t/a=6$ - Two mesons

$t/a=8$ - Three mesons



Full state overlap with one-pair states

$$p_{i,j} = |\langle \Psi_t | ij \rangle|^2$$


$t/a=3$ Nearest neighbor pair - meson

$t/a=6$ - Two mesons

$t/a=8$ - Three mesons

.....

Thermal gas of hadrons?



Conclusion

- Dynamical pair production leads to electric field screening and modification of the vacuum condensate
- Electric field and chiral condensate equilibrate in the central region
- Second Renyi entropy in the central region exhibits a transition from the area law to the volume law
- Entanglement between jets steadily grows with contributions from many Schmidt states
- At large coupling we observe a dynamical transition of Schmidt states from fermionic Fock states to bosonic Fock states