


Energy Independence of the Collins Asymmetry in pp Collisions at STAR

Yixin Zhang ,
for the STAR Collaboration
Shandong University
May.19 ~ 23



Supported in part by




U.S. DEPARTMENT OF
ENERGY

Office of
Science

Challenges in Transverse Single-Spin Asymmetry (TSSA)



- Anomalously large A_N in pp collisions observed for decades




$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow}$$

- LO QCD predicts $A_N \sim 0$

G. Kane, J. Pumplin, W. Repko, Phys. Rev. Lett 41,1689 (1978).

- Left-right asymmetries of different experiments at different beam energies



E. C. Aschenauer et al. arXiv:1602.03922

- Stable in different C.M. energies
- Interpreted by the twist-3 and transverse-momentum-dependent (TMD) formalisms

$$x_F = \frac{2p_z}{\sqrt{s}}$$


Mechanisms for Transverse Single-Spin Asymmetry

➤ Transverse Momentum Dependent (TMD) parton distributions and fragmentation functions.

- Need two scales (Q and p_T), $Q \gg p_T$

✓ **Sivers effect (*Sivers*'90):**


Nucleon spin and parton $k_\perp$ correlation in initial state (related to orbital angular momentum)



$$\vec{S} \cdot (\vec{p} \times \vec{k}_\perp)$$

✓ **Collins effect (*Collins*'93):**

Quark spin and hadron $k_\perp$ correlation in fragmentation process (coupled with transversity)




➤ Twist-3 mechanism (*Efremov-Teryaev*'82, *Qiu-Sterman*'91):

- Collinear/twist-3 multi-parton correlation + fragmentation functions
- Need one scale (Q or p_T), $Q, p_T \gg \Lambda_{QCD}$
- Consistent with TMD mechanism in the overlapping kinematics region

Relativistic Heavy Ion Collider (RHIC)




RHIC- world's first polarized proton-proton collider



The Solenoidal Tracker At RHIC (STAR)



- Subsystems used in this talk



- Time Projection Chamber (TPC)
 - $|\eta| < 1$ and $\phi \in [0, 2\pi]$
 - Main detector for tracking and PID
- Time Of Flight (TOF)
 - $|\eta| < 1.0$ and $\phi \in [0, 2\pi]$
 - Improve PID of tracks
- ElectroMagnetic Calorimeter
 - BEMC: $|\eta| < 1.0$ and $\phi \in [0, 2\pi]$
 - EEMC: $1.08 < \eta < 2.0$ and $\phi \in [0, 2\pi]$
 - Reconstruction of photon, e , π^0 and triggering
- Forward Meson Spectrometer (FMS)
 - $2.6 < \eta < 4.2, \phi \in [0, 2\pi]$
 - Detect γ , π^0 , η

TSSA of pp Collisions

- Transversely polarized proton–proton collision data in recent years at STAR


Year	2011	2012	2015	2017	2022	2024
$\sqrt{s}$ (GeV)	500	200	200	510	508	200
L_{int} (pb^{-1})	25	14	52	350	400	~170
Polarization	53%	57%	57%	55%	52%	53% / 57%

- Measurements at RHIC can reach values of Q^2 that are more than two orders of magnitude higher than current SIDIS experiments

- Collins effect for hadron within jet

- Separate initial and final state effects
- Jet- $p_T \sim$ hard scale; hadron $p_T \sim$ soft scale
- Validate universality with SIDIS and e^+e^- annihilation

F. Yuan, Phys. Rev. Lett. 100, 032003 (2008)



Angle Modulations of TSSA in pp Collisions




- For $\pi^\pm$ within jets in pp collisions, the spin dependent cross section :

$$\frac{d\sigma^\uparrow(\phi_S, \phi_H) - d\sigma^\downarrow(\phi_S, \phi_H)}{d\sigma^\uparrow(\phi_S, \phi_H) + d\sigma^\downarrow(\phi_S, \phi_H)} \propto \begin{aligned} & A_{UT}^{\sin(\phi_S)} \sin(\phi_S) && \text{related to Sivers effect} \\ & + A_{UT}^{\sin(\phi_S - \phi_H)} \sin(\phi_S - \phi_H) && \text{related to Collins effect} \\ & + A_{UT}^{\sin(\phi_S - 2\phi_H)} \sin(\phi_S - 2\phi_H) && \text{related to Collins-like effect} \\ & + A_{UT}^{\sin(\phi_S + \phi_H)} \sin(\phi_S + \phi_H) \\ & + A_{UT}^{\sin(\phi_S + 2\phi_H)} \sin(\phi_S + 2\phi_H) \end{aligned}$$

U. D'Alesio, F. Murgia, and C. Pisano, Phys. Rev. D 83,034021 (2011)



- ϕ_S : azimuthal angle between the proton transverse spin polarization vector and jet scattering plane.
- ϕ_H : azimuthal angle of pion relative to the jet scattering plane.



STAR, Phys. Rev. D 97, 032004 (2018)

Particle Identification

- Particle identification with TOF unmatched (left) and matched (right)



$$n\sigma_{dE/dx} = \frac{1}{\sigma_{\text{exp}}} \ln \left(\frac{dE/dx_{\text{meas}}}{dE/dx_{\text{theo}}} \right)$$

$$n\sigma_{TOF} = \frac{TOF_{\text{meas}} - \frac{L}{c\beta(p)}}{\sigma_{\text{eff}}}$$


- Determine particle rich region

- Asymmetries purification through Moore–Penrose inverse.

$$\begin{pmatrix} f_{\pi \text{ rich}}^{\pi TOF} & f_{\pi \text{ rich}}^{K TOF} & f_{\pi \text{ rich}}^{p TOF} \\ f_{K \text{ rich}}^{\pi TOF} & f_{K \text{ rich}}^{K TOF} & f_{K \text{ rich}}^{p TOF} \\ f_{p \text{ rich}}^{\pi TOF} & f_{p \text{ rich}}^{K TOF} & f_{p \text{ rich}}^{p TOF} \\ f_{\pi \text{ rich}}^{\pi dE/dx} & f_{\pi \text{ rich}}^{K dE/dx} & f_{\pi \text{ rich}}^{p dE/dx} \\ f_{K \text{ rich}}^{\pi dE/dx} & f_{K \text{ rich}}^{K dE/dx} & f_{K \text{ rich}}^{p dE/dx} \\ f_{p \text{ rich}}^{\pi dE/dx} & f_{p \text{ rich}}^{K dE/dx} & f_{p \text{ rich}}^{p dE/dx} \end{pmatrix} \begin{pmatrix} A_{\pi \text{ pure}} \\ A_{K \text{ pure}} \\ A_{p \text{ pure}} \end{pmatrix} = \begin{pmatrix} A_{\pi \text{ raw}}^{TOF} \\ A_{K \text{ raw}}^{TOF} \\ A_{p \text{ raw}}^{TOF} \\ A_{\pi \text{ raw}}^{dE/dx} \\ A_{K \text{ raw}}^{dE/dx} \\ A_{p \text{ raw}}^{dE/dx} \end{pmatrix}$$



- $f_{i \text{ rich}}^j$: the fraction of particle type j in the i -rich sample.
- Extract the pure asymmetry for each particle rich region

Sivers Asymmetry of Inclusive Jets at 200 GeV & 510 GeV



- Sensitive to twist-3 correlator associated with the gluon Sivers function
- High-precision measurements show that the inclusive jet asymmetry is very small.

Sivers Asymmetry of Hadron-Tagged Jet at 200 GeV & 510 GeV






- Quark jet fractions are enhanced by tagging $\pi^\pm$
- The experimental measurements are comparable to the theoretical predictions.

Z. B. Kang, A. Prokudin, F. Ringer and F. Yuan, Phys. Lett. B 774 (2017), 635–642

Sivers Asymmetry of weak bosons

➤ Preliminary results of W/Z A_N from run17 data at $\sqrt{s} = 510$ GeV



STAR, Phys. Lett. B 854 (2024) 138715

- Test sign change of Sivers function from SIDIS and Drell–Yan:

$$f_{h/q^\uparrow}^{\text{SIDIS}}(x, k_T, Q^2) = -f_{h/q^\uparrow}^{p+p \rightarrow DY}(x, k_T, Q^2)$$


- In general, the results and theoretical calculations are consistent
- Provide input to extraction of the Sivers function, especially for valence quarks at high x ($x \geq 0.1$)
- Run 22 data will further reduce the statistical uncertainty and push to larger rapidity $y^{W/Z}$

Collins Asymmetry at 510/500 GeV



➤ Collins results as a function of jet p_T

- Positive for π^+ and negative for π^- , and increase with increasing jet p_T for $x_F > 0$
- The asymmetries for $x_F < 0$ are consistent with 0.





Collins Asymmetry at 510/500 GeV



➤ Collins results as a function of jet p_T

- Positive for π^+ and negative for π^- , and increase with increasing jet p_T for $x_F > 0$
- The asymmetries for $x_F < 0$ are consistent with 0.




- New results are consistent with previous run11 data, but with 13 times more statistics

Collins Asymmetry at 510/200 GeV: Test TMD Evolution



- Comparison of Collins asymmetry *vs.* jet- x_T




- The high precision Collins results of 510 GeV and 200 GeV nicely align with jet x_T scale, giving almost no energy dependence
- At the same jet x_T , 200 GeV and 500 GeV results correspond to the same x in transversity distribution

Collins Asymmetry at 510/200 GeV: Test TMD Evolution



➤ Comparison of Collins asymmetry *vs.* jet- x_T



L. Gamberg, M. Malda, J. A. Miller, D. Pitonyak, A. Prokudin, N. Sato, [JAM], Phys. Rev. D 106 (2022), 034014


U. D'Alesio, C. Flore and M. Zacccheddu, work in preparation

- Model calculations without TMD evolution are generally consistent with measurements

Collins Asymmetry at 510/200 GeV: Test TMD Evolution




➤ Comparison of Collins asymmetry *vs.* hadron j_T



- Similar jet x_T for 200 GeV and 510 GeV, allowing for comparisons of measurements at similar parton momentum fraction regions
- Again, Collins results of two energies align with pion j_T scale, no significant energy dependence observed

j_T : pion's transverse momentum relative to jet axis



Collins Asymmetry at 510/200 GeV: Test TMD Evolution





- The Collins asymmetries as a function of z also show good consistency

Theoretical Calculations with/without TMD Evolution

Without TMD evolution:



With TMD evolution:



L. Gamberg, M. Malda, J. A. Miller, D. Pitonyak, A. Prokudin, N. Sato, [JAM], Phys. Rev. D 106 (2022), 034014

U. D'Alesio, C. Flore and M. Zaccheddu, work in preparation


Z. B. Kang, A. Prokudin, F. Ringer and F. Yuan, Phys. Lett. B 774 (2017), 635–642

C. Zeng, H. Dong, T. Lieve, P. Sun and Y. Zhao, arXiv:2412.18324

- All models are consistent with our data in general, with weak energy dependence

Collins Asymmetry of K & p

STAR, Phys. Rev. D 106, 072010 (2022)




- K^+ has a contribution from favored fragmentation of u quarks, similar in magnitude to those for π^+
- K^- can only come from unfavored fragmentation, are consistent with zero within uncertainties
- Collins asymmetry of proton is consistent zero

Collins-like Asymmetry of $\pi^\pm$




➤ Results at 200GeV, as a function of z



STAR, Phys. Rev. D 106, 072010 (2022)

- Sensitive to linearly polarized gluons inside a transversely polarized proton
- No significant asymmetry for either collision energy

➤ Results at 500GeV, as a function of z




STAR, Phys. Rev. D 97, 032004 (2018),

π^0 Asymmetry in Forward Region




STAR, Phys. Rev. D 103, 092009 (2021)




- Sizeable A_N for forward π^0

➤ Sivers effect contribution




- Magnitude of inclusive EM jet asymmetry is limited

➤ Collins effect contribution




- Collins asymmetry of π^0 inside EM jet is consistent with zero



- The A_N of the isolated π^0 was found to be significantly larger than that for non-isolated ones
- Indicating the possible contribution from the **diffractive process**


Diffractive Electromagnetic Jets Asymmetry

➤ Single diffractive process




- $p^\uparrow + p \rightarrow p + \text{EM-jet} + X$
- 1 east roman pots (RP) track , no requirement on west RP
- 1 EM-jet per event is allowed

➤ Rapidity gap event



- $p^\uparrow + p \rightarrow p + p + \text{EM-jet} + X$
- No RP track requirement
- 1 EM-jet per event is allowed





- The single diffractive and rapidity gap EM-jet A_N are consistent within uncertainty
- The observed inclusive π^0 asymmetry is unlikely to originate from diffractive processes

Outlook

➤ STAR Forward Detector Upgrade ($2.5 < \eta < 4$)

Tracking system & calorimetry system



- Successfully collected pp data at 510 and 200GeV during 2022 and 2024
- x extends up to ~ 0.5 , with charged particle tracking and electromagnetic and hadronic calorimetry
- Complement the kinematic coverage of ep data from the EIC

Year	2022	2024
$\sqrt{s}$ (GeV)	510	200
L_{int} (pb^{-1})	400	~ 170
Polarization	52%	53% / 57%

Summary




- The high-precision Collins asymmetries for π^+ and π^- results at 510 GeV, in good agreement with 200 GeV data, indicating a very weak energy dependence of the Collins effect
- No significant Sivers asymmetry or Collins-like asymmetry observed in pp collision at mid-rapidity
- W/Z boson asymmetries can provide important input to extraction of the Sivers function
- These measurements, together with pp data after STAR forward upgrade, will provide valuable insights into the three-dimensional structure of the proton

Thank you for the attention!

Back up

Jet Reconstruction



➤ Jet reconstruction :

- Anti- K_T algorithm with $R = 0.5$
- TPC tracks and EMC energy deposition as input
- Off-axis cone method to estimate underlying event contribution

➤ Simulation

- PYTHIA 6.4 with STAR adjustment of Perugia 2012
- Kinematic correction & Systematic uncertainty estimation

Extraction of Transverse Single-Spin Asymmetries




- Cross-ratio method to extract the asymmetries of different modulations.

$$A_N \sin(\phi) = \frac{1}{P} \cdot \frac{\sqrt{N^\uparrow(\phi)N^\downarrow(\phi + \pi)} - \sqrt{N^\downarrow(\phi)N^\uparrow(\phi + \pi)}}{\sqrt{N^\uparrow(\phi)N^\downarrow(\phi + \pi)} + \sqrt{N^\downarrow(\phi)N^\uparrow(\phi + \pi)}}$$

- Cross ratio method can cancel detector efficiencies and spin dependent luminosity.
- $N^\uparrow$ (or $N^\downarrow$) is the yield for a given spin state.

Collins Asymmetry from STAR 2017 Data

- Collins results as a function of z in different jet p_T regions at 510 GeV:




z : the pion's longitudinal momentum fraction in the jet

- These results provide more detailed constraints on the Collins fragmentation function

Collins Asymmetry from STAR 2017 Data

- Collins results as a function of j_T in different jet p_T regions at 510 GeV:



j_T : charged pion's transverse momentum relative to the jet axis

- These results provide more detailed constraints on the Collins fragmentation function